Monte Carlo in Action

Prof. Americo Cunha Jr

Rio de Janeiro State University - UERJ

americo.cunha@uerj.br

www.americocunha.org













Unit square sampling

Sample at random an unit square according to the distributions:

- Uniform
- Normal
- Gamma
- Chi-squared



Matlab code for square sampling (1/2)

```
clc; clear; close all;
    Ns = 1000; xmin = 0.0; xmax = 1.0; ymin = 0.0; ymax = 1.0;
    mu = 0.5; coefvar = 0.5; sigma = mu*coefvar;
6
    % uniform sample
    Xu = rand(Ns); Yu = rand(Ns);
9
    % normal sample
    Xn = mu + sigma*randn(Ns); Yn = mu + sigma*randn(Ns);
    % gamma sample
14
    Xg = gamrnd(1/coefvar^2, mu*coefvar^2, Ns);
    Yg = gamrnd(1/coefvar^2, mu*coefvar^2, Ns);
16
    % Chi-squared sample
    Xc2 = chi2rnd(mu,Ns); Yc2 = chi2rnd(mu,Ns);
```




Matlab code for square sampling (2/2)

```
figure(1)
    plot(Xu,Yu,'.b');
    axis([xmin xmax ymin ymax]);
    title(' Uniform Distribution', 'FontSize', 20)
6
    figure(2)
    plot(Xn,Yn,'.r');
    axis([xmin xmax ymin ymax]);
    title(' Normal Distribution', 'FontSize', 20)
    figure(3)
    plot(Xg,Yg,'.g');
    axis([xmin xmax ymin ymax]);
    title(' Gamma Distribution'. 'FontSize'.20)
14
15
16
    figure (4)
    plot(Xc2,Yc2,'.m');
    axis([xmin xmax ymin ymax]);
18
19
    title(' Chi-squared Distribution', 'FontSize', 20)
```



Different samplings of an unit square





MC for integration

Consider the integral

$$\alpha = \int_0^{2\pi} \sin^2(x) dx$$

$$= \int_0^{2\pi} 2\pi \sin^2(x) \frac{1}{2\pi} dx$$

$$= \int_{\mathbb{R}} h(y) p(y) dy$$

where

$$h(y) = 2\pi \sin^2(y)$$

•
$$p(y) = \frac{1}{2\pi} \mathbb{1}_{(0,2\pi)}(y)$$
 i.e., $Y \sim \mathcal{U}(0,2\pi)$.




MC for integration

```
clc: clear: close all
    Ns = 1e5;
    ymin = 0.0; ymax = 2*pi;
8
    Y = ymin + (ymax-ymin)*rand(Ns,1);
    hY = (vmax - vmin) * (sin(Y).^2):
    alpha hat = cumsum(hY)./(1:Ns)':
               = abs(pi-alpha_hat);
    error
14
    semilogy(1:Ns,error,'x')
15
    title(' Monte Carlo integration', 'FontSize', 20)
16
    set(gca, 'fontsize',18)
    xlabel('number of samples')
    ylabel('error')
19
    grid
```





MC for integration





MC for estimation of statistics







(X,Y,Z) is a random vector defined on the unit cube in \mathbb{R}^3



MC for estimation of statistics







(X, Y, Z) is a random vector defined on the unit cube in \mathbb{R}^3

$$\mathcal{P}\left\{ (X, Y, Z) \in x^2 + y^2 + z^3 \le 1/4 \right\} = \frac{1}{6}\pi$$



MC for estimation of statistics





(X,Y,Z) is a random vector defined on the unit cube in \mathbb{R}^3

$$\mathcal{P}\left\{ (X,Y,Z) \in x^2 + y^2 + z^3 \le 1/4 \right\} = \frac{1}{6}\pi$$

$$\mathcal{P}\left\{ \left(X,Y,Z\right)\in x+y+z=1\right\} =0$$



MC for estimation of statistics (sphere)

```
clear: close all: clc
    % sample size
    Ns = 1000000:
    % draw the samples
    X = -0.5 + rand(Ns.3):
8
9
    % distance from the origin for each sample
    R = sqrt(sum(X.^2,2));
    % count the number of hits
    count = sum(R \le 1/2)
14
    % compute the estimated volume
    P_sphere = count/Ns
16
    P true = pi/6
18
    rel_error = (abs(P_sphere-P_true)/abs(P_true))*100
```



MC for estimation of statistics (sphere)



MC for estimation of statistics (plane)

```
clear; close all; clc

% sample size
Ns = 100000;

% draw the samples
X = -0.5 + rand(Ns,3);

% sum of the coordinates
SumCoord = sum(X,2);

% count the number of hits
count = sum(SumCoord == 1)

% compute the estimated volume
P_plane = count/Ns
P_true = 0
```




MC for estimation of statistics (plane)

P_true =

0



MC for uncertainty propagation



Parameters support:

$$(r,h) \in [0,1] \times (0,\infty)$$

Volume:

$$v(r,h) = \frac{1}{3}\pi r^2 h$$

Input joint-distribution:

 $R \sim Beta(a,b)$ and $H \sim Exp(1/\mu_H)$ independent

$$p_{RH}(r,h) = \mathbb{1}_{[0,1]}(r) \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} r^{a-1} (1-r)^{b-1} \times \mathbb{1}_{(0,\infty)}(h) \frac{\exp(-h/\mu_H)}{\mu_H}$$

• Output distribution:


$$p_V(v) = ???$$



MC for uncertainty propagation

```
clear; close all; clc
    % random number generator (fix the seed for reproducibility)
    rng_stream = RandStream('mt19937ar', 'Seed', 30081984);
    RandStream.setGlobalStream(rng_stream);
    Ns = 5000:
6
    muR = 0.5; sigmaR = 0.1; nuR = muR*(1-muR)/(sigmaR^2)-1;
    a = muR*nuR; b = (1-muR)*nuR; muH = 1.0;
9
    R = betarnd(a.b.Ns.1): H = exprnd(muH.Ns.1):
    V_{cone}
           = (1/3)*pi*R.^2.*H;
    [bins.freq] = randvar pdf(V cone.round(sqrt(Ns)));
    alpha 2 = cumsum(V cone.^2)'./(1:Ns);
14
    figure(1)
    gplotmatrix([R,H]);
16
18
    figure(2)
19
    plot(V cone. 'o'): vlim([0 5])
20
    figure(3)
    bar(bins.freg):
    xlim([0 4]); ylim([0 5])
24
25
    figure (4)
26
    plot(1:Ns.alpha 2. 'LineWidth'.2)
```

MC for uncertainty propagation






MC convergence analysis (finite mean)

```
clc; clear; close all
    Ns = 10^3:
    X1 = randn(Ns, 1); X2 = randn(Ns, 1); X3 = randn(Ns, 1);
    Y1 = 1./(1+X1.^2); Y2 = 1./(1+X2.^2); Y3 = 1./(1+X3.^2);
6
    meanY1 = mean(Y1)
    mean Y2 = mean (Y2)
9
    meanY3 = mean(Y3)
    figure(1)
    plot(1:Ns,X1, 'xb',1:Ns,X2, 'xg',1:Ns,X3, 'xm', 'LineWidth',1)
    set(gca, 'fontsize', 18)
14
    xlabel('index'); ylabel('X'); xlim([1,Ns]); ylim([-4 4]);
16
    figure(2)
    plot(1:Ns,Y1,'xb',1:Ns,Y2,'xg',1:Ns,Y3,'xm','LineWidth',1)
18
    hold on
    plot([1 Ns],[meanY1 meanY1], 'b-', 'LineWidth', 4);
19
20
    plot([1 Ns],[meanY2 meanY2], 'g-', 'LineWidth', 4);
    plot([1 Ns], [meanY3 meanY3], 'm-', 'LineWidth', 4);
    hold off
    xlabel('index'); ylabel('Y'); xlim([1,Ns]); ylim([0 1]);
```



MC convergence analysis (finite mean)






MC convergence analysis (infinite mean)

```
clc; clear; close all;
    Ns = 10^3:
    X1 = randn(Ns,1); X2 = randn(Ns,1); X3 = randn(Ns,1);
    Z1 = 1./(1+X1); Z2 = 1./(1+X2); Z3 = 1./(1+X3);
6
    meanZ1 = mean(Z1)
    mean 72 = mean (72)
9
    mean Z3 = mean (Z3)
    figure(1)
    plot(1:Ns,X1,'xb',1:Ns,X2,'xg',1:Ns,X3,'xm','LineWidth',1)
    xlabel('index'): vlabel('X'): xlim([1.Ns]): vlim([-4.4]):
14
    figure(2)
    plot(1:Ns.Z1, 'xb'.1:Ns.Z2, 'xg'.1:Ns.Z3, 'xm', 'LineWidth'.1)
16
    hold on
18
    plot([1 Ns],[meanZ1 meanZ1],'b-', 'LineWidth', 4);
    plot([1 Ns],[meanZ2 meanZ2], 'g-', 'LineWidth', 4);
19
    plot([1 Ns],[meanZ3 meanZ3],'m-', 'LineWidth', 4);
    hold off
    xlabel('index'); ylabel('Y'); xlim([1,Ns]); ylim([-2000 12000]);
```



MC convergence analysis (infinite mean)





References



S. B. Volchan, What Is a Random Sequence? The American Mathematical Monthly, 109: 46-63, 2002.



T. Homem-de-Mello and G. Bayraksan, *Monte Carlo sampling-based methods for stochastic optimization*, **Surveys in Operations Research and Management Science**, 19: 56–85, 2014.



A. Cunha Jr, R. Sampaio, R. Nasser, H. Lopes, and K. Breitman, *Uncertainty quantification through Monte Carlo method in a cloud computing setting*, Computer Physics Communications, 185: 1355-1363, 2014.



R. W. Shonkwiler, and F. Mendivil, Explorations in Monte Carlo Methods, Springer, 2009.



C. Robert, and G. Casella, Monte Carlo Statistical Methods, Springer, 2nd edition, 2005.



S. Asmussen, and P. W. Glynn, Stochastic Simulation: Algorithms and Analysis, Springer, 2007.



J. E. Gentle, Random Number Generation and Monte Carlo Methods, Springer, 2nd Edition, 2004.



J. S. Liu, Monte Carlo Strategies in Scientific Computing, Springer, 2009.



R. Y. Rubinstein, and Dirk P. Kroese, Simulation and the Monte Carlo Method, Wiley, 3rd Edition, 2016.



D. P. Kroese, T. Taimre, and Z. I. Botev, Handbook of Monte Carlo Methods, Wiley, 2011.


















How to cite this material?

A. Cunha Jr, *Monte Carlo in Action*, Rio de Janeiro State University – UERJ, 2021.











These class notes may be shared under the terms of Creative Commons BY-NC-ND 4.0 license, for educational purposes only.



