

# Probabilistic Modeling of Uncertainties in Physical Systems

Prof. Americo Cunha Jr

Rio de Janeiro State University – UERJ

[americo.cunha@uerj.br](mailto:americo.cunha@uerj.br)

[www.americocunha.org](http://www.americocunha.org)



@AmericoCunhaJr



@AmericoCunhaJr




@AmericoCunhaJr



@AmericoCunhaJr



# General framework for uncertainty quantification (UQ)



1. **Stochastic Modeling**: characterize inputs uncertainties
2. **Uncertainty Propagation**: quantify output uncertainties
3. **Certification**: establish acceptable levels of uncertainty
4. **Sensitivity Analysis**: explain the output variability



B. Sudret *A short review of computational methods for uncertainty quantification in engineering*, 2013.

# How to model uncertainties in physical systems?

Several approaches are available:

- Probability theory
- Evidency theory
- Interval analysis
- Fuzzy logic



# How to model uncertainties in physical systems?

Several approaches are available:

- Probability theory
- Evidency theory
- Interval analysis
- Fuzzy logic



# How to model uncertainties in physical systems?

Several approaches are available:

- Probability theory
- Evidency theory
- Interval analysis
- Fuzzy logic

Probability theory is the most used approach to describe uncertainties in physical systems (best suited for high-dimensions)



# How to model uncertainties in physical systems?

Several approaches are available:

- Probability theory
- Evidency theory
- Interval analysis
- Fuzzy logic

Probability theory is the most used approach to describe uncertainties in physical systems (best suited for high-dimensions)

In general other approaches are used only when the probability theory can not be applied




# Different probabilistic approaches

- Parametric probabilistic approach: model parameters as random objects (deal with data uncertainties)
- Nonparametric probabilistic approach: model operators as random operators (deal with data and model uncertainties)

In any approach the probability distribution of the random objects must be constructed, and not arbitrarily chosen.



# A simple mechanical system



Parameter:  $k$  – spring stiffness

Input:  $f$  – external force

Response:  $u$  – displacement

- Mathematical model:

$$k u = f$$

- Model response:

$$u = \underbrace{k^{-1} f}_{g(k)}$$

(nonlinear mapping of  $k$ )




C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.





# A simple mechanical system



Parameter:  $k$  – spring stiffness

Input:  $f$  – external force

Response:  $u$  – displacement

- Mathematical model:

$$k u = f$$

- Model response:

$$u = \underbrace{k^{-1} f}_{g(k)}$$

(nonlinear mapping of  $k$ )

What happens if the model parameter is random?



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# Parametric probabilistic approach

Probability space:  $(\Omega, \Sigma, \mathcal{P})$

Stiffness:

$$K : \Omega \rightarrow \mathbb{R}$$

Displacement:

$$U : \Omega \rightarrow \mathbb{R} \quad \text{such that} \quad K U = f$$

To ensure the consistency of the stochastic model:

- $\mathbb{E} \left\{ K^2 \right\} < +\infty$  “finite variance random variables”
- $\mathbb{E} \left\{ U^2 \right\} < +\infty$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# The importance of know parameters distribution

Hypotheses about random parameter  $K$ :

- finite variance –  $\mathbb{E} \left\{ K^2 \right\} < +\infty$
- known mean –  $\mathbb{E} \left\{ K \right\} = \mu_K$
- unknown distribution –  $p_K(k)$  is not known

Can we compute the model response mean value ?

$$\mathbb{E} \left\{ U \right\} = \mathbb{E} \left\{ K^{-1} f \right\}$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# The importance of know parameters distribution

Hypotheses about random parameter  $K$ :

- finite variance –  $\mathbb{E} \left\{ K^2 \right\} < +\infty$
- known mean –  $\mathbb{E} \left\{ K \right\} = \mu_K$
- unknown distribution –  $p_K(k)$  is not known

Can we compute the model response mean value ?

$$\begin{aligned} \mathbb{E} \{ U \} &= \mathbb{E} \left\{ K^{-1} f \right\} \\ &= \int_{\mathbb{R}} f k^{-1} p_K(k) dk \end{aligned}$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# The importance of know parameters distribution

Hypotheses about random parameter  $K$ :

- finite variance –  $\mathbb{E} \{ K^2 \} < +\infty$
- known mean –  $\mathbb{E} \{ K \} = \mu_K$
- unknown distribution –  $p_K(k)$  is not known

Can we compute the model response mean value ?

$$\begin{aligned} \mathbb{E} \{ U \} &= \mathbb{E} \{ K^{-1} f \} \\ &= \int_{\mathbb{R}} f k^{-1} p_K(k) dk \neq \frac{f}{\mu_K} \quad \text{No, we can't !} \end{aligned}$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# The importance of know parameters distribution

Hypotheses about random parameter  $K$ :

- finite variance –  $\mathbb{E} \{ K^2 \} < +\infty$
- known mean –  $\mathbb{E} \{ K \} = \mu_K$
- unknown distribution –  $p_K(k)$  is not known

Can we compute the model response mean value ?

$$\begin{aligned} \mathbb{E} \{ U \} &= \mathbb{E} \{ K^{-1} f \} \\ &= \int_{\mathbb{R}} f k^{-1} p_K(k) dk \neq \frac{f}{\mu_K} \quad \text{No, we can't !} \end{aligned}$$

**The input distribution is essential to obtain output statistics!**



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# Can we arbitrate distributions?



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# Can we arbitrate distributions?

Suppose  $p_K(k)$  is arbitrarily chosen Gaussian

$$p_K(k) = \frac{1}{\sqrt{2\pi} \sigma_K} \exp \left\{ -\frac{(k - \mu_K)^2}{2 \sigma_K^2} \right\}$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.





# Can we arbitrate distributions?

Suppose  $p_K(k)$  is arbitrarily chosen Gaussian

$$p_K(k) = \frac{1}{\sqrt{2\pi} \sigma_K} \exp \left\{ -\frac{(k - \mu_K)^2}{2 \sigma_K^2} \right\}$$

Does the model response a finite variance random variable ?



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# Can we arbitrate distributions?

Suppose  $p_K(k)$  is arbitrarily chosen Gaussian

$$p_K(k) = \frac{1}{\sqrt{2\pi} \sigma_K} \exp \left\{ -\frac{(k - \mu_K)^2}{2 \sigma_K^2} \right\}$$

Does the model response a finite variance random variable ?

$$\mathbb{E} \left\{ U^2 \right\} = \mathbb{E} \left\{ K^{-2} f^2 \right\}$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# Can we arbitrate distributions?

Suppose  $p_K(k)$  is arbitrarily chosen Gaussian

$$p_K(k) = \frac{1}{\sqrt{2\pi} \sigma_K} \exp \left\{ -\frac{(k - \mu_K)^2}{2 \sigma_K^2} \right\}$$

Does the model response a finite variance random variable ?

$$\begin{aligned} \mathbb{E} \left\{ U^2 \right\} &= \mathbb{E} \left\{ K^{-2} f^2 \right\} \\ &= \int_{-\infty}^{+\infty} f^2 k^{-2} \frac{1}{\sqrt{2\pi} \sigma_K} \exp \left\{ -\frac{(k - \mu_K)^2}{2 \sigma_K^2} \right\} dk \end{aligned}$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# Can we arbitrate distributions?

Suppose  $p_K(k)$  is arbitrarily chosen Gaussian

$$p_K(k) = \frac{1}{\sqrt{2\pi} \sigma_K} \exp \left\{ -\frac{(k - \mu_K)^2}{2 \sigma_K^2} \right\}$$

Does the model response a finite variance random variable ?

$$\begin{aligned} \mathbb{E} \left\{ U^2 \right\} &= \mathbb{E} \left\{ K^{-2} f^2 \right\} \\ &= \int_{-\infty}^{+\infty} f^2 k^{-2} \frac{1}{\sqrt{2\pi} \sigma_K} \exp \left\{ -\frac{(k - \mu_K)^2}{2 \sigma_K^2} \right\} dk \\ &= +\infty \quad \text{No, it doesn't} \end{aligned}$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# Can we arbitrate distributions?

Suppose  $p_K(k)$  is arbitrarily chosen Gaussian

$$p_K(k) = \frac{1}{\sqrt{2\pi} \sigma_K} \exp \left\{ -\frac{(k - \mu_K)^2}{2 \sigma_K^2} \right\}$$

Does the model response a finite variance random variable ?

$$\begin{aligned} \mathbb{E} \left\{ U^2 \right\} &= \mathbb{E} \left\{ K^{-2} f^2 \right\} \\ &= \int_{-\infty}^{+\infty} f^2 k^{-2} \frac{1}{\sqrt{2\pi} \sigma_K} \exp \left\{ -\frac{(k - \mu_K)^2}{2 \sigma_K^2} \right\} dk \\ &= +\infty \quad \text{No, it doesn't} \end{aligned}$$

**And make no sense from the physical point of view!**



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.




# 1st consequence of the unphysical distribution

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 1/2$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$





# 1st consequence of the unphysical distribution

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 1/2$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$





# 1st consequence of the unphysical distribution

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 1/2$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$







# 1st consequence of the unphysical distribution

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 1/2$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



Scenarios that do not respect the second law of thermodynamics may appear! ☹





## 2nd consequence of the unphysical distribution

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 1/2$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 2.3358$$





## 2nd consequence of the unphysical distribution

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 1/2$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 2.6816$$





## 2nd consequence of the unphysical distribution

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 1/2$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 1.4487$$





## 2nd consequence of the unphysical distribution

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 1/2$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 1.4487$$





## 2nd consequence of the unphysical distribution

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 1/2$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 1.4487$$

Statistical estimates don't make sense! ☹





## 2nd consequence of the unphysical distribution

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 1/2$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 1.4487$$

Statistical estimates don't make sense! ☹

What happens if the negative samples could be avoided ?





# Trying to avoid the negative samples

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 3/10$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 2.2514$$







# Trying to avoid the negative samples

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 3/10$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 2.3050$$





# Trying to avoid the negative samples

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 3/10$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 2.2137$$





# Trying to avoid the negative samples

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 3/10$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 2.2137$$

Now the statistical estimates seem to make sense! ☺





# Trying to avoid the negative samples

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 3/10$$

$$K \sim \mathcal{N}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$





$$\widehat{\mu_U} = 2.2137$$

Now the statistical estimates seem to make sense! ☺

**Is this really the case?**





# What is the big consequence of an infinite 2nd moment?



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# What is the big consequence of an infinite 2nd moment?





**Apparently there is nothing wrong !**



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.





# But note the convergence of the estimators



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# But note the convergence of the estimators



**The Monte Carlo simulation does not converge !**





C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.





# But note the convergence of the estimators



**The Monte Carlo simulation does not converge !**

**The obtained response is not statistically significant ! ☹**



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# So we can not arbitrate distributions !

The Gaussian is a bad choice since  $K$  must be a positive-valued random variable.



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# So we can not arbitrate distributions !

The Gaussian is a bad choice since  $K$  must be a positive-valued random variable.

Suppose then:

- positive support –  $\text{Supp } p_K \subset (0, +\infty) \implies K > 0 \text{ a.s.}$
- finite variance –  $\mathbb{E} \{ K^2 \} < +\infty$
- known mean –  $\mathbb{E} \{ K \} = \mu_K$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# So we can not arbitrate distributions !

The Gaussian is a bad choice since  $K$  must be a positive-valued random variable.

Suppose then:

- positive support –  $\text{Supp } p_K \subset (0, +\infty) \implies K > 0 \text{ a.s.}$
- finite variance –  $\mathbb{E} \{ K^2 \} < +\infty$
- known mean –  $\mathbb{E} \{ K \} = \mu_K$

All these requirements are verified by the exponential distribution

$$p_K(k) = \mathbb{1}_{(0,+\infty)}(k) \frac{1}{\mu_K} \exp \left\{ -\frac{k}{\mu_K} \right\}.$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# So we can not arbitrate distributions !

Do we have  $\mathbb{E} \left\{ U^2 \right\} < +\infty$  for the exponential distribution?



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# So we can not arbitrate distributions !

Do we have  $\mathbb{E} \left\{ U^2 \right\} < +\infty$  for the exponential distribution?

$$\mathbb{E} \left\{ U^2 \right\} = \mathbb{E} \left\{ K^{-2} f^2 \right\}$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# So we can not arbitrate distributions !

Do we have  $\mathbb{E} \left\{ U^2 \right\} < +\infty$  for the exponential distribution?

$$\begin{aligned} \mathbb{E} \left\{ U^2 \right\} &= \mathbb{E} \left\{ K^{-2} f^2 \right\} \\ &= \int_0^{+\infty} f^2 k^{-2} \frac{1}{\mu_K} \exp \left\{ -\frac{k}{\mu_K} \right\} dk \end{aligned}$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# So we can not arbitrate distributions !

Do we have  $\mathbb{E} \left\{ U^2 \right\} < +\infty$  for the exponential distribution?

$$\begin{aligned}\mathbb{E} \left\{ U^2 \right\} &= \mathbb{E} \left\{ K^{-2} f^2 \right\} \\ &= \int_0^{+\infty} f^2 k^{-2} \frac{1}{\mu_K} \exp \left\{ -\frac{k}{\mu_K} \right\} dk \\ &= +\infty\end{aligned}$$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.





# So we can not arbitrate distributions !

Do we have  $\mathbb{E} \left\{ U^2 \right\} < +\infty$  for the exponential distribution?

$$\begin{aligned} \mathbb{E} \left\{ U^2 \right\} &= \mathbb{E} \left\{ K^{-2} f^2 \right\} \\ &= \int_0^{+\infty} f^2 k^{-2} \frac{1}{\mu_K} \exp \left\{ -\frac{k}{\mu_K} \right\} dk \\ &= +\infty \end{aligned}$$

The function  $k \mapsto k^{-2}$  diverges in  $k = 0$ .



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# So we can not arbitrate distributions !

Do we have  $\mathbb{E} \left\{ U^2 \right\} < +\infty$  for the exponential distribution?

$$\begin{aligned}\mathbb{E} \left\{ U^2 \right\} &= \mathbb{E} \left\{ K^{-2} f^2 \right\} \\ &= \int_0^{+\infty} f^2 k^{-2} \frac{1}{\mu_K} \exp \left\{ -\frac{k}{\mu_K} \right\} dk \\ &= +\infty\end{aligned}$$

The function  $k \mapsto k^{-2}$  diverges in  $k = 0$ .

**In order to  $\mathbb{E} \left\{ U^2 \right\} < +\infty$  we must have  $\mathbb{E} \left\{ K^{-2} \right\} < +\infty$**



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# An acceptable choice

With the following requirements:

- positive support –  $\text{Supp } p_K \subset (0, +\infty) \implies K > 0 \text{ a.s.}$
- finite variance –  $\mathbb{E} \{ K^2 \} < +\infty$
- known mean –  $\mathbb{E} \{ K \} = \mu_K$
- inverse finite variance –  $\mathbb{E} \{ K^{-2} \} < +\infty$



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# An acceptable choice

With the following requirements:

- positive support –  $\text{Supp } p_K \subset (0, +\infty) \implies K > 0 \text{ a.s.}$
- finite variance –  $\mathbb{E} \{ K^2 \} < +\infty$
- known mean –  $\mathbb{E} \{ K \} = \mu_K$
- inverse finite variance –  $\mathbb{E} \{ K^{-2} \} < +\infty$

$$p_K(k) = \mathbb{1}_{(0,+\infty)}(k) \frac{1}{\mu_K} \frac{\delta_K^{-2\delta_K^{-2}}}{\Gamma(\delta_K^{-2})} \left( \frac{k}{\mu_K} \right)^{\delta_K^{-2}-1} \exp \left\{ -\frac{k/\mu_K}{\delta_K^2} \right\}$$

**The gamma distribution is an acceptable choice !**



C. Soize, *Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering*, Springer, 2017.



# How to safely specify a distribution?

Scenario 1: significant amount of experimental data is available

- Nonparametric statistical estimation

Scenario 2: few or none experimental data is available

- Maximum Entropy Principle  
(a tool from information theory)





# Physically consistent simulation

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 3/10$$

$$K \sim \text{Gamma}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 2.163$$





# Physically consistent simulation

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 3/10$$

$$K \sim \text{Gamma}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 2.2111$$





# Physically consistent simulation

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 3/10$$

$$K \sim \text{Gamma}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 2.1701$$







# Physically consistent simulation

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 3/10$$

$$K \sim \text{Gamma}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$



$$\widehat{\mu_U} = 2.1701$$

Now the statistical estimates seem to make sense! 😊





# Physically consistent simulation

$$f = 2 \quad \mu_K = 1 \quad \sigma_K = 3/10$$

$$K \sim \text{Gamma}(\mu_K, \sigma_K)$$

$$U = K^{-1} f$$





$$\widehat{\mu_U} = 2.1701$$

Now the statistical estimates seem to make sense! 😊



**Is this really the case?**



# Physically consistent simulation





# Physically consistent simulation



**Now Monte Carlo simulation converges !**

# Physically consistent simulation





**Now Monte Carlo simulation converges !**



**The obtained response is statistically significant ! ☺**



# Physically consistent simulation



# Physically consistent simulation



**The gamma (input) distribution is mapped to an inverse-gamma (output) distribution !**



# References



A. Cunha Jr, *Modeling and quantification of physical systems uncertainties in a probabilistic framework*, In: S. Ekwaro-Osire; A. C. Gonçalves; F. M. Alemayehu (Org.), **Probabilistic Prognostics and Health Management of Energy Systems**, Springer, 2017. [http://dx.doi.org/10.1007/978-3-319-55852-3\\_8](http://dx.doi.org/10.1007/978-3-319-55852-3_8)



E. T. Jaynes, **Probability Theory: The Logic of Science**, Cambridge University Press, 2003.



R. C. Smith, **Uncertainty Quantification: Theory, Implementation, and Applications**, SIAM, 2013.








T. J. Sullivan, **Introduction to Uncertainty Quantification**, Springer, 2015.



C. Soize **Uncertainty Quantification: An Accelerated Course with Advanced Applications in Computational Engineering**, Springer, 2017.



R. Ghanem, D. Higdon and H. Owhadi (Editors) **Handbook of Uncertainty Quantification**, Springer, 2017.





## How to cite this material?


A. Cunha Jr, *Probabilistic Modeling of Uncertainties in Physical Systems*, Rio de Janeiro State University – UERJ, 2021.



 @AmericoCunhaJr

 @AmericoCunhaJr

 @AmericoCunhaJr

 @AmericoCunhaJr

These class notes may be shared under the terms of  
Creative Commons BY-NC-ND 4.0 license,  
for educational purposes only.

