

1 Axioms for the natural numbers

Axiom. (Peano Postulates). *There exists a set \mathbb{N} with an element $1 \in \mathbb{N}$ and a function $s : \mathbb{N} \rightarrow \mathbb{N}$ that satisfy the following three properties.*

- a. *There is no $n \in \mathbb{N}$ such that $s(n) = 1$.*
- b. *The function is injective.*
- c. *Let $G \subseteq \mathbb{N}$ be a set. Suppose that $1 \in G$ and that if $g \in G$ then $s(g) \in G$. Then $G = \mathbb{N}$.*

Definition. *The set of **natural numbers**, denoted \mathbb{N} , is the set the existence of which is given in the Peano Postulates.*

Theorem 1. (Definition by Recursion). *Let H be a set, and let $e \in H$ and let $k : H \rightarrow H$ be a function. Then there is a unique function $f : \mathbb{N} \rightarrow H$ such that $f(1) = e$ and that $f \circ s = k \circ f$.*

Theorem 2. *There is a unique binary operation $+$: $\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ that satisfies the following two properties for all $n, m \in \mathbb{N}$.*

- a. $n + 1 = s(n)$
- b. $n + s(m) = s(n + m)$

Theorem 3. *There is a unique binary operation \cdot : $\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ that satisfies the following two properties for all $n, m \in \mathbb{N}$.*

- a. $n \cdot 1 = n$
- b. $n \cdot s(m) = (n \cdot m) + n$

Theorem 4. *Let $a, b, c \in \mathbb{N}$.*

- 1. *If $a + c = b + c$ then $a = b$ (Cancellation Law for Addition).*
- 2. *$(a + b) + c = a + (b + c)$ (Associative Law for Addition).*
- 3. *$1 + a = s(a) = a + 1$.*
- 4. *$a + b = b + a$ (Commutative Law for Addition).*
- 5. *$a + b \neq 1$.*
- 6. *$a + b \neq a$.*
- 7. *$a \cdot 1 = a = 1 \cdot a$ (Identity Law for Multiplication).*
- 8. *$(a + b)c = ac + bc$ (Distributive Law).*
- 9. *$ab = ba$ (Commutative Law for Multiplication).*
- 10. *$(ab)c = a(bc)$ (Associative Law for Multiplication).*
- 11. *If $ac = bc$ then $a = b$ (Cancellation Law for Multiplication).*
- 12. *$ab = 1$ if and only if $a = 1 = b$.*

Definition. The relation $<$ on \mathbb{N} is defined by $a < b$ if and only if there is some $p \in \mathbb{N}$ such that $a + p = b$ for all $a, b \in \mathbb{N}$. The relation \leq on \mathbb{N} is defined by $a \leq b$ if and only if $a < b$ or $a = b$, for all $a, b \in \mathbb{N}$.

Theorem 5. Let $a, b, c, d \in \mathbb{N}$.

1. $a \leq a$ and $a \not< a$, and $a < a + 1$.
2. $1 \leq a$.
3. If $a < b$ and $b < c$, then $a < c$; if $a \leq b$ and $b < c$ then $a < c$; if $a < b$ and $b \leq c$ then $a < c$; if $a \leq b$ and $b \leq c$ then $a \leq c$.
4. $a < b$ if and only if $a + c < b + c$.
5. $a < b$ if and only if $ac < bc$.
6. Precisely one of $a < b$ or $a = b$ or $a > b$ holds (Trichotomy Law).
7. $a \leq b$ or $b \leq a$
8. If $a \leq b$ and $b \leq a$ then $a = b$.
9. It cannot be that $b < a < b + 1$.
10. $a \leq b$ if and only if $a < b + 1$.
11. $a < b$ if and only if $a + 1 \leq b$.

Theorem 6. (Well-Ordering Principle). Let $G \subseteq \mathbb{N}$ be a non-empty set. Then there is some $m \in G$ such that $m \leq g$ for all $g \in G$.

Exercises

1. Fill in the missing details in the proof of Theorem 1.2.6.

Proof. To prove uniqueness, suppose that there are two binary operations \cdot and \times on \mathbb{N} that satisfy the two properties of the theorem. Let

$$G = \{x \in \mathbb{N} \mid n \cdot x = n \times x \text{ for all } n \in \mathbb{N}\}.$$

Then $G \subseteq \mathbb{N}$. By part (a) applied to both \cdot and \times we see that $n \cdot 1 = 1 = n \times 1$ for all $n \in \mathbb{N}$, so $1 \in G$. Now let $q \in G$ and $n \in \mathbb{N}$. Then $n \cdot q = n \times q$. Then it follows from part (b) that

$$n \cdot s(q) = (n \cdot q) + n = (n \times q) + n = n \times s(q).$$

Then $s(q) \in G$, and therefore we can conclude using part (c) of the Peano Postulates that $G = \mathbb{N}$.

Let $q \in \mathbb{N}$. Let $h_q : \mathbb{N} \mapsto \mathbb{N}$ be defined by $h_q(m) = m + q$ for all $m \in \mathbb{N}$. Applying theorem 1.2.4 to the set \mathbb{N} , the element $q \in \mathbb{N}$ and the function $h_q : \mathbb{N} \mapsto \mathbb{N}$ implies that there is a unique function $g_q : \mathbb{N} \mapsto \mathbb{N}$ such that $g_q(1) = q$ and $g_q \circ s = h_q \circ g_q$. Let $\cdot : \mathbb{N} \times \mathbb{N} \mapsto \mathbb{N}$ be defined by $c \cdot d = g_c(d)$ for all $(c, d) \in \mathbb{N} \times \mathbb{N}$. Let $n, m \in \mathbb{N}$. Then $n \cdot 1 = g_n(1) = n$ which is part (a), and $n \cdot s(m) = g_n(s(m)) = (g_n \circ s)(m) = (h_n \circ g_n)(m) = h_n(g_n(m)) = (n \cdot m) + n$ which is part (b). \square