

# Algorithms

## Runtime Analysis And Order Statistics

Dr. Mudassir Shabbir

LUMS University

January 22, 2026



# Selection Problem

## Problem

Given an array  $A$  of  $n$  distinct numbers, find the  $k$ -th smallest element.

- Sorting gives  $O(n \log n)$  time
- Can we do better?
- **Yes:** Deterministic linear-time selection

## Median of Medians Algorithm



# High-Level Idea

- Use a **good pivot** for partitioning
- Pivot should be guaranteed to be “near the middle”
- Avoid worst-case  $O(n^2)$  behavior



# High-Level Idea

- Use a **good pivot** for partitioning
- Pivot should be guaranteed to be “near the middle”
- Avoid worst-case  $O(n^2)$  behavior


## Key Insight

We can find a pivot whose rank is guaranteed to be between  $\frac{3n}{10}$  and  $\frac{7n}{10}$ .



# Step 1: Grouping

- Divide the array into groups of 5
- Each group has at most 5 elements



## Step 2: Medians of Groups

- Sort each group of 5 (constant time per group)
- Extract the median of each group



## Step 2: Medians of Groups

- Sort each group of 5 (constant time per group)
- Extract the median of each group

### Observation

Number of medians =  $\lceil n/5 \rceil$



## Step 2: Medians of Groups

- Sort each group of 5 (constant time per group)
- Extract the median of each group

### Observation

Number of medians =  $\lceil n/5 \rceil$

- Recursively find the median of these medians
- This value is called the **median of medians**





## Step 3: Partitioning

- Use the median of medians as a pivot
- Partition the array around this pivot



## Step 3: Partitioning

- Use the median of medians as a pivot
- Partition the array around this pivot

### Guarantee

At least  $\frac{3n}{10}$  elements are  $\leq$  pivot At least  $\frac{3n}{10}$  elements are  $\geq$  pivot



## Step 3: Partitioning

- Use the median of medians as a pivot
- Partition the array around this pivot

### Guarantee

At least  $\frac{3n}{10}$  elements are  $\leq$  pivot At least  $\frac{3n}{10}$  elements are  $\geq$  pivot

- Recurse only on the relevant side



# Resulting Recurrence

- One recursive call to find median of medians:

$$T(n/5)$$

- One recursive call after partitioning:

$$T(7n/10)$$

- Linear work for grouping, sorting, partitioning



# Resulting Recurrence

- One recursive call to find median of medians:

$$T(n/5)$$

- One recursive call after partitioning:

$$T(7n/10)$$

- Linear work for grouping, sorting, partitioning

## Recurrence

$$T(n) \leq T(n/5) + T(7n/10) + cn$$




# Convex Functions

## Definition

A function  $f$  is **convex** if for all  $a, b$  and  $\alpha \in [0, 1]$ ,

$$f(\alpha a + (1 - \alpha)b) \leq \alpha f(a) + (1 - \alpha)f(b)$$



# A Useful Property

## Assumption

Assume  $f$  is convex and  $f(0) = 0$



# A Useful Property

## Assumption

Assume  $f$  is convex and  $f(0) = 0$

## Claim

For  $0 \leq \alpha \leq 1$ ,

$$f(\alpha a) \leq \alpha f(a)$$





# A Useful Property

## Assumption

Assume  $f$  is convex and  $f(0) = 0$

## Claim

For  $0 \leq \alpha \leq 1$ ,

$$f(\alpha a) \leq \alpha f(a)$$

- Apply convexity with  $b = 0$
- Use  $f(0) = 0$



# Subadditivity of Convex Functions

## Claim

For all  $x, y \geq 0$ ,

$$f(x) + f(y) \leq f(x + y)$$



# Subadditivity of Convex Functions

## Claim

For all  $x, y \geq 0$ ,

$$f(x) + f(y) \leq f(x + y)$$

- Let  $\alpha = \frac{x}{x+y}$
- Use  $f(\alpha(x+y)) \leq \alpha f(x+y)$
- Rearranging gives the result



# Applying This to the Recurrence

- Let  $f(n) = T(n)$
- Assume  $T$  is non-decreasing and convex



# Applying This to the Recurrence

- Let  $f(n) = T(n)$
- Assume  $T$  is non-decreasing and convex

$$T(n/5) + T(7n/10) \leq T\left(\frac{n}{5} + \frac{7n}{10}\right) = T\left(\frac{9n}{10}\right)$$



# Applying This to the Recurrence

- Let  $f(n) = T(n)$
- Assume  $T$  is non-decreasing and convex

$$T(n/5) + T(7n/10) \leq T\left(\frac{n}{5} + \frac{7n}{10}\right) = T\left(\frac{9n}{10}\right)$$

$$T(n) \leq T(9n/10) + cn$$



# Solving the Final Recurrence

## Final Recurrence

$$T(n) \leq T(9n/10) + cn$$



# Solving the Final Recurrence

## Final Recurrence

$$T(n) \leq T(9n/10) + cn$$

- Recursion depth:  $O(\log n)$
- Total work:

$$cn + c(9n/10) + c(9^2n/10^2) + \dots$$





# Solving the Final Recurrence

## Final Recurrence

$$T(n) \leq T(9n/10) + cn$$

- Recursion depth:  $O(\log n)$
- Total work:

$$cn + c(9n/10) + c(9^2n/10^2) + \dots$$

$$T(n) = O(n)$$



# Takeaway

- Median of Medians gives a **deterministic linear-time** selection algorithm
- Careful pivot choice guarantees balanced recursion
- Convexity provides a clean way to reason about the recurrence

Worst-case linear time is achievable!

