

WASSERMAN 8457-8458

Terms & Conditions

Electronic Supporting Information files are available without a subscription to ACS Web Editions. The American Chemical Society holds a copyright ownership interest in any copyrightable Supporting Information. Files available from the ACS website may be downloaded for personal use only. Users are not otherwise permitted to reproduce, republish, redistribute, or sell any Supporting Information from the ACS website, either in whole or in part, in either machine-readable form or any other form without permission from the American Chemical Society. For permission to reproduce, republish and redistribute this material, requesters must process their own requests via the RightsLink permission system. Information about how to use the RightsLink permission system can be found at <http://pubs.acs.org/page/copyright/permissions.html>.



ACS Publications

MOST TRUSTED. MOST CITED. MOST READ.

J-8458-m1


Experimental Section

General Information

Melting points were determined on a Thomas Hoover capillary melting point apparatus and were uncorrected. Infrared spectra were recorded on a Perkin Elmer 1420 Spectrophotometer or Nicolet S-5X FT-IR. Proton and carbon nuclear magnetic resonance spectra were determined in the indicated solvent on a Bruker WM-250 spectrometer. In most cases, trimethyl silane was used as the internal standard (δ 0.00 ppm). Low-resolution mass spectra were recorded on a Hewlett Packard 5985A or 5989A mass spectrometer using electron ionization (20 EV). High-resolution mass spectra, using either electron ionization (EI) or chemical ionization (CI), were performed by Mr. Dan Pentek on a Kratos MS-80RFA at Yale University, Instrument Center. Elemental analyses were performed by Atlantic Microlabs, Norcross, Georgia.


Reactions were generally carried out in oven-dried or flame-dried glassware under a dry nitrogen or argon atmosphere. Flash chromatography was performed on silica gel 60 (230-400 mesh, EM laboratories). All chromatographic solvents were distilled prior to use with the exception of diethyl ether which was obtained in anhydrous form. Thin layer chromatography (TLC) was performed on glass plates pre-coated with silica gel 60 F₂₅₄ (0.25 mm, EM laboratories). Visualization was carried out using UV lamp, iodine, phosphomolybdic acid, or *p*-anisaldehyde stain.

Anhydrous solvents were distilled prior to use as follows: THF was distilled from sodium/benzophenone ketyl; benzene, methylene chloride and hexane were distilled from calcium hydride; ethyl acetate was distilled from K₂CO₃.



J-8458-m2


Phosporanylidene. A quantity of 47.6 mL (0.19 mole, 1.8 equivalents) of Bis (trimethylsilyl) acetamide (BSA) was added in a dropwise fashion over a period of 5 minutes to a vigorously stirred solution of 37.5 g (0.11 mole, 1.0 equivalent) of methyl (triphenylphosphoranylidene) acetate in 500 mL of benzene at 8° C, followed by 10.5 mL (0.11 mole, 1.0 equivalent) of 3-chloropropionyl chloride. The resulting solution was stirred for an additional 10 min. at 8° C and 30 min. at room temperature. The mixture was then washed 2 X 200 mL of H₂O, the combined aqueous layers were extracted 2 X 200 mL of ether, and the combined organic extracts were washed 1 X 200 mL of brine, dried over anhydrous MgSO₄ and solvent removed under reduced pressure. The resulting pale orange solid was washed 3 X 300 mL hexanes which resulted in 41.0 g (88%) of phosporanylidene as a pale yellow solid, mp 114-115° C from ethyl acetate/hexane; ¹H NMR (CDCl₃) δ 3.16 (s, 3H), 3.39 (t, *J* = 7.9 Hz, 2H), 3.81 (t, *J* = 7.9 Hz, 2H), 7.40-7.58 (m, 9H), 7.61-7.72 (m, 6H); IR (CCl₄) 3060, 2940, 1668, 1432, 1380, 1300, 1110, 1090 cm⁻¹; Anal. Calcd for C₂₄H₂₂ClO₃P: C, 67.85; H, 5.22. Found C, 67.73; H, 5.23.



VTC-methyl ester 7. A quantity of 9.0 g (0.021 mole) of **phosporanylidene** in 150 mL of 2:1 CH₂Cl₂/CH₃OH at -78° C was treated with O₃ until a light blue color persisted. The solution was then warmed to room temperature and concentrated under reduced pressure. The residue was dissolved in 100 mL THF and treated, with vigorous stirring, with 70 mL of saturated aqueous NaHCO₃ at 0° C for 10 min., and warmed to room temperature for 2.5 hr. The reaction mixture was then diluted with 100 mL of H₂O, extracted 3 X 150 mL ethyl acetate, the combined organic extracts were washed 1 X 100 mL brine, dried over anhydrous MgSO₄, and concentrated under reduced pressure. The residue was gradient column chromatographed (silica gel, 10% ethyl acetate hexane to 40% ethyl acetate hexane)


J-84/58-m3

to afford 3.0 g (89%) of **VTC-methyl ester 7** as a light yellow oil R_f 0.43 (silica gel, 1:1 ethyl acetate/hexane); ^1H NMR (CDCl_3) δ 3.85 (s, 3H), 5.07 (s, 2H), 6.04 (M, 1H), 6.63 (m, 2H); ^{13}C NMR (CDCl_3) δ 53.56, 60.36, 129.02, 133.38, 169.49, 191.33; IR (neat) 3430, 2970, 1750, 1710, 1615, 1440, 1410, 1380, 1270, 1120, 1060, 990 cm^{-1} ; Mass Spec (CI) m/z , 161 (M^+), 143, 139, 138, 129, 114, 85. Exact Mass Calcd For $\text{C}_6\text{H}_8\text{O}_5$ (hydrate) 161.0449. Found: 161.0450. Exact Mass Calcd For $\text{C}_6\text{H}_6\text{O}_4$ (tricarbonyl) 143.0344. Found: 1143.0342.




N-methyl-2-methoxycarbonyl-3-hydroxy pyrrole (8). A vigorously stirred solution of 2.0 g (0.0125 mole) of **VTC-methyl ester 7** in 400 mL of CH_2Cl_2 at 0°C was treated in a dropwise fashion over a period of 5 minutes with 1.1 mL (0.0125 mole) 40% methyl amine in H_2O . The resulting solution was stirred for 15 min. at room temperature, at which time no methyl-VTC **7** was visible by TLC, 25 g SiO_2 was added in one portion and stirred for an additional hour. The resulting mixture was filtered through a Celite pad and washed with 200 mL CH_2Cl_2 followed by 200 mL ethyl acetate, the filtrate was concentrated under reduced pressure. The residue was column chromatographed (silica gel, 1:1 ethyl acetate/hexane) to afford 0.6 g (32%) of **N-methyl-2-methoxycarbonyl-3-hydroxy pyrrole (8)** as a light brown solid mp $43\text{--}44^\circ\text{C}$. R_f 0.46 (silica gel, 30% ethyl acetate/hexane); ^1H NMR (CDCl_3) δ 3.74 (s, 3H), 3.89 (s, 3H), 5.72 (d, $J = 2.8$ Hz, 1H), 6.53 (d, $J = 2.8$ Hz, 1H), 7.98 (bs, 1H); IR (CHCl_3) 3500, 2300, 1720, 1660, 1470, 1450, 1410, 1390, 1335, 1120, 1090, 1075 cm^{-1} . Anal. Calcd for $\text{C}_7\text{H}_9\text{NO}_3$: C, 54.19; H, 5.85; N, 9.03. Found: C, 54.35; H, 5.92; N, 9.10. Mass Spec (EI, 20 EV) (m/e) 155 (M^+), 124, 123, 95, 86, 84; Exact Mass Calcd for $\text{C}_7\text{H}_9\text{NO}_3$ 155.0583. Found 155.0587.

J-8458-m4




N-methyl-4-bromo-2-methoxycarbonyl-3-hydroxy pyrrole (8a). A vigorously stirred solution of 0.57 g (3.67 mmole) of **N-methyl-2-methoxycarbonyl-3-hydroxy pyrrole (8)** in 40 mL of CH_2Cl_2 at -78°C was treated with 0.58 g (3.30 mmole) of N-bromosuccinimide for 2 hr. The reaction was quenched by addition of 15 mL 10% aqueous Na_2SO_3 . The mixture was then warmed to room temperature and extracted 2 X 150 mL CH_2Cl_2 , the combined organic extracts were dried over anhydrous MgSO_4 , concentrated under reduced pressure, and the residue was column chromatographed (silica gel, 20% ethyl acetate / hexanes) to afford 0.64 g (83%) of **N-methyl-4-bromo-2-methoxycarbonyl-3-hydroxy pyrrole (8a)** as a white solid mp $85\text{--}87^\circ\text{C}$. R_f 0.38 (silica gel, 20% ethyl acetate hexane); ^1H NMR (CDCl_3) δ 3.75 (s, 3H), 3.91 (s, 3H), 6.59 (s, 1H), 8.05 (bs, 1H); IR (CCl_4) 3495, 2950, 1715, 1664, 1485, 1470, 1450, 1405, 1370, 1335, 1290, 1160, 1110, 1100, 1050 cm^{-1} ; Mass Spectrum (EI, 20EV), (m/e) 235 (M^+), 233, 203, 201; Exact Mass Calcd for $\text{C}_7\text{H}_8\text{BrNO}_3$ 234.9667. Found: 234.9668. Anal Calcd for $\text{C}_7\text{H}_8\text{BrNO}_3$ C, 35.92; H, 3.45; N, 5.98. Found C, 36.17; H, 3.50; N, 6.08.



N-methyl-4-bromo-2-methoxycarbonyl-3-methoxy pyrrole (8b). A vigorously stirred solution of 0.42 g (1.79 mmole) of **N-methyl-4-bromo-2-methoxycarbonyl-3-hydroxy pyrrole (8a)** in 10 mL of THF at 0°C was treated in one portion with 48 mg (1.96 mmole) of NaH, and the

J-8458.ms


resulting solution was stirred for 5 minutes. At that time 0.2 mL (1.96 mmole) of dimethylsulfate (DMS) was added. The resulting solution stirred overnight at room temperature at which time 20 mL of H₂O was added. The reaction mixture was extracted 2 X 100 mL CH₂Cl₂ and the combined organic extracts were dried over anhydrous MgSO₄, the solvent was removed under reduced pressure and residue chromatographed (silica gel, 20% ethyl acetate/hexanes) to afford 0.43 g (99%) of **N-methyl-4-bromo-2-methoxycarbonyl-3-methoxy pyrrole (8b)** as a light yellow solid mp 55-56° C. *R_f* 0.51 (silica gel, 20% ethyl acetate/hexanes); ¹H NMR (CDCl₃) δ 3.82 (s, 3H), 3.87 (s, 6H), 6.65 (s, 1H); IR (CCl₄) 3000, 2945, 2920, 1700, 1440, 1405, 1360, 1275, 1170, 1105, 1060 cm⁻¹; Mass Spectrum (EI, 20EV), (*m/e*) 249 (M⁺), 247, 216, 214, 202, 200; Exact Mass Calcd for C₈H₁₀BrNO₃: 248.9824. Found: 248.9829. Anal Calcd for C₈H₁₀BrNO₃: C, 38.73; H, 4.06; N, 5.65. Found: C, 38.83; H, 4.07; N, 5.60.



4,4'-Dimethoxy-5,5'-dimethoxycarbonyl-1,1'-dimethyl-3,3'-bipyrrole (6a). A solution of 66 mg (0.266 mmole) of **N-methyl-4-bromo-2-methoxycarbonyl-3-methoxy pyrrole (8b)** and 78 mg (0.334 mmole) of CuBr₂ (activated by heating with a flame under aspirator vacuum until slight Br₂ evolution occurs and backfilling with N₂, that process repeated 3 times) in 1 mL of THF at -78° C was treated with 0.23 mL (0.346 mmole) 1.51 M *n*-BuLi in pentane. The resulting mixture was stirred at -78° C for 30 min. warmed to 0° C for 1 hr and finally to room temperature for 1 hr. The reaction was quenched by addition of 10 mL dilute aqueous NH₄OH (2 mL conc. NH₄OH in 8 mL H₂O), extracted 3 X 20 mL CH₂Cl₂, the combined organic extracts were dried over anhydrous MgSO₄, the solvent was removed under reduced pressure and residue chromatographed (silica gel, 20% ethyl acetate/hexanes) to afford 37 mg (83%) of **4,4'-Dimethoxy-5,5'-dimethoxycarbonyl-1,1'-dimethyl-3,3'-bipyrrole**

J 8458-176

(6a) as a white solid mp 172-173° C. R_f 0.25 (silica gel, 20% ethyl acetate/hexanes); ^1H NMR (CDCl_3) δ 3.80 (s, 6H), 3.86 (s, 6H), 3.89 (s, 6H), 7.06 (s, 2H); IR (CCl_4) 3000, 2945, 1695, 1440, 1400, 1360, 1350, 1270, 1160, 1100, 1060 cm^{-1} ; Exact Mass Calcd for $\text{C}_{16}\text{H}_{20}\text{NO}_6$: 3336.1321. Found: 336.1325.



Isochrysohermidin (5). A quantity of 9.0 g (0.034 mole) of PPh_3 in 150 mL of CH_2Cl_2 at -78°C was treated with O_3 (to generate $\text{PPh}_3\cdot\text{O}_3$) until a light blue color persisted. The solution was then purged with N_2 (until the light blue color faded) to remove the excess O_3 . The reaction flask was connected through a glass tube to another flask containing 14 mg (0.042 mmole) of **4,4'-Dimethoxy-5,5'-dimethoxycarbonyl-1,1'-dimethyl-3,3'-bipyrrole (6a)** in 20 mL of CH_2Cl_2 at 0°C . N_2 was bubbled through the $\text{PPh}_3\cdot\text{O}_3$ solution while it was allowed to warm to room temperature, thus generating singlet oxygen, which was bubbled through the bipyrrole/ CH_2Cl_2 solution for 30 min. The solvent was then removed from the bipyrrole/ CH_2Cl_2 solution under reduced pressure and the residue chromatographed (silica gel, ethyl acetate) the fractions with R_f 0.2 to 0.5 were collected to afford 7 mg (42%) of a 1:1 mixture of *d,l* and *meso* diastereomers of **isochrysohermidin (5)**. Purification of the *d,l*-**5** isomer was accomplished by selective recrystallization from ethyl acetate to afford 2.9 mg (18%) of pure *d,l*-**5**. Purification of *meso*-**5** was accomplished by column chromatography (silica gel, ethyl acetate) of the mother liquor.

d,l-**5**: mp 264-267° C from ethyl acetate (white needles) lit^{5a} mp 265-268° C from ethyl acetate. R_f 0.31 (silica gel, ethyl acetate); ^1H NMR (CDCl_3) δ 2.81 (s, 6H), 3.82 (s, 6H), 3.97 (s, 6H); IR (CH_2Cl_2) 3478, 3407, 3186, 1760, 1740, 1700, 1684, 1644, 1393, 1363, 1253, 1147, 1052 cm^{-1} ;

J- 8458-m7

Mass Spec (EI, 20EV), (m/e) 400 (M^+), 368, 341, 323, 277, 236, 196, 129, 83; Exact Mass Calcd for $C_{16}H_{20}N_2O_{10}$: 400.1118. Found: 400.1117.

meso-5: mp 207-210° C from ethyl acetate (white needles) lit^{5b} mp 207-209° C. R_f 0.21 (silica gel, ethyl acetate); 1H NMR ($CDCl_3$) δ 2.79 (s, 6H), 3.88 (s, 6H), 3.97 (s, 6H); IR (CH_2Cl_2) 3478, 1750, 1700, 1684, 1644, 1363, 1300, 1253, 1147, 1052 cm^{-1} ; Mass Spec (EI, 20EV), (m/e) 400 (M^+), 355, 341, 323, 309, 295, 277, 265, 236, 208, 201; Exact Mass (CI) Calcd for $C_{16}H_{20}N_2O_{10}$: 401.1196. Found: 401.1174.

J-8458-m8

5,5'-Dimethoxy-3,3',6,6'-tetramethoxycarbonyl-4,4'-bi-1,2-diazine (11). Mp 124-125°C (Et₂O, pale yellow plates); ¹H NMR (CDCl₃, 500 MHz) δ 4.12 (s, 3H, OMe), 3.89 (s, 3H, CO₂Me), 3.82 (s, 3H, CO₂Me); ¹³C NMR (CDCl₃, 50 MHz) δ 164.4 (e, C-3 CO₂Me), 163.8 (e, C6 CO₂Me), 155.9 (e, C-5), 150.4 (e, C-4), 147.3 (e, C-3), 125.8 (e, C-6), 61.6 (o, OMe), 53.7 (o, CO₂Me), 53.4 (o, CO₂Me); UV (CHCl₃) λ_{max} 248 nm (ε 58000); IR (KBr) ν_{max} 3745, 1735, 1697, 1438, 1385, 1288, 1249, 1211, 1098, 1054 cm⁻¹; EIMS, *m/e* (relative intensity) 450 (M⁺, 4) 378 (14), 377 (95), 325 (35), 211 (91), 179 (39), 59 (base); CIMS (2-methylpropane), *m/e* 451 (M + H⁺, base); EIHRMS, *m/e* 450.1021 (C₁₈H₁₈N₄O₁₀ requires 450.1023).

Anal. Calcd for C₁₈H₁₈N₄O₁₀: C, 48.01; H, 4.03; N, 12.44. Found: C, 47.72; H, 3.93; N, 12.22.

4,4'-Dimethoxy-2,2',5,5'-tetramethoxycarbonyl-3,3'-bipyrrole (12). Mp 199-200°C (*i*PrOH, colorless needles); ¹H NMR (CDCl₃, 500 MHz) δ 9.44 (bs, 1H, NH), 3.98 (s, 3H, OMe), 3.78 (s, 3H, CO₂Me), 3.72 (s, 3H, CO₂Me); ¹³C NMR (CDCl₃, 50 MHz) δ 160.6 (e, C-2 CO₂Me), 160.3 (e, C-5 CO₂Me), 150.7 (e, C-4), 120.6 (e, C-3), 114.1 (e, C-2), 112.6 (e, C-5), 62.0 (o, OMe), 51.8 (o, CO₂Me), 51.8 (o, CO₂Me); UV (CHCl₃) λ_{max} 282 nm (ε 46000); IR (KBr) ν_{max} 3315, 3286, 2956, 1727, 1705, 1556, 1491, 1440, 1303, 1267, 1247, 1200, 1143, 1110, 1017, 957, 784 cm⁻¹; EIMS, *m/e* 424 (M⁺, base); CIMS (2-methylpropane), *m/e* 425 (M + H⁺, base); EIHRMS *m/e* 424.1119 (C₁₈H₂₀N₂O₁₀ requires 424.1118).

Anal. Calcd for C₁₈H₂₀N₂O₁₀: C, 50.95; H, 4.75; N, 6.60. Found: C, 50.79; H, 4.88; N, 6.54.

4,4'-Dimethoxy-2,2',5,5'-tetramethoxycarbonyl-1,1'-dimethyl-3,3'-bipyrrole (13). Mp 119-120°C (*i*PrOH, colorless prisms); ¹H NMR (CDCl₃, 200 MHz) δ 4.16 (s, 3H, NMe),

J 8458 m9

3.91 (s, 3H, OMe), 3.63 (s, 3H, CO₂Me), 3.60 (s, 3H, CO₂Me); ¹³C NMR (CDCl₃, 50 MHz) δ 161.7 (e, C-2 CO₂Me), 161.4 (e, C-5 CO₂Me), 150.6 (e, C-4), 124.3 (e, C-3), 117.2 (e, C-2), 113.7 (e, C-5), 62.1 (o, OMe), 51.5 (o, CO₂Me), 51.4 (o, CO₂Me), 34.9 (o, NMe); UV (CHCl₃) λ_{max} 286 nm (ε 22000); IR (KBr) ν_{max} 2955, 1719, 1483, 1458, 1433, 1414, 1399, 1350, 1283, 1241, 1206, 1116, 1032 cm⁻¹; EIMS, *m/e* 452 (M⁺, base); CIMS (2-methylpropane), *m/e* 453 (M + H⁺, base); EIHRMS, *m/e* 452.1431 (C₂₀H₂₄N₂O₁₀ requires 452.1431).

Anal. Calcd for C₂₀H₂₄N₂O₁₀: C, 53.10; H, 5.35; N, 6.19. Found: C, 52.89; H, 5.50; N, 6.17.

4,4'-Dimethoxy-1,1'-dimethyl-3,3'-bipyrrole-2,2',5,5'-tetracarboxylic Acid (14). Mp 190-192°C (dec, *i*PrOH-Et₂O, white solid); ¹H NMR (*d*₆-acetone, 200 MHz) δ 4.14 (s, 3H, NMe), 3.66 (s, 3H, OMe); ¹³C NMR (CD₃OD, 50 MHz) δ 164.2 (e, C-2 CO₂H), 163.6 (e, C-5 CO₂H), 152.2 (e, C-4), 126.6 (e, C-3), 118.5 (e, C-2), 115.5 (e, C-5), 47.7 (o, OMe), 35.4 (o, NMe); UV (CH₃CN) λ_{max} 288 nm (ε 22700); IR (KBr) ν_{max} 3855, 3745, 3630, 3423, 2966, 2616, 1685, 1526, 1444, 1419, 1352, 1262, 1207, 1114 cm⁻¹; FABHRMS (NBA), *m/e* 397.0810 (C₁₆H₁₆N₂O₁₀ + H⁺ requires 397.0883).

Anal. calcd for C₁₆H₁₆N₂O₁₀: C, 48.49; H, 4.07; N, 7.07. Found: C, 48.09; H, 4.44; N, 7.38.

4,4'-Dimethoxy-5,5'-dimethoxycarbonyl-1,1'-dimethyl-3,3'-bipyrrole-2,2'-dicarboxylic Acid (6b). Mp 175-177°C (dec, *i*PrOH-Et₂O, white solid); ¹H NMR (*d*₆-DMSO, 400 MHz) δ 4.02 (s, 3H, NMe), 3.81 (s, 3H, OMe), 3.52 (s, 3H, CO₂Me); ¹³C NMR (*d*₆-DMSO, 100 MHz) δ 161.8 (e, CO₂H), 149.9 (e, CO₂Me), 149.8 (e, C-4), 124.9 (e, C-3), 116.1 (e, C-2), 113.6 (e, C-5), 61.7 (o, OMe), 51.5 (o, CO₂Me), 34.8 (o, NMe); UV (THF) λ_{max} 280 nm (ε 24800); IR (KBr) ν_{max} 3418, 2923, 1720, 1655, 1443, 1384, 1261, 1101, 800 cm⁻¹;

J-8458m10

FABHRMS (NBA), m/e 424.1118 ($C_{18}H_{20}N_2O_{10}$ requires 424.1118).

Anal. Calcd for $C_{18}H_{20}N_2O_{10}$: C, 50.95; H, 4.75; N, 6.60. Found: C, 51.30; H, 5.07; N, 6.50.

Isocrysohermidin (5). A solution of **6b** (40.0 mg, 0.094 mmol) in collidine (30 mL), H_2O (90 mL), and 2-propanol (15 mL) containing Rose Bengal (800 μg , 7.9×10^{-4} mmol, 8 mequiv) in a pyrex tube was treated with a steady stream of O_2 (5 psi). The solution was irradiated with a Hanovia high-pressure mercury lamp (450 W) through a uranium yellow glass filter (transmits > 330 nm) at $22^\circ C$ for 1 h. The reaction mixture was partitioned between 10% aqueous HCl and EtOAc. The organic layer was dried (Na_2SO_4) and concentrated under reduced pressure. Chromatography (SiO_2 , 1 x 10 cm, EtOAc) afforded 35 mg of crude product (R_f 0.2-0.5, EtOAc). The crude product which contained a mixture of d,l- and meso-**5** as well as potential ring opened isomers was dissolved in hot EtOAc which was allowed to slowly concentrate (recrystallization of the mother liquor was repeated two times) to afford d,l-**5** as a white solid (14 mg, 38 mg theoretical, 37%; typically 37-43%). The mother liquor was concentrated and the residue purified by chromatography (SiO_2 , 1 x 10 cm, EtOAc, 3x) to afford pure meso-**5** (3 mg, 38 mg theoretical, 8%). For d,l-**5**: mp $266-268^\circ C$ (EtOAc, colorless needles); lit⁶ mp $265-268^\circ C$ (EtOAc); 1H NMR ($CDCl_3$, 400 MHz) δ 6.62 (bs, 1H, OH), 3.98 (s, 3H, OMe), 3.83 (s, 3H, CO_2Me), 2.84 (s, 3H, NMe); ^{13}C NMR (d_6 -DMSO, 100 MHz) δ 169.7 (e, CONMe), 168.6 (e, CO_2Me), 167.9 (e, C-4), 97.2 (e, C-3), 86.8 (e, C-5), 58.8 (o, CO_2Me), 53.3 (o, OMe), 24.1 (o, NMe); IR (KBr) ν_{max} 3478, 3408, 3187, 1760, 1740, 1700, 1684, 1644, 1393, 1363, 1253, 1147, 1052 cm^{-1} ; FABHRMS (NBA-CsI), m/e 533.0178 ($C_{16}H_{20}N_2O_{10} + Cs^+$ requires 533.0172).

For meso-**5**: mp $207-209^\circ C$ (EtOAc, white powder); lit⁵ mp $207-209^\circ C$ (1:1, EtOAc-

J-8458-m11

Et₂O); ¹H NMR (CDCl₃, 400 MHz) δ 4.65 (bs, 1H, OH), 3.98 (s, 3H, OMe), 3.87 (s, 3H, CO₂Me), 2.78 (s, 3H, NMe); ¹³C NMR (d₆-DMSO, 100 MHz) δ 169.6 (e, CONMe), 168.5 (e, CO₂Me), 167.7 (e, C-4), 97.1 (e, C-3), 86.7 (e, C-5), 58.7 (o, CO₂Me), 53.2 (o, OMe), 23.9 (o, NMe); IR (KBr) ν_{max} 3413, 3180, 2954, 1753, 1730, 1706, 1692, 1638, 1440, 1366, 1307, 1253, 1146, 1053, 938 cm⁻¹; FABHRMS (NBA-CsI), *m/e* 533.0172 (C₁₆H₂₀N₂O₁₀ + Cs⁺ requires 533.0172).

Singlet Oxygen Reaction of **6b**.^a

| solvent | d,l- 5 |
|--|--------------------------------------|
| <i>i</i> PrOH-H ₂ O (3:1) | --- |
| CH ₃ CN-H ₂ O (3:1) | --- |
| <i>i</i> PrOH-Pyridine-H ₂ O (3:1:1) | 6% ^b 16% ^c |
| CH ₃ CN-Pyridine-H ₂ O (3:1:1) | 5% ^b |
| <i>i</i> PrOH-H ₂ O-DBU (6:2:1) | --- |
| Pyridine-H ₂ O (1:1) | 10% ^b 23% ^c |
| Pyridine-H ₂ O (1:3) | 28% ^c |
| Pyridine-H ₂ O (3:1) | 11% ^c |
| Collidine-H ₂ O (1:10) | 26% ^c |
| Collidine-H ₂ O (1:3) | 40% ^c |
| Collidine-H ₂ O (1:1) | 36% ^c |

^a ¹O₂ source: rose bengal (8 mequiv), Hanovia high-pressure lamp (450 W), uranium yellow glass filter (transmits > 330 nm), O₂, 22°C, 1 h. ^b Product (R_f 0.39, SiO₂, EtOAc) was isolated by chromatography (SiO₂, EtOAc) and recrystallized (EtOAc) to provide pure d,l-**5**. ^c Product (R_f 0.2-0.5) isolated by chromatography (SiO₂, EtOAc) and recrystallized (EtOAc) to provide pure d,l-**5**. Meso-**5** and isomers of **5** remained in mother liquor.