

CUSHING 9323-9324


Terms & Conditions

Electronic Supporting Information files are available without a subscription to ACS Web Editions. The American Chemical Society holds a copyright ownership interest in any copyrightable Supporting Information. Files available from the ACS website may be downloaded for personal use only. Users are not otherwise permitted to reproduce, republish, redistribute, or sell any Supporting Information from the ACS website, either in whole or in part, in either machine-readable form or any other form without permission from the American Chemical Society. For permission to reproduce, republish and redistribute this material, requesters must process their own requests via the RightsLink permission system. Information about how to use the RightsLink permission system can be found at <http://pubs.acs.org/page/copyright/permissions.html>.



Supplementary material

J-9324-m1




^1H NMR (300MHz) (CDCl_3) δ TMS; 1.22(3H, s); 1.56(3H, s); 3.03(1H, d, $J = 11.4\text{Hz}$, D_2O exch.); 3.63(1H, ddd, $J = 4.0, 0.9, 11.3\text{Hz}$); 4.19(1H, dd, $J = 0.9, 12.3\text{Hz}$); 4.31(1H, dd, $J = 4.0, 12.3\text{Hz}$); 6.49(1H, dd, $J = 2.2, 3.1\text{Hz}$); 6.78(1H, d, $J = 8.4\text{Hz}$); 7.16-7.19(2H, m); 8.29(1H, s, D_2O exch.).

IR(KBr) 3340,2984,1580,1504,1444,1338,1224,1133,1057,814, 753 cm^{-1}

microanalysis calc'd for $\text{C}_{13}\text{H}_{15}\text{NO}_3$: C, 66.94; H, 6.48; N, 6.00; Found: C, 67.16; H, 6.63; N, 5.79.


m.p. 202-205 $^\circ\text{C}$



^1H NMR (300Hz) (CDCl_3) δ TMS; 0.14(6H, s); 0.89(9H, s); 1.12(3H, s); 1.48(3H, s); 3.88(1H, dd, $J = 9.2, 11.5\text{Hz}$); 3.98(1H, dd, $J = 3.2, 9.2\text{Hz}$); 4.22(1H, dd, $J = 3.2, 11.5\text{Hz}$); 6.48(1H, dd, $J = 2.2, 3.1\text{Hz}$); 6.76(1H, d, $J = 8.4\text{Hz}$); 7.14(2H, ddd, $J = 2.4, 3.4, 3.5\text{Hz}$); 8.21(1H, s, D_2O exch.)

IR(neat) 3412,2936,1500,1438,1234,1093,833 cm^{-1} .

m.p. 118-119°C




^1H NMR (300Hz) (CDCl_3) δ TMS; 0.90(9H, s); 1.13(3H, s); 1.48(3H, s); 2.28(6H, s); 3.58(2H, s); 3.58(2H, s); 3.88(1H, dd, $J = 9.2, 11.4\text{Hz}$); 3.98(1H, dd, $J = 3.2, 9.1\text{Hz}$); 4.21(1H, dd, $J = 3.2, 11.5\text{Hz}$); 6.76(1H, d, $J = 8.4\text{Hz}$); 8.44(1H, s, D_2O exch.)

IR (neat) 2932, 1502, 1458, 1360, 1251, 1218, 1093, 837, 777 cm^{-1} .

microanalysis calc'd for $\text{C}_{22}\text{H}_{36}\text{N}_2\text{O}_3\text{Si}$: C, 65.31; H, 8.97; N, 6.92; found: C, 65.09; H, 8.77; N, 6.73.

m.p. 152°C



^1H NMR (300MHz) (CDCl_3) δ TMS 1.76(3H, s); 1.99-2.10(2H, br s); 2.17-2.26(2H, m); 2.78(1H, dd, $J = 7.3, 14.5\text{Hz}$); 2.90(1H, dd, $J = 8.0, 14.8\text{Hz}$); 3.54-3.63(1H, m); 3.84(1H, dt, $J = 12.3, 8.4\text{Hz}$); 3.95(1H, $d\frac{1}{2}\text{ABq}$, $J = 3.4, 17.6\text{Hz}$); 4.10(1H, $\frac{1}{2}\text{ABq}$, $J = 17.6\text{Hz}$); 6.55(1H, t, $J = 7.2\text{Hz}$); 7.96(1H, br s, D_2O exch.); 9.45(1H, s),

IR (neat) 3246, 1684, 1448, 1326, 1107 cm^{-1} .

$[\alpha]_{25}^{\text{D}} = -1.51/1.92 \times 10^{-2} = -78.4 (\text{CH}_2\text{Cl}_2, c = 0.164)$


J-9324-m3

IR (neat) 3246, 1684, 1448, 1326, 1107 cm^{-1} .

$[\alpha]_{25}^D = -1.51/1.92 \times 10^{-2} = -78.4 (\text{CH}_2\text{Cl}_2, c = 0.164)$

microanalysis calc'd for $\text{C}_{12}\text{H}_{15}\text{N}_2\text{O}_3$ C, 61.00; H, 6.83; N, 11.86; found: C, 60.88; H, 6.66; N, 11.71.

EI HRMS 236.1155 ($\text{C}_{12}\text{H}_{15}\text{N}_2\text{O}_3$ requires 236.11609).




^1H NMR (300MHz) (CDCl_3) δ 1.03(9H, s); 1.54(3H, s); 1.92-2.19(4H, m); 2.49(1H, dd, $J = 8.6, 14.1\text{Hz}$); 2.58(1H, dd, $J = 7.5, 14.1\text{Hz}$); 3.44-3.53(1H, m); 3.73(1H, $d^{1/2}\text{ABq}$, $J = 4.1, 16.9\text{Hz}$); 3.78-3.85(1H, m); 4.01(2H, s); 4.06(1H, $^{1/2}\text{ABq}$, $J = 16.9\text{Hz}$); 5.56-5.62(1H, m); 6.38(1H, d, $J = 3.7\text{Hz}$, D_2O exch.); 7.32-7.43(6H, m); 7.62(4H, dd, $J = 1.8, 7.6\text{Hz}$).

IR (neat) 3232(br) 2930, 2857, 1664, 1446, 1435, 1113, 822, 733, 702 cm^{-1} .

$[\alpha]_{25}^D = -1.24/1.92 \times 10^{-2} = -63.3 (\text{CDCl}_3, c = 0.0822)$

microanalysis calc'd for $\text{C}_{28}\text{H}_{36}\text{N}_2\text{O}_3\text{Si}$ C, 70.55; H, 7.61; N, 5.88; found C, 70.60; H, 7.56; N, 5.91.


m.p. 132°C.



^1H NMR (300MHz) (CDCl_3) δ 1.04(9H, s); 1.40(3H, s); 1.86-2.03(2H, m); 2.12-2.31(2H, m); 2.55(1H, d, $J = 7.4\text{Hz}$); 3.43-3.52(1H, m); 3.74-3.82(1H, m); 3.83(3H, s); 3.88(3H, s); 4.03(2H, br s); 5.48-5.53(2H, m); 7.34-7.41(6H, m); 7.57-7.66(4H, m).

IR (neat) 2960, 1790, 1740, 1681, 1430, 1366, 1272, 1223, 1109, 735, 705 cm^{-1} .

microanalysis calcd for $\text{C}_{32}\text{H}_{40}\text{N}_2\text{O}_7\text{Si}$ C, 68.06; H, 7.14; N, 4.96; found C, 67.87; H, 7.27; N, 4.77.




^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 0.10(6H, s); 0.115(3H, s); 0.12(3H, s); 0.87(9H, s); 0.88(9H, s); 1.02(18H, s); 1.096(3H, s); 1.10(3H, s); 1.45(3H, s); 1.46(3H, s); 1.54(6H, s); 1.60-1.88(6H, m); 2.02-2.11(2H, m); 2.92(2H, dd, $J = 7.1, 14.4\text{Hz}$); 2.44(2H, dd, $J = 8.1, 14.5\text{Hz}$); 3.32-3.44(4H, m); 3.60(3H, s); 3.62(3H, s); 3.72-3.93(8H, m); 3.98(4H, br s); 4.18(2H, dd, $J = 2.9, 8.4\text{Hz}$); 5.43(2H, m); 6.38(1H, s, D_2O exch.); 6.41(1H, s, D_2O exch.); 6.74(1H, d, $J = 8.5\text{Hz}$); 6.75(1H, d, $J = 8.5\text{Hz}$); 6.89(1H, d, $J = 2.3\text{Hz}$); 6.92(1H, d, $J = 2.3\text{Hz}$); 7.08(2H, d, $J = 8.5\text{Hz}$); 7.33-7.41(12H, m); 7.61-7.63(8H, m); 8.43(1H, d, $J = 2.9\text{Hz}$, D_2O exch.); 8.64(1H, d, $J = 1.9\text{Hz}$, D_2O exch.).

IR (neat) 3281(br), 2954, 2932, 2856, 1747, 1670, 1665, 1649, 1431, 1251, 1224, 1109, 1088, 733, 706 cm^{-1} .

EI HRMS 893.4457($\text{C}_{50}\text{H}_{67}\text{N}_3\text{O}_8\text{Si}_2$ requires 893.4467).

microanalysis calc'd for $C_{50}H_{67}N_3O_8Si_2$ C, 67.16; H, 7.55; N, 4.70; found C, 66.93; H, 7.36; N, 4.51.

m.p. 106-108°C.



1H NMR (300MHz) ($CDCl_3$) (syn, mixture of two diastereomers) δ TMS 0.146(6H, s); 0.904(18H, s); 1.04(18H, s); 1.126(3H, s); 1.13(3H, s); 1.48(6H, s); 1.64(6H, s); 1.94-2.06(6H, m); 2.20-2.24(2H, m); 2.36-2.46(2H, m); 2.60-2.72(2H, m); 2.98(2H, dd, J = 11.6, 14.1Hz); 3.44-3.57(4H, m); 3.88(2H, dd, J = 6.7, 9.2Hz); 3.97(2H, dd, J = 3.1, 9.1Hz); 4.02-4.06(2H, m); 4.10(4H, s); 4.17-4.25(4H, m); 5.58(2H, m); 5.68(2H, br s, D_2O exch.); 6.75(2H, d, J = 8.5Hz); 6.86(1H, d, J = 2.2Hz); 6.88(1H, J = 2.2Hz); 7.14(2H, d, J = 8.4Hz); 7.26-7.44(12H, m); 7.60-7.64(8H, m); 8.04(1H, s, D_2O exch.); 8.06(1H, s, D_2O exch.).

syn diastereomers separated;

1H NMR (300MHz) ($CDCl_3$) (syn #1, less polar) δ TMS 0.12(3H, s); 0.13(3H, s); 0.88(9H, s); 1.03(9H, s); 1.11(3H, s); 1.46(3H, s); 1.63(3H, s); 1.92-2.04(3H, m); 2.18-2.23(1H, m); 2.39(1H, dd, J = 7.2, 14.2Hz); 2.64(1H, dd, J = 8.7, 14.2Hz); 2.99(1H, dd, J = 11.4, 14.2Hz); 3.42-3.46(1H, m); 3.51(1H, dd, J = 2.7, 14.2Hz); 3.85(1H, dd, J = 9.2, 11.3Hz); 3.94(1H, dd, J = 3.0, 9.1Hz); 3.99-4.06(1H, m); 4.08(2H, s); 4.11-4.15(1H, m); 4.19(1H, dd, J = 3.0, 11.3Hz); 5.58(1H, t, J = 7.8Hz);

5.76(1H, d, J = 2.7Hz, D₂O exch.); 6.73(1H, d, J = 8.4Hz); 6.85(1H, d, J = 2.1Hz);
7.11(1H, d, J = 8.5Hz); 7.26-7.42(6H, m); 7.57-7.63(4H, m); 8.15(1H, s, D₂O exch.).

¹H NMR (300MHz) (CDCl₃) (syn #2, more polar) δTMS 0.12(3H, s); 0.14(3H, s);
0.88(9H, s); 1.03(9H, s); 1.11(3H, s); 1.46(3H, s); 1.62(3H, s); 1.91-2.04(3H, m);
2.18-2.22(1H, m); 2.36(1H, dd, J = 7.3, 14.2Hz); 2.60(1H, dd, J = 8.6, 14.3Hz);
2.97(1H, dd, J = 11.3, 14.2Hz); 3.41-3.44(1H, m); 3.50(1H, dd, J = 3.1, 14.2Hz);
3.86(1H, dd, J = 9.3, 11.3Hz); 3.95(1H, dd, J = 3.0, 9.1Hz); 3.99-4.03(1H, m);
4.08(2H, s); 4.14-4.16(1H, m); 4.20(1H, dd, J = 2.9, 11.6Hz); 5.56(1H, t, J = 7.5Hz);
5.72(1H, d, J = 2.6Hz, D₂O exch.); 6.73(1H, d, J = 8.4Hz); 6.84(1H, d, J = 2.1Hz);
7.11(1H, d, J = 8.4Hz); 7.26-7.42(6H, m); 7.57-7.62(4H, m); 8.07(1H, s, D₂O exch.).

¹H NMR (300MHz) (CDCl₃) (anti, mixture of two diastereomers) δTMS 0.14(6H, s);
0.16(6H, s); 0.90(18H, s); 1.04(9H, s); 1.045(9H, s); 1.09(3H, s); 1.13(3H, s);
1.47(6H, s); 1.53(3H, m); 1.54(3H, m); 1.97-2.17(8H, m); 2.47-2.62(4H, m); 2.78-
2.88(2H, m); 3.54-3.65(4H, m); 3.82-3.99(6H, m); 4.02(4H, s); 4.21(2H, dd, J = 3.1,
11.0Hz); 4.35-4.39(2H, m); 5.52-5.54(2H, m); 5.69(2H, br s, D₂O exch.); 6.60(1H, d, J
= 8.4Hz); 6.63(1H, d, J = 8.4Hz); 6.89(2H, d, J = 2.1Hz); 6.98(2H, d, J = 8.4Hz);
7.36-7.42(12H, m); 7.62-7.69(4H, m); 8.08(2H, br s, D₂O exch.).

IR (neat) (anti) 3289(br), 2929, 2855, 1666, 1444, 1428, 1254, 1222, 1111, 857, 836,
704 cm⁻¹.


IR (neat) (syn) 3274(br), 2929, 2858, 1666, 1651, 1453, 1428, 1250, 1224, 1112, 1052,
858, 838, 777 cm⁻¹.

microanalysis calcd for C₄₉H₆₅N₃O₆Si₂(syn) C, 68.94; H, 7.84; N, 5.02; Found
C, 69.06; H, 7.76; N, 5.03.

m.p. (syn) 167-168°C.

microanalysis calcd for C₄₈H₆₅N₃O₆Si₂(anti) C, 68.94; H, 7.84; N, 5.02; Found
C, 68.76; H, 7.60; N, 4.82.

m.p.(anti) 95-99°C




^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 0.120(12H, s); 0.875(18H, s); 1.02(18H, s); 1.06(3H, s); 1.07(3H, s); 1.45(12H, s); 1.65-2.08(12H, m); 3.07-3.15(2H, m); 3.26(2H, dd, $J = 6.2, 12.6\text{Hz}$); 3.32-3.40(2H, m); 3.61(6H, s); 3.70-3.86(2H, m); 3.91-3.95(4H, m); 3.99(2H, s); 4.15(2H, dd, $J = 3.6, 11.7\text{Hz}$); 4.36-4.40(2H, m); 5.37-5.44(2H, br m); 6.69(2H, d, $J = 8.4\text{Hz}$); 7.01(2H, d, $J = 1.7\text{Hz}$); 7.15(2H, d, $J = 8.4\text{Hz}$); 7.26-7.41(12H, m); 7.58-7.62(8H, m); 8.06(2H, s, D_2O exch.).
IR (neat) 3292, 2932, 1687, 1643, 1447, 1251, 1218, 1109, 837 cm^{-1} .

MS (EI) 849 *m/e* (rel intensity) 849 (M^+ , 8.9) 361 (26) 360 (95) 167(100).

microanalysis calc'd for $\text{C}_{49}\text{H}_{67}\text{N}_3\text{O}_6\text{Si}_2$; C, 69.02; H, 7.94; N, 4.94; Found: C, 69.02; H, 7.88; N, 4.79.

m.p. 74-76°C.




^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 0.13(3H, s); 0.14(9H, s); 0.89(18H, s); 1.03(9H, s); 1.04(9H, s); 1.087(3H, s); 1.093(3H, s); 1.28-1.43(3H, m); 1.48(3H, s); 1.50(6H, s); 1.79-1.89(4H, m); 2.24-2.38(4H, m); 3.22-3.42(6H, m); 3.60(3H, s); 3.62(3H, s); 3.68-3.76(2H, m); 3.79-3.87(2H, m); 3.94(2H, d, $J = 3.4\text{Hz}$); 3.97(4H, br s); 4.15-4.20(2H, m); 4.26-4.32(2H, m); 5.41(2H, t, $J = 7.8\text{Hz}$); 6.701(1H, d, $J = 8.5\text{Hz}$); 6.703(2H, d, $J = 8.4\text{Hz}$); 6.96(1H, d, $J = 2.6\text{Hz}$); 6.97(1H, d, $J = 2.6\text{Hz}$); 7.28(2H, d, $J = 8.5\text{Hz}$); 7.32-7.44(12H, m); 7.60-7.64(8H, m); 7.97(2H, br s, D_2O exch.).

IR (neat) 3304, 2930, 1695, 1645, 1447, 1249, 1221, 836 cm^{-1} .

EI HRMS 849.4550 ($\text{C}_{49}\text{H}_{67}\text{N}_3\text{O}_6\text{Si}_2$ requires 849.4568)

microanalysis calc'd for $\text{C}_{49}\text{H}_{67}\text{N}_3\text{O}_6\text{Si}_2$: C, 69.22; H, 7.94; N, 4.94; found C, 59.06; H, 8.04; N, 4.89.

m.p. 54-58°C.



^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 1.18(6H, s); 1.52(3H, s); 1.53(1H, s); 1.56(1H, s); 1.57(21H, s); 1.61-2.07(10H, m); 2.14(2H, dd, J = 8.6, 14.5Hz); 2.85(2H, br s, D_2O Exch.); 2.92-3.01(2H, m); 3.18-3.35(6H, m); 3.56(2H, br s, D_2O exch.); 3.62(3H, s); 3.64(3H, s); 3.88(4H, br s); 3.91-4.00(2H, m); 4.25(4H, br s); 4.30-4.39(2H, m); 4.98-5.01(2H, m); 6.87(1H, d, J = 8.3Hz); 6.88(1H, d, J = 8.3Hz); 7.16(1H, d, J = 8.3Hz); 7.17(1H, d, J = 8.3Hz); 7.34(1H, s); 7.35(1H, s).


^{13}C NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 13.44, 19.47, 19.68, 23.51, 23.63, 25.13, 25.32, 27.86, 30.34, 30.52, 34.38, 34.84, 35.12, 35.26, 43.38, 43.63, 52.63, 52.70, 62.03, 62.40, 65.26, 65.45, 67.67, 67.76, 70.63, 75.37, 82.57, 82.64, 114.53, 114.73, 116.75, 116.88, 118.18, 118.32, 119.04, 119.12, 126.34, 128.04, 128.10, 129.94, 130.07, 138.64, 138.71, 140.68, 146.15, 148.49, 161.32, 161.50, 168.53, 168.66

IR (neat) 3390(br), 2976, 1752, 1692, 1632, 1491, 1453, 1371, 1251, 1158, 733 cm^{-1} .

microanalysis calc'd for $\text{C}_{32}\text{H}_{43}\text{N}_3\text{O}_8$; C, 64.30; H, 7.25; N, 7.03; found: C, 64.12; H, 7.41; N, 6.88.

EI HRMS m/e 597.3065 ($\text{C}_{32}\text{H}_{43}\text{N}_3\text{O}_8$ requires 597.3050).

m.p. 72-85°C.




^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 1.16(3H, s); 1.18(3H, s); 1.51(3H, s); 1.52(3H, s); 1.55(6H, s); 1.57(18H, s); 1.60-2.14(10H, m, 2H D_2O exch.); 2.22-2.37(4H, m); 3.06-3.18(3H, m, 1H D_2O exch.); 3.26-3.36(5H, m, 1H D_2O exch.); 3.55(3H, s); 3.56(2H, br s); 3.60(3H, s); 3.63-3.72(2H, m); 3.89(4H, m); 4.18-4.23(2H, m); 4.25(4H, br s); 5.21-5.27(2H, m); 6.857(1H, d, $J = 8.3\text{Hz}$); 6.861(1H, d, $J = 8.3\text{Hz}$); 7.22(2H, d, $J = 8.3\text{Hz}$); 7.24(2H, s).

IR (neat) 3401(br), 2976, 1747, 1692, 1632, 1496, 1436, 1371, 1251, 1158, 733 cm^{-1} .

EI HRMS 597.3050 ($\text{C}_{32}\text{H}_{43}\text{N}_3\text{O}_8$ requires 597.3050)

m.p.72-80°C.




^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 1.17(6H, s); 1.52(6H, s); 1.57(18H, s); 1.65(6H, s); 1.73-2.20(12H, s); 2.84(2H, dd, $\tau = 9.0, 14.4\text{Hz}$); 3.06(1H, br s, D_2O exch.); 3.10(1H, br s, D_2O exch.); 3.26-3.36(4H, m); 3.55-3.58(4H, m); 3.62(2H, s); 3.63(2H, s); 3.91(4H, s); 3.95-4.05(2H, m); 4.24-4.25(4H, m); 4.30-4.36(2H, m); 5.28(2H, m); 6.88(2H, d, $J = 8.3\text{Hz}$); 7.14(1H, d, $J = 8.3\text{Hz}$); 7.15(1H, d, $J = 8.3\text{Hz}$); 7.376(1H, s); 7.384(1H, s).

IR (neat) 3403, 2979, 1750, 1716, 1642, 1348, 1154 cm^{-1} .

EI HRMS 615.2709 ($\text{C}_{32}\text{H}_{42}\text{N}_3\text{O}_7\text{Cl}$ requires 615.2711).


microanalysis calc'd for $\text{C}_{32}\text{H}_{42}\text{N}_3\text{O}_7\text{Cl}$ C, 62.38; H, 6.87; N, 6.82; found C, 62.53; H, 6.86; N, 6.67.



^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 1.17(3H, s); 1.18(3H, s); 1.52(3H, s); 1.54(3H, s); 1.57(18H, s); 1.65(6H, s); 1.71-1.92(6H, m); 2.24-2.39(4H, m); 3.03-3.19(4H, m, 2H D_2O exch.); 3.28-3.37(4H, m); 3.56(3H, s); 3.60(3H, s); 3.59-3.75(4H, m); 3.89(4H, s); 4.21-4.29(6H, m); 5.35(2H, t, $J = 7.5\text{Hz}$); 6.86(1H, d, $J = 8.3\text{Hz}$); 6.87(1H, d, $J = 8.3\text{Hz}$); 7.23(2H, d, $J = 8.3\text{Hz}$); 7.22(1H, s); 7.27(1H, s).

IR (neat) 3412(br), 2976, 1752, 1698, 1638, 1365, 1251, 1158 cm^{-1} .

EI HRMS 615.2714 ($\text{C}_{32}\text{H}_{42}\text{N}_3\text{O}_7\text{Cl}$ requires 615.2711)




^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 0.12(6H, s); 0.13(6H, s); 0.88(18H, s); 1.06(6H, s); 1.47(6H, s); 1.59(18H, s); 1.65(6H, s); 1.78-1.98(8H, s); 2.02-2.12(4H, m); 2.86(2H, dd, $J = 9.0, 14.6\text{Hz}$); 3.31-3.34(2H, m); 3.33(2H, dd, $J = 4.0, 13.6\text{Hz}$); 3.62(3H, s); 3.64(3H, s); 3.71-3.79(2H, m); 3.73(1H, dd, $J = 4.2, 9.8\text{Hz}$); 3.77(1H, dd, $J = 4.4, 9.7\text{Hz}$); 3.92(4H, s); 3.94-4.01(4H, m); 4.15(2H, dd, $J = 3.8, 12.4\text{Hz}$); 4.32-4.37(2H, m); 5.28-5.30(2H, m); 6.87(2H, d, $J = 8.3\text{Hz}$); 7.12(2H, d, $J = 8.3\text{Hz}$); 7.13(1H, d, $J = 8.3\text{Hz}$); 7.38(2H, s).

IR (neat) 2930, 1750, 1691, 1652, 1494, 1424, 1366, 1248, 1159, 1088 cm^{-1} .

MS (EI) 729 m/e (rel intensity) 731 ($M+2$, 2.1) 729 (M^+ , 4.2) 629 (9.4) 361 (24.1) 360 (100) 167(94.8) 57.2 (63).

microanalysis calc'd for $\text{C}_{38}\text{H}_{56}\text{N}_3\text{O}_7\text{SiCl}$; C, 62.49; H, 7.73; N, 5.75; found: C, 62.57; H, 7.71, N, 5.55.





^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 0.12(6H, s); 0.13(6H, s); 0.87(18H, s); 1.05(3H, s); 1.06(3H, s); 1.47(6H, s); 1.50-1.53(2H, m); 1.58(18H, s); 1.65(6H, s); 1.72-1.91(4H, m); 2.21-2.37(4H, m); 3.06-3.19(2H, m); 3.28-3.36(4H, m); 3.56(3H, s); 3.60(3H, s); 3.63-3.87(4H, m); 3.89(4H, s); 3.93(2H, dd, $J = 3.9, 9.8\text{Hz}$); 4.13-4.18(2H, m); 4.22-4.35(2H, m); 5.30-5.40(2H, m); 6.85(1H, d, $J = 8.3\text{Hz}$); 6.86(1H, d, $J = 8.3\text{Hz}$); 7.19-7.26(4H, m).

IR (neat) 2949, 1751, 1693, 1652, 1493, 1424, 1369, 1250, 1156, 1086 cm^{-1} .

microanalysis calc'd for $\text{C}_{38}\text{H}_{56}\text{N}_3\text{O}_7\text{SiCl}$; C, 62.49; H, 7.73; N, 5.75; found: C, 62.29; H, 7.61; N, 5.76.

EI HRMS 729.3555($\text{C}_{38}\text{H}_{56}\text{N}_3\text{O}_7\text{SiCl}$ requires 729.3576)






^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 0.12(6H, s); 0.13(6H, s); 0.872(9H, s); 0.875(9H, s); 1.46(6H, s); 1.58(18H, s); 1.61(3H, s); 1.64(3H, s); 1.72-2.03(8H, m); 2.25-2.42(2H, m); 2.47(2H, dd, $J = 5.1, 9.7\text{Hz}$); 2.54(2H, dd, $J = 5.8, 9.7\text{Hz}$); 3.05(1H, $\frac{1}{2}$ ABq, $J = 15.0\text{Hz}$); 3.07(1H, $\frac{1}{2}$ ABq, $J = 15.0\text{Hz}$); 3.31-3.53(6H, m); 3.57(3H, s); 3.64(3H, s); 3.73-3.89(2H, m); 3.94(2H, dd, $J = 3.7, 9.7\text{Hz}$); 4.17(2H, dd, $J = 3.1, 11.6\text{Hz}$); 4.62(1H, s); 4.75(1H, s); 4.78(1H, s); 4.85(1H, s); 6.82(2H, d, $J = 8.4\text{Hz}$); 7.31(1H, d, $J = 8.4\text{Hz}$); 7.38(1H, d, $J = 8.4\text{Hz}$); 7.44(1H, s); 7.52(1H, s).

IR (neat) 2935, 1752, 1684, 1637, 1496, 1418, 1365, 1350, 1250, 1220, 1156, 1083 cm^{-1} .

EI HRMS m/e 693.3834 ($\text{C}_{38}\text{H}_{55}\text{N}_3\text{O}_7\text{Si}$ requires 693.3809).

microanalysis calc'd for $\text{C}_{38}\text{H}_{55}\text{N}_3\text{O}_7\text{Si}$ C, 65.77; H, 7.99; N, 6.05; found C, 65.85; H, 7.99; N, 5.91.

m.p. 105-108°C.




^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 0.081(6H,s); 0.11(6H, s); 0.87(9H, s); 0.88(9H, s); 1.08(3H, s); 1.17(3H, s); 1.26(3H, s); 1.27(3H, s); 1.34(3H, s); 1.35(3H, s); 1.44(3H, s); 1.46(3H, s); 1.56(9H, s); 1.58(9H, s); 1.81–1.90(2H, m); 1.96–2.06(6H, m); 2.20(2H, dd, $J = 10.3, 13.5\text{Hz}$); 2.52–2.60(4H, m); 2.78(2H, dt, $J = 6.5, 12.9\text{Hz}$); 3.36–3.49(2H, m); 3.51–3.57(2H, m); 3.51–3.57(2H, m); 3.63–3.84(4H, m); 3.88–3.92(2H, m); 4.04–4.16(2H, m); 6.24(1H, s, D_2O exch.); 6.26(1H, s, D_2O exch.) 6.78(1H, d, $J = 8.3\text{Hz}$); 6.80(1H, d, $J = 8.5\text{Hz}$); 6.98(1H, d, $J = 8.2\text{Hz}$); 6.99(1H, d, $J = 8.4\text{Hz}$).

^{13}C NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ –5.2, –5.1, –5.0, –4.5, –4.3, 17.6, 18.7, 19.3, 19.7, 19.9, 24.3, 25.5, 25.6, 26.9, 26.2, 27.2, 27.8, 27.9, 28.3, 28.5, 29.1, 31.1, 36.2, 43.8, 50.5, 50.6, 53.3, 54.8, 55.7, 59.4, 60.16, 60.22, 66.3, 67.6, 71.1, 72.7, 75.9, 78.0, 80.5, 84.1, 84.3, 108.3, 112.4, 112.5, 113.6, 117.9, 118.5, 124.6, 124.9, 128.7, 128.9, 129.4, 137.7, 138.3, 139.4, 139.6, 143.0, 143.2, 152.9, 153.0, 168.3, 174.1.


IR(neat) 3214, 2928, 2856, 1745, 1556, 1496, 1443, 1368, 1252, 1233, 1154, 1141, 1091, 1052, 994, 859, 838, 777, 733.

MS (EI) 679 m/e (rel intensity) 679 (M^+ , 0.3) 580 (20.4) 579 (51) 73(100).

EI HRMS m/e 679.3661 ($\text{C}_{37}\text{H}_{53}\text{N}_3\text{O}_7\text{Si}$ requires 679.3653).



^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 0.085(6H, s); 0.11(6H, s); 0.87((H, s); 0.88(9H, s); 1.12(3H, s); 1.15(3H, s); 1.23(3H, s); 1.24(3H, s); 1.36(3H, s); 1.37(3H, s); 1.45(6H, s); 1.59(9H, s); 1.61(9H, s); 1.88–1.92(3H, s); 1.97–2.10(2H, m); 2.17–2.26(2H, m); 2.54–2.63(2H, m); 2.70(2H, $\frac{1}{2}$ ABq, $J = 15.5\text{Hz}$); 2.829(1H, $\frac{1}{2}$ ABq, $J = 15.4\text{Hz}$); 2.835(1H, $\frac{1}{2}$ ABq, $J = 15.6\text{Hz}$); 3.06–3.09(2H, m); 3.45–3.49(2H, m); 3.67–3.85(4H, m); 3.90(2H, dd, $J = 3.4, 8.7\text{Hz}$); 4.09–4.18(2H, m); 6.03(2H, s, D_2O exch.); 6.78(1H, d, $J = 8.3\text{Hz}$); 6.79(1H, d, $J = 8.3\text{Hz}$); 6.89(2H, d, $J = 8.3\text{Hz}$); IR(neat) 3227, 2928, 1746, 1683, 1597, 1371, 1254, 1233, 1154, 1138, 1090, 836. MS (EI) 665 m/e (rel intensity) 665 (M^+ , 0.3) 565 (30.6) 521 (40.1) 164(100). EI HRMS m/e 665.38365 ($\text{C}_{37}\text{H}_{55}\text{N}_3\text{O}_6\text{Si}$ requires).




^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 0.089(6H, s); 0.11(6H, s); 0.87(9H, s); 0.88(9H, s); 1.13(3H, s); 1.15(3H, s); 1.36(3H, s); 1.37(3H, s); 1.46(6H, s); 1.59(9H, s); 1.61(9H, s); 1.86–2.06(10H, m); 2.09–2.20(6H, m); 2.61–2.70(2H, m); 2.747(1H, $\frac{1}{2}$ ABq, $J = 15.4\text{Hz}$); 2.754(1H, $\frac{1}{2}$ ABq, $J = 15.4\text{Hz}$); 2.30–3.05(2H, m); 3.05(6H, s); 3.14(2H, $\frac{1}{2}$ ABq, $J = 15.4\text{Hz}$); 3.39(2H, d, $J = 10.5\text{Hz}$); 3.74–3.85(2H,

m); 3.89–3.93(2H, m); 4.07–4.18(2H, m); 6.797(1H, d, $J = 8.3\text{Hz}$); 6.804(1H, d, $J = 8.3\text{Hz}$); 6.93(2H, d, $J = 8.3\text{Hz}$).

IR(neat) 2921, 1747, 1665, 1496, 1371, 1251, 1235, 1158, 1142, 1108, 1093, 837, 755.

MS (EI) 679 m/e (rel intensity) 679 (M^+ , 2.1) 579(4.2) 520(4.2) 178 (100).

EI HRMS m/e 679.4008 ($C_{38}H_{57}N_3O_6Si$ requires 679.4017).



12


^1H NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 1.21((3H, s); 1.23(3H, s); 1.29(2H, s); 1.32(3H, s); 1.42(3H, s); 1.45(3H, s); 1.54(6H, s); 1.88–2.00(10H, m); 2.07–2.22(6H, m); 2.63–2.72(2H, m); 2.79(1H, $\frac{1}{2}$ ABq, $J = 15.1\text{Hz}$); 2.80(1H, $\frac{1}{2}$ ABq, $J = 15.1\text{Hz}$); 3.01–3.07(4H, m, 2H D_2O exch.); 3.07(6H, s); 3.17(1H, $\frac{1}{2}$ ABq, $J = 15.1\text{Hz}$); 3.19(1H, $\frac{1}{2}$ ABq, $J = 15.4\text{Hz}$); 3.37–3.43(2H, m); 3.62(2H, br. s); 4.20(2H, dd, $J = 4.4, 12.3\text{Hz}$); 4.29(1H, dd, $J = 4.0, 12.3\text{Hz}$); 4.31(1H, dd, $J = 4.0, 12.3\text{Hz}$); 6.750(1H, d, $J = 8.4\text{Hz}$); 6.753(1H, d, $J = 8.3\text{Hz}$); 7.01(2H, d, $J = 8.4\text{Hz}$); 8.01(2H, s, D_2O exch.).

^{13}C NMR (300MHz) (CDCl_3) (mixture of two diastereomers) δ 14.0, 20.8, 22.6, 23.9, 24.4, 24.5, 24.7, 25.1, 27.7, 27.9, 30.2, 30.3, 31.3, 34.4, 45.9, 54.3, 57.4, 60.0, 60.2, 64.0, 71.0, 75.5, 76.6, 77.0, 77.4, 79.5, 104.6, 112.2, 116.17, 116.22, 125.0, 129.2, 137.2, 140.4, 141.6, 171.0, 174.3.


IR (neat) 3324, 2954, 1654, 1507, 1474, 1365, 1235, 1071, 1049, 908, 733.

MS (EI) 465 m/e (rel intensity) 465 (M^+ , 9.7) 406 (14.5) 287 (11.8) 178 (100).

EI HRMS m/e 465.2625 ($C_{27}H_{35}N_3O_4Si$ requires 465.2628).



12




13

^1H NMR (300MHz) (CDCl_3) mixture of two diastereomers δ TMS 0.80(3H, s); 0.83(3H, s); 1.08(3H, s); 1.10(3H, s); 1.22(3H, s); 1.26(3H, s); 1.50(3H, s); 1.52(3H, s); 1.40–1.60(8H, m); 1.77–1.93(8H, m); 2.05–2.21(2H, m); 2.55–2.71(4H, m); 3.02–3.10(4H, m); 3.06(6H, s); 3.63(4H, br s, 2H D_2O exch.); 4.05–4.24(4H, m); 6.60(1H, d, $J = 8.1\text{Hz}$); 6.62(1H, d, $J = 8.2\text{Hz}$); 6.78(1H, d, $J = 8.1\text{Hz}$); 6.79(1H, d, $J = 8.2\text{Hz}$); 7.42(1H, s, D_2O exch.); 7.45(1H, s, D_2O exch.)


IR (neat) 3333, 2974, 2933, 1703, 1651, 1646, 1631, 1456, 1395, 1323, 1200, 1046 903 cm^{-1}

MS (EI) 481 m/e (rel intensity) 481 (M^+ , 0.7) 422 (20.7) 421 (15) 135 (48).133 (100).

CI HRMS m/e 481 ($\text{C}_{27}\text{H}_{35}\text{N}_3\text{O}_5$ requires 481.2578), $[\text{M} + \text{H}]$ 482.2645 ($\text{C}_{27}\text{H}_{36}\text{N}_3\text{O}_5$ requires 482.2655).



13



(+)-Paraherquamide B

2

^1H NMR (300MHz) (CDCl_3) δ TMS 0.82(3H, s); 1.09(3H, s); 1.40(3H, s); 1.41(3H, s); 1.64(1H, dd, $J = 9.7, 12.4\text{Hz}$); 1.73–1.92(4H, m); 1.82(1H, $\frac{1}{2}$ ABq, $J = 15.5\text{Hz}$); 2.16(1H, dd, $J = 8.6, 17.8\text{Hz}$); 2.54–2.59(1H, m); 2.61(1H, $\frac{1}{2}$ ABq, $J = 11.1\text{Hz}$); 2.66(1H, $\frac{1}{2}$ ABq, $J = 15.5\text{Hz}$); 3.03–3.10(2H, m); 3.05(3H, s); 3.60(1H, $\frac{1}{2}$ ABq, $J =$

11.1Hz); 4.87(1H, d, J = 7.7Hz); 6.30(1H, d, J = 7.7Hz); 6.64(1H, d, J = 8.2Hz);
6.78(1H, d, J = 8.2Hz); 8.5(1H, br s, D₂O exch.).

¹³C NMR (300MHz) (CDCl₃) δ 20.7(q); 23.8(q); 26.2(q); 28.2(q); 28.8(t); 29.8(t);
29.9(q); 37.2(t); 46.1(s); 52.8(d); 53.8(t); 59.5(t); 63.0(s); 65.2(s); 67.4(s); 79.7(s);
115.0(d); 117.2(d); 120.3(d); 125.3(s); 132.5(s); 135.3(s); 139.0(d); 146.0(s); 172.9(s);
183.1(s).


IR (neat) 3190, 2974, 2933, 1703, 1697, 1651, 1631, 1503, 1456, 1328, 1195, 1046 728
cm⁻¹.

u.v. λ_{max} 226nm (ε = 30200).

[α]₂₅^D = + 0.4/7.75x10⁻³ = + 51.6(CHCl₃, c = 0.008)

MS (EI) 463 *m/e* (rel intensity) 463 (M⁺, 0.5) 404 (15.6) 135 (41.5) 133 (100).

EI HRMS *m/e* 463.2456 (C₂₇H₃₃N₃O₄ requires 463.2471).



CD Spectra (CH₂Cl₂)

J9324-m20