

CORRESPONDENCE

SARS-CoV-2 Infection among Travelers Returning from Wuhan, China

TO THE EDITOR: As severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infections (causing coronavirus disease 2019 [Covid-19]) spread globally, uncertainty surrounds estimates of the true number of infected persons, which is crucial to determining the severity of infection and the incidence of mild or asymptomatic cases and their possible transmission.¹ Modeling estimates suggest that in Wuhan, China, the city with the most Covid-19 cases, there are substantially




Figure 1. Follow-up of 94 Persons who Boarded an Evacuation Flight from Wuhan, China, to Singapore on January 30, 2020.

Of the 6 persons who had been transferred to a hospital before February 2, 2020, 3 returned to the government quarantine facility and are included with the 87 persons who underwent testing at the facility on quarantine day 6 (February 5, 2020). SARS-CoV-2 denotes severe acute respiratory syndrome coronavirus 2.

more cases than were officially reported, because milder cases may not have been captured in hospital-based surveillance.^{2,3} Data on travelers returning from areas with cases of Covid-19 could be useful in estimating its incidence.⁴

We followed up on 94 persons who boarded an evacuation flight from Wuhan to Singapore on January 30, 2020. Screening for body temperature was conducted at check-in and before boarding, and 3 febrile persons were prevented from boarding (Fig. 1); no additional information regarding the status of these 3 febrile persons was available. Surgical masks were provided to passengers on board the plane. On arrival in Singapore, the passengers underwent repeat screening for body temperature (fever was defined as a body temperature $\geq 38^{\circ}\text{C}$), and 2 persons (a woman 48 years of age [Patient 1] and a woman 47 years of age [Patient 2]) had a fever. The 2 febrile women were transferred immediately to a hospital, and they tested positive for SARS-CoV-2 (their clinical course is described in the Supplementary Appendix, available with the full text of this letter at NEJM.org).

The remaining 92 afebrile passengers (age range, 2 to 82 years) were quarantined for 14 days at a government quarantine facility, where they were checked for symptoms and fever three times daily. Six persons reported symptoms (4 on quarantine day 2 and 2 on quarantine day 3) and were placed in isolation in a hospital and underwent polymerase-chain-reaction (PCR) testing; all 6 persons tested negative for SARS-CoV-2. On quarantine day 3, samples from 76 of the 86 asymptomatic persons (75 nasopharyngeal swab samples and 1 nasal swab sample) were obtained and tested by means of PCR. A 17-year-old boy (the son of Patient 1) tested positive for Covid-19 and continued to have PCR-positive status for 2 weeks, and a 41-year-old man had an inconclusive result (positive for *N* gene and negative for *ORF1ab* gene). On quarantine day 6, samples from all 87 quarantined asymptomatic persons (85 nasopharyngeal swab samples and 2 nasal swab samples [3 of the 6 persons who had been transferred to the hospital before February 2 had returned to the government quarantine facility]) were obtained and tested; all tested negative. All persons who were not isolated in the hospital were released from quarantine on day 14, and all remained uninfected with Covid-19. Understanding the impli-

cations of transmission of SARS-CoV-2 infection from persons with asymptomatic or very mild symptomatic cases of Covid-19 is vital for the formulation of containment strategies.

Oon-Tek Ng, M.P.H.

Kalisvar Marimuthu, M.B., B.S.

Po-Ying Chia, M.B., B.S.

Vanessa Koh, Ph.D.

National Centre for Infectious Diseases
Singapore, Singapore
oon_tek_ng@ncid.sg
kalisvar_marimuthu@ncid.sg

Calvin J. Chiew, M.P.H.

Ministry of Health
Singapore, Singapore

Liang De Wang, M.Sc.

Barnaby E. Young, M.B., B.Chir.

Monica Chan, B.M., B.S.

Shawn Vasoo, M.B., B.S.

Li-Min Ling, M.B., B.S.

David C. Lye, M.B., B.S.

National Centre for Infectious Diseases
Singapore, Singapore

Kai-qian Kam, M.Med.(Ped.)

Koh-Cheng Thoon, M.Med.(Ped.)

KK Women's and Children's Hospital
Singapore, Singapore

Lalitha Kurupatham, M.P.H.

Zubaidah Said, M.P.H.

Ethan Goh, M.P.H.

Constance Low, M.P.H.

Soon-Kok Lim, M.P.H.

Pream Raj, M.P.H.

Olivia Oh, Ph.D.

Valerie T.J. Koh, M.P.H.

Cuiqin Poh, M.P.H.

Ministry of Health
Singapore, Singapore

Tze-Minn Mak, Ph.D.

Lin Cui, Ph.D.

National Centre for Infectious Diseases
Singapore, Singapore

Alex R. Cook, Ph.D.

Saw Swee Hock School of Public Health
Singapore, Singapore

Raymond T.P. Lin, M.B., B.S.

Yee-Sin Leo, M.P.H.

National Centre for Infectious Diseases
Singapore, Singapore
yee_sin_leo@ncid.sg

Vernon J.M. Lee, Ph.D.

Ministry of Health
Singapore, Singapore
vernon_lee@moh.gov.sg

Drs. Ng, Marimuthu, Leo, and Lee contributed equally to the study.

Supported by the following grants from the National Medical Research Council, Ministry of Health, Singapore: Collaborative Solutions Targeting Antimicrobial Resistance Threats in Health Systems (CoSTAR-HS) (NMRC CGAug16C005), NMRC Clinician Scientist Award (MOH-000276), and NMRC Clinician Scientist Individual Research Grant (MOH-CIRG18nov-0006).

Disclosure forms provided by the author are available with the full text of this letter at NEJM.org.

Any opinions, findings, conclusions, or recommendations expressed in this material are those of the authors and do not reflect the views of the Ministry of Health or the National Medical Research Council.

This article was published on March 12, 2020, at NEJM.org.

1. World Health Organization. Novel coronavirus (2019-nCoV): situation report - 12. February 2020 (https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200201-sitrep-12-ncov.pdf?sfvrsn=273c5d35_2).
2. Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet* 2020;395:689-97.
3. Imai N, Dorigatti I, Cori A, Riley S, Ferguson NM. Estimating the potential total number of novel Coronavirus cases in Wuhan City, China. London: Imperial College London, January 17, 2020 (<https://www.imperial.ac.uk/media/imperial-college/medicine/sph/ide/gida-fellowships/2019-nCoV-outbreak-report-17-01-2020.pdf>).
4. Hoehl S, Rabenau H, Berger A, et al. Evidence of SARS-CoV-2 infection in returning travelers from Wuhan, China. *N Engl J Med*. DOI: 10.1056/NEJMc2001899.

DOI: 10.1056/NEJMc2003100

Correspondence Copyright © 2020 Massachusetts Medical Society.