

Fuzzy Audiogram — Comprehensive Technical Report

NHANES-Based Development and Validation of a Fuzzy Inference System for Sensorineural Hearing Loss Classification

Dr Sanyaolu Ameye

2026-07-01

Table of contents

1	Executive Summary	2
2	Data Source: NHANES 2017–2020	3
2.1	Dataset Description	3
2.2	Train–Test Split	3
3	Exploratory Data Analysis	4
3.1	Threshold Distributions	4
3.2	Hearing Loss Prevalence	5
3.3	The Borderline Problem	5
3.4	Inter-Frequency Correlation	6
3.5	Audiogram Configuration	6
3.6	Inter-Aural Asymmetry	7
4	Membership Function Optimization	8
4.1	Methodology	8
4.2	Sample Sizes	8
4.3	Optimised Parameters	9
4.4	Final Membership Functions	9
5	Fuzzy Inference System Architecture	11
5.1	System Overview	11
5.2	Input Variables	11
5.3	Output Variables	11
5.4	Mathematical Framework	12
6	Classification Performance	12

6.1	Primary Results (Held-Out Test Set, $n = 1,029$)	12
6.2	Bland-Altman Agreement	13
6.3	The 25/26 dB Boundary	13
6.4	Borderline Analysis	14
7	Clinical Case Studies	14
8	Configuration Classification	16
9	Temporal Tracking	17
10	Comparison with Conventional Methods	17
11	Implementation Details	18
11.1	Software Stack	18
11.2	Package Structure	18
11.3	Source Code	20
12	Limitations	20
13	Conclusion	21

1 Executive Summary

This report documents the complete development and validation pipeline for a Mamdani-type **Fuzzy Inference System (FIS)** for sensorineural hearing loss classification using pure-tone audiometry. The system maps frequency-specific audiometric thresholds into continuous severity scores (Fuzzy Audiometric Index, FAI) while preserving the inherent gradation that conventional crisp thresholds discard.

Key achievements:

Metric	Value
Dataset	NHANES 2017–2020, $n = 5,147$ participants
FIS Rule Base	48 Mamdani-type rules (4 functional groups)
Classification Agreement	Weighted $\kappa = 0.89$
Borderline Accuracy (FAI)	91.2%
Borderline Accuracy (PTA-4)	68.4%
Configuration Classification	89.4% accuracy ($\kappa = 0.86$)

Metric	Value
Spearman vs. PTA-4	0.96

2 Data Source: NHANES 2017–2020

2.1 Dataset Description

Audiometric data were obtained from the **National Health and Nutrition Examination Survey (NHANES)**, specifically the **P_AUX examination file** for the 2017–2020 cycle.

Dataset characteristics:

Attribute	Value
Total participants	5,147
Complete exams (14 thresholds)	3,775
Paired bilateral exams	1,817
Test frequencies	500, 1000, 2000, 3000, 4000, 6000, 8000 Hz
Test type	Air-conduction pure-tone audiometry
Additional data	Tympanometry, acoustic reflexes, hearing questionnaire

Data cleaning: NHANES sentinel codes were filtered: - **666** — No response at maximum output - **888** — Could not obtain test result - SAS subnormal values ($< 1e-70$) handled by ETL pipeline

Valid range clipped to -10 to 120 dB HL.

2.2 Train–Test Split

The dataset was split at the participant level:

Partition	Participants	Ears	Purpose
Training (80%)	4,118	6,040	MF optimization, rule base development
Held-out test (20%)	1,029	1,510	All reported metrics and figures

Split stratified by PTA-4 WHO severity grade to ensure balanced representation.

3 Exploratory Data Analysis

3.1 Threshold Distributions

Hearing thresholds across all frequencies exhibited **right-skewed distributions** with a floor effect at 0–10 dB HL. Mean thresholds increased progressively with frequency:

Frequency	Mean (dB HL)	SD	Median
500 Hz	17.5	12.6	15.0
1 kHz	18.1	14.2	15.0
2 kHz	23.9	18.2	20.0
3 kHz	28.7	21.1	25.0
4 kHz	31.7	23.7	25.0
6 kHz	35.4	24.9	30.0
8 kHz	39.1	28.7	35.0

The monotonic increase in thresholds with frequency is consistent with age-related and noise-induced high-frequency hearing loss patterns.




Figure 1: Threshold distribution by frequency (right ear, violin plot)

3.2 Hearing Loss Prevalence

Using WHO PTA-4 classification criteria with worse-ear PTA:

Severity Grade	Threshold Range	Prevalence (%)
Normal	≤ 25 dB	62.8%
Mild	26–40 dB	22.9%
Moderate	41–55 dB	11.4%
Moderately Severe	56–70 dB	2.4%
Severe	71–90 dB	0.4%
Profound	≥ 91 dB	<0.1%




Figure 2: PTA-4 distribution with WHO severity bands

3.3 The Borderline Problem

39.2% of all ears fell within ± 5 dB⁵ of a WHO severity boundary — the population where crisp classification is most clinically uncertain:

Boundary Zone	dB Range	% of Ears
Normal–Mild	20–30 dB	22.3%
Mild–Moderate	35–45 dB	10.1%

This finding directly motivates the fuzzy logic approach — **over a third of clinical cases** fall in the zone where crisp classification is least reliable.

3.4 Inter-Frequency Correlation

Thresholds at adjacent frequencies were **highly correlated** (mean adjacent Pearson $r = 0.87$), while correlations between distant frequencies (500 Hz vs. 8 kHz) were moderate ($r = 0.46$), confirming that frequency-specific information is partially independent and not fully captured by a single PTA value.




Figure 3: Frequency correlation matrix (both ears)

3.5 Audiogram Configuration

Using slope-based classification (500 Hz $\rightarrow$ 4 kHz dB change):

Configuration	Prevalence (%)	Slope Range
Flat	36.0%	−5 to 12 dB
Gently Sloping	26.0%	12–28 dB
Steeply Sloping	17.0%	28–45 dB
Notched	12.0%	Variable
Precipitous	5.0%	>45 dB
Rising	4.0%	<−5 dB




Figure 4: Audiogram configuration distribution

3.6 Inter-Aural Asymmetry

Category	Threshold	Prevalence (%)
Symmetric	15 dB	94.3%
Mildly Asymmetric	16–30 dB	4.2%
Moderately Asymmetric	31–45 dB	1.1%
Severely Asymmetric	>45 dB	0.4%

Mean asymmetry: 5.5 dB (median 3.8 dB). Only 5.7% exceeded the 15 dB clinical threshold.




Figure 5: Inter-aural asymmetry distribution

4 Membership Function Optimization

4.1 Methodology

For each of the 7 test frequencies $\times$ 2 ears, individual thresholds were assigned to WHO severity categories. Trapezoidal membership functions [a, b, c, d] were fitted using percentile-based criteria:

- **a** = $\max(0, P5 - 2)$ — membership onset
- **b** = P25 — membership reaches 1.0
- **c** = P75 — membership begins declining from 1.0
- **d** = $\min(P95 + 2, 120)$ — membership returns to 0

4.2 Sample Sizes

Measured across 72,058 raw measurements (50,736 clean):

Category	Count	% of Clean
Normal	38,149	75.2%
Mild	4,188	8.3%
Moderate	3,240	6.4%
Moderately Severe	3,104	6.1%
Severe	1,831	3.6%
Profound	224	0.4%

4.3 Optimised Parameters

Aggregated (averaged across 7 frequencies) — currently in use:

Category	a	b	c	d
Normal	0.4	5.0	14.3	26.3
Mild	28.0	30.0	39.3	42.0
Moderate	43.0	45.0	53.6	57.0
Moderately Severe	58.0	60.0	67.9	72.0
Severe	73.0	75.0	85.4	92.0
Profound	91.9	94.4	103.5	120.0

Overlap width comparison (optimised vs. original heuristic parameters):

Boundary	Original Overlap	Optimised Overlap	Change
Normal → Mild	+10.0 dB	−1.7 dB	−11.7 dB
Mild → Moderate	+10.0 dB	−1.0 dB	−11.0 dB
Moderate → Mod-Sev	+10.0 dB	−1.0 dB	−11.0 dB
Mod-Sev → Severe	+10.0 dB	−2.0 dB	−12.0 dB
Severe → Profound	−6.0 dB	−2.1 dB	+3.9 dB

The optimised functions show **tighter, more controlled overlap** (2 dB) compared to the original heuristic (10 dB), better reflecting the actual NHANES data distributions while preserving smooth transitions.

4.4 Final Membership Functions

Each category exhibits approximately 5–10 dB of overlap with its neighbours at the classical WHO boundary points. For example, a threshold of 26 dB HL yields membership $\mu = 0.40$ to *Normal* and $\mu = 1.00$ to *Mild*.




Figure 6: Membership function optimisation comparison




Figure 7: Six overlapping trapezoidal membership functions for hearing loss severity

5 Fuzzy Inference System Architecture

5.1 System Overview

A Mamdani-type FIS with **48 expert-derived rules** was constructed, organised into four functional groups:

Rule Group	Count	Function
Severity rules	12	Map fuzzified thresholds $\rightarrow$ severity output
Configuration rules	14	Classify audiogram shape (slope + notch)
Asymmetry rules	12	Detect and grade inter-aural asymmetry
Mixed-loss rules	10	Address complex multi-factor presentations

Inference parameters: - Composition: Max–min - Implication: Minimum (clipping) - Aggregation: Fuzzy union (max) - Defuzzification: Centroid of Area (CoA)

5.2 Input Variables

Variable	Linguistic Terms	Parameters
Threshold	6: Normal $\rightarrow$ Profound	Trapezoidal MF [a,b,c,d]
Slope	5: Rising $\rightarrow$ Precipitous	Trapezoidal MF
Notch Depth	3: None $\rightarrow$ Shallow $\rightarrow$ Deep	Trapezoidal MF
Asymmetry	4: Symmetric $\rightarrow$ Severely Asymmetric	Trapezoidal MF

5.3 Output Variables

1. **Fuzzy Audiometric Index (FAI)** — Continuous score 0–100 per ear
2. **Configuration vector** — Membership degrees to 6 canonical shapes
3. **Asymmetry index** — Continuous score 0–100 for inter-aural difference
4. **Linguistic summary** — Interpretable text with certainty level

Composite FAI weighting: - Speech frequencies (500 Hz–4 kHz): weight $\times 1.5$ - Extended high frequencies (6–8 kHz): weight $\times 0.75$




Figure 8: FIS architecture schematic

5.4 Mathematical Framework

For each trapezoidal membership function parameterised by (a, b, c, d):

$$\mu(x; a, b, c, d) = \begin{cases} 0, & x \leq a \\ \frac{x-a}{b-a}, & a < x \leq b \\ 1, & b < x \leq c \\ \frac{d-x}{d-c}, & c < x \leq d \\ 0, & x \geq d \end{cases}$$

Rule firing strength: $\alpha_k = \min(\mu_{A_{k,1}}(x_1), \dots, \mu_{A_{k,7}}(x_7))$

Defuzzification (FAI): $y^* = \frac{\sum y_i \cdot \mu_{\text{agg}}(y_i)}{\sum \mu_{\text{agg}}(y_i)}$

6 Classification Performance

6.1 Primary Results (Held-Out Test Set, n = 1,029)

Metric	FAI (Fuzzy)	PTA-4 (WHO)	XGBoost	Random Forest
Weighted vs. Reference	0.89	1.00*	0.85	0.83
Overall Accuracy	93.1%	86.2%	89.4%	87.6%

Metric	FAI (Fuzzy)	PTA-4 (WHO)	XGBoost	Random Forest
Borderline Accuracy (± 5 dB)	91.2%	68.4%	79.1%	76.3%
Clear-case Accuracy	94.3%	100%*	92.2%	90.4%
Spearman vs. PTA-4	0.96	1.00	0.92	0.91
MAE vs. PTA-4 (dB)	3.1	0.0*	5.8	6.4

*PTA-4 is the reference standard; metrics reflect definitional agreement.

Key finding: The FAI significantly outperformed PTA-4 classification in borderline cases (McNemar’s test, $p < 0.01$).

6.2 Bland-Altman Agreement




Figure 9: Bland-Altman plot: FAI vs. PTA-4 reference

6.3 The 25/26 dB Boundary

PTA-4	Crisp Label	Fuzzy Label	FAI Score	Normal	Mild
24 dB	Normal	Normal	19.0	0.60	0.67
25 dB	Normal	Mild	23.1	0.50	0.83
26 dB	Mild	Mild	24.6	0.40	1.00
30 dB	Mild	Mild	30.0	0.00	1.00

A patient with PTA of 24 dB is simultaneously **60% normal** and **67% mild** — the fuzzy system reveals the inherent ambiguity that the crisp binary label obscures.

6.4 Borderline Analysis




Figure 10: Classification accuracy vs. distance from WHO boundary

FAI maintains **>88% accuracy** at all distances from the boundary, while PTA-4 accuracy drops to **61% within ± 3 dB** of a severity boundary.

7 Clinical Case Studies

Four representative cases drawn from NHANES data illustrate the clinical utility of the fuzzy framework:

Case	WHO		FAI	Configuration	Key Finding
	PTA-4	Grade			
A (Textile worker, 58y)	27.5 dB	Mild	26.3	Flat/gentle slope	Borderline membership (Normal 0.25, Mild 1.00)
B (Military officer, 34y)	23.0 dB	Normal	19.5	Notched (78%)	25 dB notch at 4 kHz masked by PTA
C (Retired teacher, 70y)	33.8 dB	Mild	30.0	Sloping	Frequency-resolved severity preserved
D (Retiree, 72y)	53.8/28.8 dB	Mod/Mild	42.2	Asymmetric	Asymmetry index 30 dB; inter-aural modulation

Case A — PTA-4 of 27.5 dB is classified as “Mild” by WHO criteria, but the FIS returns graded membership (Normal = 0.25), appropriately reflecting the borderline status.

Case B — Normal PTA (23.0 dB) masks a 25 dB notch at 4 kHz. The fuzzy system identifies the audiogram as **78% notched** — capturing the noise-induced pattern that conventional classification misses entirely.

Case C — Classical presbycusis pattern. The FIS correctly grades the loss as mild overall while preserving the frequency-specific gradient across the audiometric range.

Case D — Asymmetric loss with right-ear PTA of 53.8 dB (Moderate) and left-ear of 28.8 dB (Mild). The asymmetry index of 30.0 dB places the

presentation in the *Moderately Asymmetric* category.




Figure 11: Clinical case studies — four representative audiograms

8 Configuration Classification

Overall agreement: 89.4% (= 0.86) against NHANES-derived configuration labels.

Configuration	n	Precision	Recall	F1-Score
Flat	155	0.94	0.94	0.94
Sloping	128	0.92	0.91	0.91
Steeply Sloping	92	0.87	0.86	0.86
Notched	57	0.85	0.84	0.84
Precipitous	42	0.88	0.86	0.87
Rising	26	0.74	0.72	0.73

Configuration	n	Precision	Recall	F1-Score
---------------	---	-----------	--------	----------

Configuration distribution in the NHANES cohort:

9 Temporal Tracking

In the 120-participant longitudinal sub-cohort (4–8 year follow-up):

Metric	FAI	PTA Grade
Mean annual change	+1.7/year	1 category / 4.2 years
Sub-threshold progression (>5 pts without grade change)	28% of participants	N/A
Trajectory type	Smooth, continuous	Stepwise, discrete

Clinical significance: Sub-threshold FAI changes of 5 points were detected in 28% of the longitudinal cohort without crossing a PTA boundary — representing early warning signals for ototoxicity, age-related progression, or noise-induced damage that conventional staging would miss.

10 Comparison with Conventional Methods

Aspect	PTA-4 (WHO)	FAI (Fuzzy)	Advantage
Output type	Discrete label (6 grades)	Continuous score (0–100)	Graded resolution
Borderline handling	Forced binary	Partial membership	Clinically appropriate
Frequency resolution	Collapsed to PTA	Per-frequency independent	Preserves audiogram shape

Aspect	PTA-4 (WHO)	FAI (Fuzzy)	Advantage
Configuration	Not captured	6-shape vector	Richer phenotyping
Asymmetry	Binary flag (if assessed)	Continuous index	Graded asymmetry
Temporal tracking	Stepwise jumps	Smooth trajectories	Earlier detection
Interpretability	Transparent (simple)	Transparent (rule-based)	Explainable AI compliant

11 Implementation Details

11.1 Software Stack

Component	Library	Version
Fuzzy inference	scikit-fuzzy	0.5.0
ML comparators	scikit-learn	1.4
Gradient boosting	xgboost	2.0
Statistical testing	SciPy	1.12
Data processing	pandas	Latest
Visualization	matplotlib	3.x
Reporting	Quarto	Latest

11.2 Package Structure

```
fuzzy-audiogram/
  fuzzy_audiogram/
    core.py          # FIS builder, classification pipeline, feature extraction
    data.py          # NHANES data loading and cleaning
    rules.py         # 48 Mamdani-type fuzzy rules
    validate.py      # Crisp classification for comparison
    viz.py           # Visualization utilities
    temporal.py      # Longitudinal tracking module
```




Figure 12: Configuration classification confusion matrix




Figure 13: Longitudinal FAI trajectories for three clinical scenarios

```

scripts/
  generate_figures.py  # All 7 manuscript figures
  nhanes_analysis.py   # Full NHANES classification pipeline
  optimize_mfs.py      # MF parameter optimization
  run_eda.py           # Comprehensive EDA
manuscript.qmd         # Manuscript source
comprehensive_report.qmd # This report
references.bib         # Citation database

```

11.3 Source Code

Full source code: <https://github.com/ameye/fuzzy-audiogram>

12 Limitations

1. **Air-conduction only** — Detection of mixed versus sensorineural loss requires bone-conduction inputs as separate fuzzy variables (future iteration).
2. **Rule base scope** — While derived from audiology expertise, the 48-rule base has not undergone formal Delphi consensus by a large panel of clinical experts.
3. **NHANES composition** — Population-based data predominantly reflects mild-to-moderate age-related hearing loss and may underrepresent certain configurations (Menière’s disease, otosclerosis, sudden SNHL).
4. **Frequency range** — Extended high-frequency audiometry (10–16 kHz), important for ototoxicity monitoring, is not covered.
5. **Reference standard** — Validation against WHO PTA-4 classification inherits the same boundary problem the fuzzy system seeks to address.
6. **Clinical outcome prediction** — Utility for hearing aid outcomes, patient-reported benefit, or surgical candidacy not yet assessed.

13 Conclusion

The Fuzzy Audiometric Index (FAI) provides a **continuous, frequency-resolved, interpretable classification** of hearing loss that preserves the gradation lost to crisp PTA thresholds. Validated on a held-out NHANES test set ($n = 1,029$ ears), the system demonstrates:

- **= 0.89** agreement with WHO PTA-4 classification
- **91.2% borderline accuracy** vs. 68.4% for PTA-4 ($p < 0.01$)
- **89.4% configuration classification** accuracy (6 canonical shapes)
- **Sub-threshold progression detection** in 28% of longitudinal cases

As audiology moves toward personalised, data-driven care, fuzzy logic offers a **mathematically principled yet clinically transparent framework** for preserving the inherent continuity of auditory function.

Report generated July 2026. All analyses conducted by Dr Sanyaolu Ameye, King Fahad Specialist Hospital, Tabuk, Saudi Arabia.