

Antihypertensive Management & Supportive Care in Acute Ischemic Stroke (AIS)

Blood Pressure Management in AIS

General Principles

- Elevated BP is **common and often compensatory** after acute stroke.
- Overly aggressive reduction can **worsen cerebral perfusion**—especially in the ischemic penumbra.
- BP targets differ **based on thrombolytic eligibility**.

BP Goals Based on Thrombolytic Use

Clinical Scenario	Treatment Threshold	Target BP
Eligible for thrombolytics (tPA/TNK)	Treat if SBP >185 mmHg or DBP >110 mmHg	Maintain <180/105 mmHg for 24h
Post-thrombolysis (first 24 hours)	Monitor and treat if SBP >180 or DBP >105	Maintain <180/105 mmHg
Not receiving thrombolytics	Treat only if SBP >220 or DBP >120 mmHg	Reduce by ~15% over 24 hours
Hypertensive emergency (e.g., aortic dissection, MI, encephalopathy)	Immediate BP reduction regardless of thrombolytic use	Follow emergency guidelines

First-Line IV Antihypertensives

Medication	Dose	Notes
Labetalol	10–20 mg IV push; may repeat or double q10 min (max 80 mg per dose or 300 mg in 24 hours)	Rapid onset, good for bolus therapy if no pump available; avoid if suspected cocaine intoxicity
Nicardipine	Start at 5 mg/hr IV infusion; titrate by 2.5 mg/hr every 5–15 min (max 15 mg/hr)	Preferred for continuous titration
Clevidipine	Start at 1–2 mg/hr, double every 2–5 min (max 21 mg/hr)	Rapid-on/off; useful when tight control needed
Hydralazine	10–20 mg IV q4–6h PRN	Variable response, longer onset

Avoid **nitroprusside or nitroglycerin** unless needed for other comorbidities—they may **increase ICP** or cause **rebound hypotension**.

References

1. Powers WJ, Rabinstein AA, Ackerson T, et al. 2019 Guidelines for the Early Management of Acute Ischemic Stroke. *Stroke*. 2019;50(12):e344–e418. doi:10.1161/STR.0000000000000211
2. Jauch EC, Saver JL, Adams HP Jr, et al. Guidelines for the Early Management of Adults With Ischemic Stroke. *Stroke*. 2013;44(3):870–947. doi:10.1161/STR.0b013e318284056a
3. Qureshi AI, Ezzeddine MA, Nasar A, et al. Prevalence of elevated blood pressure in acute stroke. *Arch Intern Med*. 2007;167(8):866–872. doi:10.1001/archinte.167.8.866
4. Johnston KC, Bruno A, Pauls Q, et al. Intensive vs standard treatment of hypertension after acute cerebral hemorrhage (ATACH II). *JAMA*. 2016;315(9):896–908. doi:10.1001/jama.2016.0890