

# Tenecteplase (TNKase) Quick-Use Sheet – Acute Ischemic Stroke

Dose | Prep | Admin | PK | Clinical Considerations | Wastage Info

## Dosing Chart (IV Bolus Over 5 Seconds)

| Weight (kg) | Dose (mg) | Volume (mL at 5 mg/mL) |
|-------------|-----------|------------------------|
| < 60        | 15        | 3.0                    |
| 60–<70      | 17.5      | 3.5                    |
| 70–<80      | 20        | 4.0                    |
| 80–<90      | 22.5      | 4.5                    |
| ≥ 90        | 25        | 5.0                    |

Max dose in Acute Ischemic Stroke: 25 mg (5 mL)

Administer as single IV bolus over 5 seconds

## Vial Sizes, Reconstitution & Wastage

| Vial Size | Diluent Volume | Final Conc. | Max Dose Drawn | Wastage               |
|-----------|----------------|-------------|----------------|-----------------------|
| 25 mg     | 5.2 mL SWFI    | 5 mg/mL     | 5 mL (25 mg)   | Dose dependent        |
| 50 mg     | 10 mL SWFI     | 5 mg/mL     | 5 mL (25 mg)   | At least 5 mL (25 mg) |

Wastage Key Points:

- 25 mg vial: Minimally wasted, preferred for AIS use
- 50 mg vial: Significant wastage (at least 25 mg for **all AIS patients**)
- Do not pool excess drug from multiple vials
- Document discarded volume per institutional policy

## **Preparation & Administration**

### Reconstitution:

- Use only accompanying SWFI (5.2 mL for 25 mg; 10 mL for 50 mg)
- Swirl gently, do not shake
- Final concentration: 5 mg/mL

### Administration:

- Draw up appropriate dose based on patient weight
- Push IV bolus over 5 seconds
- Flush IV line with NS only (avoid D5W)

Because TNKase contains no antibacterial preservatives, reconstitute immediately before use. If not used immediately, refrigerate solution at 2°C to 8°C (35°F to 46°F) and use within 8 hours. DO NOT FREEZE. Final concentration of TNKase is 5 mg/mL.

## **Onset, Duration, and Half-Life**

Onset: Within minutes

Half-life: Biphasic

- Initial ~20 min
- Terminal ~90–130 min

Duration of action: ~2–4 hours

## **Clinical Considerations**

Guidelines: Class IIB recommendation (AHA/ASA 2019) as alteplase alternative

Contraindications:

- Refer to Inclusion/Exclusion Decision tree (linkout here?)

Monitoring:

- Neuro status q15 min for first 2 hours
- BP management & bleeding precautions
- No anticoagulants/antiplatelets for 24h

References:

<https://www.tnkase.com/content/dam/gene/tnkase/pdfs/tnkase-ais-dosing-card.pdf>

<https://www.tnkase.com/stemi.html>

[https://www.gene.com/download/pdf/tnkase\\_prescribing.pdf](https://www.gene.com/download/pdf/tnkase_prescribing.pdf)

