

# Modern Optimization Techniques

## 1. Theory

Lars Schmidt-Thieme

Information Systems and Machine Learning Lab (ISMLL)  
Institute for Computer Science  
University of Hildesheim, Germany

# Syllabus

Mon. 1.11.	(1)	0. Overview
		<b>1. Theory</b>
Mon. 8.11.	(2)	1. Convex Sets and Functions
		<b>2. Unconstrained Optimization</b>
Mon. 15.11.	(3)	2.1 Gradient Descent
Mon. 22.11.	(4)	2.2 Stochastic Gradient Descent
Mon. 29.11.	(5)	2.3 Newton's Method
Mon. 6.12.	(6)	2.4 Quasi-Newton Methods
Mon. 13.12.	(7)	2.5 Subgradient Methods
Mon. 20.12.	(8)	2.6 Coordinate Descent
	—	— <i>Christmas Break</i> —
		<b>3. Equality Constrained Optimization</b>
Mon. 3.1.	(9)	3.1 Duality
Mon. 10.1.	(10)	3.2 Methods
		<b>4. Inequality Constrained Optimization</b>
Mon. 17.1.	(11)	4.1 Primal Methods
Mon. 24.1.	(12)	4.2 Barrier and Penalty Methods
Mon. 31.1.	(13)	4.3 Cutting Plane Methods
Mon. 7.2.	(14)	Q & A

# Outline

1. Introduction
2. Convex Sets
3. Convex Functions
4. Recognizing Convex Functions

# Outline


1. Introduction

2. Convex Sets


3. Convex Functions

4. Recognizing Convex Functions

# A convex function



# A non-convex function



# Convex Optimization Problem

An **optimization problem**

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & h_q(x) \leq 0, \quad q = 1, \dots, Q \\ & Ax = b\end{array}$$

is said to be convex if  $f, h_1 \dots h_Q$  are convex.

Note: The equality constraints also are convex, even linear.

# Convex Optimization Problem

An **optimization problem**

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & h_q(x) \leq 0, \quad q = 1, \dots, Q \\ & Ax = b\end{array}$$

is said to be convex if  $f, h_1 \dots h_Q$  are convex.

How do we know if a function is convex or not?

Note: The equality constraints also are convex, even linear.



# Outline

1. Introduction

2. Convex Sets

3. Convex Functions

4. Recognizing Convex Functions

# Affine Sets

For any two points  $x_1, x_2$  we can define the line through them as:

# Affine Sets

For any two points  $x_1, x_2$  we can define the line through them as:

$$x = \theta x_1 + (1 - \theta)x_2, \quad \theta \in \mathbb{R}$$

# Affine Sets

For any two points  $x_1, x_2$  we can define the line through them as:

$$x = \theta x_1 + (1 - \theta)x_2, \quad \theta \in \mathbb{R}$$

Example:

$x_1$   
○


$x_2$   
○

# Affine Sets

For any two points  $x_1, x_2$  we can define the line through them as:

$$x = \theta x_1 + (1 - \theta)x_2, \quad \theta \in \mathbb{R}$$

Example:




# Affine Sets

For any two points  $x_1, x_2$  we can define the line through them as:

$$x = \theta x_1 + (1 - \theta)x_2, \quad \theta \in \mathbb{R}$$

Example:




# Affine Sets

For any two points  $x_1, x_2$  we can define the line through them as:

$$x = \theta x_1 + (1 - \theta)x_2, \quad \theta \in \mathbb{R}$$

Example:



# Affine Sets - Definition

An **affine set** is a set containing the line through any two distinct points in it.



# Affine Sets - Definition

An **affine set** is a set containing the line through any two distinct points in it.

## Examples:

►  $\mathbb{R}^N$  for  $N \in \mathbb{N}^+$

► solution set of linear equations  $X := \{x \in \mathbb{R}^N \mid Ax = b\}$

# Convex Sets

The **line segment** between any two points  $x_1, x_2$  is the set of all points:

# Convex Sets

The **line segment** between any two points  $x_1, x_2$  is the set of all points:

$$x = \theta x_1 + (1 - \theta)x_2, \quad 0 \leq \theta \leq 1$$

# Convex Sets

The **line segment** between any two points  $x_1, x_2$  is the set of all points:

$$x = \theta x_1 + (1 - \theta)x_2, \quad 0 \leq \theta \leq 1$$

Example:

# Convex Sets

The **line segment** between any two points  $x_1, x_2$  is the set of all points:

$$x = \theta x_1 + (1 - \theta)x_2, \quad 0 \leq \theta \leq 1$$

Example:

$x_1$   
○


$x_2$   
○

# Convex Sets

The **line segment** between any two points  $x_1, x_2$  is the set of all points:

$$x = \theta x_1 + (1 - \theta)x_2, \quad 0 \leq \theta \leq 1$$

Example:




# Convex Sets

The **line segment** between any two points  $x_1, x_2$  is the set of all points:

$$x = \theta x_1 + (1 - \theta)x_2, \quad 0 \leq \theta \leq 1$$

Example:




# Convex Sets

The **line segment** between any two points  $x_1, x_2$  is the set of all points:

$$x = \theta x_1 + (1 - \theta)x_2, \quad 0 \leq \theta \leq 1$$





Example:



A **convex set** contains the line segment between any two points in the set.





# Convex Sets - Examples: Which ones are Convex?





# Convex Sets - Examples

## Convex Sets:



## Non-convex Sets:



# Convex Sets - Examples

All affine sets are also convex:

- ▶  $\mathbb{R}^N$  for  $N \in \mathbb{N}^+$
- ▶ solution set of linear equations  $X := \{x \in \mathbb{R}^N \mid Ax = b\}$

Convex sets (but in general not affine sets):

- ▶ solution set of linear inequalities  $X := \{x \in \mathbb{R}^N \mid Ax \leq b\}$ 
  - ▶ half spaces, e.g.  $X := \{x \in \mathbb{R}^N \mid a^T x \leq b\}$   
e.g.,  $X := \{x \in \mathbb{R}^N \mid x_1 \geq 0\}$
  - ▶ convex polygons (2d) / polyhedrons (3d) / polytopes (nd)

# Convex Combination and Convex Hull

**(standard) simplex:**

$$\begin{aligned}\Delta^N &:= \{\theta \in \mathbb{R}^N \mid \theta_n \geq 0, n = 1, \dots, N; \sum_{n=1}^N \theta_n = 1\} \\ &= \{\theta \in [0, 1]^N \mid \mathbb{1}^T \theta = 1\}\end{aligned}$$

**convex combination** of some points  $x_1, \dots, x_N \in \mathbb{R}^M$ : any point  $x$  with

$$x = \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_N x_N, \quad \theta \in \Delta^N$$

**convex hull** of a set  $X \subseteq \mathbb{R}^M$  of points:

$$\text{conv}(X) := \{\theta_1 x_1 + \theta_2 x_2 + \dots + \theta_N x_N \mid N \in \mathbb{N}, x_1, \dots, x_N \in X, \theta \in \Delta^N\}$$

i.e., the set of all convex combinations of points in  $X$ .

Note:  $\mathbb{1} := (1, 1, \dots, 1)^T$  vector of all ones.

# Outline

1. Introduction

2. Convex Sets

3. Convex Functions

4. Recognizing Convex Functions

# Convex Functions

A function  $f : X \rightarrow \mathbb{R}$ ,  $X \subseteq \mathbb{R}^N$  is **convex** iff:

# Convex Functions

A function  $f : X \rightarrow \mathbb{R}$ ,  $X \subseteq \mathbb{R}^N$  is **convex** iff:

- ▶  $\text{dom } f := X$  is a convex set

# Convex Functions

A function  $f : X \rightarrow \mathbb{R}$ ,  $X \subseteq \mathbb{R}^N$  is **convex** iff:

- ▶  $\text{dom } f := X$  is a convex set
- ▶ for all  $x_1, x_2 \in \text{dom } f$  and  $0 \leq \theta \leq 1$  it satisfies

$$f(\theta x_1 + (1 - \theta)x_2) \leq \theta f(x_1) + (1 - \theta)f(x_2)$$

(the function is below of its secant segments/chords.)




# Convex Functions

A function  $f : X \rightarrow \mathbb{R}$ ,  $X \subseteq \mathbb{R}^N$  is **convex** iff:


- ▶  $\text{dom } f := X$  is a convex set
- ▶ for all  $x_1, x_2 \in \text{dom } f$  and  $0 \leq \theta \leq 1$  it satisfies

$$f(\theta x_1 + (1 - \theta)x_2) \leq \theta f(x_1) + (1 - \theta)f(x_2)$$


(the function is below of its secant segments/chords.)



# Convex functions




# Convex functions




►  $\theta x_1 + (1 - \theta)x_2$

# Convex functions



- ▶  $\theta x_1 + (1 - \theta)x_2$
- ▶  $(\theta x_1 + (1 - \theta)x_2, f(\theta x_1 + (1 - \theta)x_2))$

# Convex functions



- ▶  $\theta x_1 + (1 - \theta)x_2$
- ▶  $(\theta x_1 + (1 - \theta)x_2, f(\theta x_1 + (1 - \theta)x_2))$
- ▶  $(\theta x_1 + (1 - \theta)x_2, \theta f(x_1) + (1 - \theta)f(x_2))$

# How are Convex Functions Related to Convex Sets?

**epigraph** of a function  $f : X \rightarrow \mathbb{R}, X \subseteq \mathbb{R}^N$ :

$$\text{epi}(f) := \{(x, y) \in X \times \mathbb{R} \mid y \geq f(x)\}$$

# How are Convex Functions Related to Convex Sets?

**epigraph** of a function  $f : X \rightarrow \mathbb{R}, X \subseteq \mathbb{R}^N$ :

$$\text{epi}(f) := \{(x, y) \in X \times \mathbb{R} \mid y \geq f(x)\}$$

$f$  is convex (as function)  $\iff$   $\text{epi}(f)$  is convex (as set).

proof is straight-forward (try it!)

# Concave Functions

A function  $f$  is called **concave** if  $-f$  is convex




# Concave Functions

A function  $f$  is called **concave** if  $-f$  is convex

## A Concave Function

$f(x)$




# Concave Functions

A function  $f$  is called **concave** if  $-f$  is convex


## A Concave Function

$f(x)$



## A Convex Function

$f(x)$



# Strictly Convex Functions

A function  $f : X \rightarrow \mathbb{R}, X \subseteq \mathbb{R}^N$  is **strictly convex** if:

# Strictly Convex Functions

A function  $f : X \rightarrow \mathbb{R}, X \subseteq \mathbb{R}^N$  is **strictly convex** if:

- ▶  $\text{dom } f$  is a convex set

# Strictly Convex Functions

A function  $f : X \rightarrow \mathbb{R}, X \subseteq \mathbb{R}^N$  is **strictly convex** if:

- ▶  $\text{dom } f$  is a convex set
- ▶ for all  $x_1, x_2 \in \text{dom } f, x_1 \neq x_2$  and  $0 < \theta < 1$  it satisfies

$$f(\theta x_1 + (1 - \theta)x_2) < \theta f(x_1) + (1 - \theta)f(x_2)$$

# Examples

## **Examples of Convex functions:**

# Examples

## Examples of Convex functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$

# Examples

## Examples of Convex functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$
- ▶ exponential:  $f(x) = e^{ax}$ , with  $a \in \mathbb{R}$



# Examples

## Examples of Convex functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$
- ▶ exponential:  $f(x) = e^{ax}$ , with  $a \in \mathbb{R}$
- ▶ powers:  $f(x) = x^a$ , with  $\text{dom } f = \mathbb{R}_0^+$  and  $a \geq 1$  or  $a \leq 0$

# Examples

## Examples of Convex functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$
- ▶ exponential:  $f(x) = e^{ax}$ , with  $a \in \mathbb{R}$
- ▶ powers:  $f(x) = x^a$ , with  $\text{dom } f = \mathbb{R}_0^+$  and  $a \geq 1$  or  $a \leq 0$
- ▶ powers of absolute value:  $f(x) = |x|^a$ , with  $\text{dom } f = \mathbb{R}$  and  $a \geq 1$

# Examples

## Examples of Convex functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$
- ▶ exponential:  $f(x) = e^{ax}$ , with  $a \in \mathbb{R}$
- ▶ powers:  $f(x) = x^a$ , with  $\text{dom } f = \mathbb{R}_0^+$  and  $a \geq 1$  or  $a \leq 0$
- ▶ powers of absolute value:  $f(x) = |x|^a$ , with  $\text{dom } f = \mathbb{R}$  and  $a \geq 1$
- ▶ negative entropy:  $f(x) = x \log x$ , with  $\text{dom } f = \mathbb{R}^+$

# Examples

## Examples of Convex functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$
- ▶ exponential:  $f(x) = e^{ax}$ , with  $a \in \mathbb{R}$
- ▶ powers:  $f(x) = x^a$ , with  $\text{dom } f = \mathbb{R}_0^+$  and  $a \geq 1$  or  $a \leq 0$
- ▶ powers of absolute value:  $f(x) = |x|^a$ , with  $\text{dom } f = \mathbb{R}$  and  $a \geq 1$
- ▶ negative entropy:  $f(x) = x \log x$ , with  $\text{dom } f = \mathbb{R}^+$

## Examples of Concave Functions:

# Examples

## Examples of Convex functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$
- ▶ exponential:  $f(x) = e^{ax}$ , with  $a \in \mathbb{R}$
- ▶ powers:  $f(x) = x^a$ , with  $\text{dom } f = \mathbb{R}_0^+$  and  $a \geq 1$  or  $a \leq 0$
- ▶ powers of absolute value:  $f(x) = |x|^a$ , with  $\text{dom } f = \mathbb{R}$  and  $a \geq 1$
- ▶ negative entropy:  $f(x) = x \log x$ , with  $\text{dom } f = \mathbb{R}^+$

## Examples of Concave Functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$

# Examples

## Examples of Convex functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$
- ▶ exponential:  $f(x) = e^{ax}$ , with  $a \in \mathbb{R}$
- ▶ powers:  $f(x) = x^a$ , with  $\text{dom } f = \mathbb{R}_0^+$  and  $a \geq 1$  or  $a \leq 0$
- ▶ powers of absolute value:  $f(x) = |x|^a$ , with  $\text{dom } f = \mathbb{R}$  and  $a \geq 1$
- ▶ negative entropy:  $f(x) = x \log x$ , with  $\text{dom } f = \mathbb{R}^+$

## Examples of Concave Functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$
- ▶ powers/roots:  $f(x) = x^a$ , with  $\text{dom } f = \mathbb{R}_0^+$  and  $0 \leq a \leq 1$

# Examples

## Examples of Convex functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$
- ▶ exponential:  $f(x) = e^{ax}$ , with  $a \in \mathbb{R}$
- ▶ powers:  $f(x) = x^a$ , with  $\text{dom } f = \mathbb{R}_0^+$  and  $a \geq 1$  or  $a \leq 0$
- ▶ powers of absolute value:  $f(x) = |x|^a$ , with  $\text{dom } f = \mathbb{R}$  and  $a \geq 1$
- ▶ negative entropy:  $f(x) = x \log x$ , with  $\text{dom } f = \mathbb{R}^+$

## Examples of Concave Functions:

- ▶ affine:  $f(x) = ax + b$ , with  $\text{dom } f = \mathbb{R}$  and  $a, b \in \mathbb{R}$
- ▶ powers/roots:  $f(x) = x^a$ , with  $\text{dom } f = \mathbb{R}_0^+$  and  $0 \leq a \leq 1$
- ▶ logarithm:  $f(x) = \log x$ , with  $\text{dom } f = \mathbb{R}^+$

# Examples

## Examples of Convex functions:

All norms are convex!

- Immediate consequence of the triangle inequality and absolute homogeneity.

$$\|\theta x + (1 - \theta)y\| \leq \|\theta x\| + \|(1 - \theta)y\| = \theta\|x\| + (1 - \theta)\|y\|$$



# Examples

## Examples of Convex functions:

All norms are convex!

- ▶ Immediate consequence of the triangle inequality and absolute homogeneity.

$$\|\theta x + (1 - \theta)y\| \leq \|\theta x\| + \|(1 - \theta)y\| = \theta\|x\| + (1 - \theta)\|y\|$$

- ▶ For  $\mathbf{x} \in \mathbb{R}^N$ ,  $p \geq 1$ :

**p-norm:**  $\|\mathbf{x}\|_p := (\sum_{n=1}^N |x_n|^p)^{\frac{1}{p}},$

# Examples

## Examples of Convex functions:

All norms are convex!

- ▶ Immediate consequence of the triangle inequality and absolute homogeneity.

$$\|\theta x + (1 - \theta)y\| \leq \|\theta x\| + \|(1 - \theta)y\| = \theta\|x\| + (1 - \theta)\|y\|$$

- ▶ For  $\mathbf{x} \in \mathbb{R}^N$ ,  $p \geq 1$ :

**p-norm:**  $\|\mathbf{x}\|_p := (\sum_{n=1}^N |x_n|^p)^{\frac{1}{p}},$

- ▶  $\|\mathbf{x}\|_\infty := \max_{n=1:N} |x_n|$

# Examples

## Examples of Convex functions:

All norms are convex!

- ▶ Immediate consequence of the triangle inequality and absolute homogeneity.

$$\|\theta x + (1 - \theta)y\| \leq \|\theta x\| + \|(1 - \theta)y\| = \theta\|x\| + (1 - \theta)\|y\|$$

- ▶ For  $\mathbf{x} \in \mathbb{R}^N$ ,  $p \geq 1$ :

**p-norm:**  $\|\mathbf{x}\|_p := (\sum_{n=1}^N |x_n|^p)^{\frac{1}{p}},$

- ▶  $\|\mathbf{x}\|_\infty := \max_{n=1:N} |x_n|$

Affine functions on vectors are also convex:  $f(\mathbf{x}) = \mathbf{a}^T \mathbf{x} + b$

# Outline

1. Introduction

2. Convex Sets

3. Convex Functions

4. Recognizing Convex Functions

# 1st-Order Condition

$f$  is **differentiable** if  $\text{dom } f$  is open and the gradient

$$\nabla f(\mathbf{x}) = \left( \frac{\partial f(\mathbf{x})}{\partial x_1}, \frac{\partial f(\mathbf{x})}{\partial x_2}, \dots, \frac{\partial f(\mathbf{x})}{\partial x_n} \right)^T$$

exists everywhere.

**1st-order condition:** a differentiable function  $f$  is convex iff

# 1st-Order Condition

$f$  is **differentiable** if  $\text{dom } f$  is open and the gradient

$$\nabla f(\mathbf{x}) = \left( \frac{\partial f(\mathbf{x})}{\partial x_1}, \frac{\partial f(\mathbf{x})}{\partial x_2}, \dots, \frac{\partial f(\mathbf{x})}{\partial x_n} \right)^T$$

exists everywhere.

**1st-order condition:** a differentiable function  $f$  is convex iff

- $\text{dom } f$  is a convex set

# 1st-Order Condition

$f$  is **differentiable** if  $\text{dom } f$  is open and the gradient

$$\nabla f(\mathbf{x}) = \left( \frac{\partial f(\mathbf{x})}{\partial x_1}, \frac{\partial f(\mathbf{x})}{\partial x_2}, \dots, \frac{\partial f(\mathbf{x})}{\partial x_n} \right)^T$$

exists everywhere.

**1st-order condition:** a differentiable function  $f$  is convex iff

- ▶  $\text{dom } f$  is a convex set
- ▶ for all  $\mathbf{x}, \mathbf{y} \in \text{dom } f$

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x})$$


(the function is above any of its tangents.)

# 1st-Order Condition

**1st-order condition:** a differentiable function  $f$  is convex iff

- ▶  $\text{dom } f$  is a convex set
- ▶ for all  $\mathbf{x}, \mathbf{y} \in \text{dom } f$

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x})$$





# 1st-Order Condition / Proof

Let  $\text{dom } f = X$  be convex.

$$f : X \rightarrow \mathbb{R} \text{ convex} \Leftrightarrow f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}) \quad \forall \mathbf{x}, \mathbf{y}$$

# 1st-Order Condition / Proof

Let  $\text{dom } f = X$  be convex.

$$f : X \rightarrow \mathbb{R} \text{ convex} \Leftrightarrow f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}) \quad \forall \mathbf{x}, \mathbf{y}$$

$$\text{"} \Rightarrow \text{"} : f(x + t(y - x)) \leq (1 - t)f(x) + tf(y) \quad | : t$$

# 1st-Order Condition / Proof

Let  $\text{dom } f = X$  be convex.

$$f : X \rightarrow \mathbb{R} \text{ convex} \Leftrightarrow f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}) \quad \forall \mathbf{x}, \mathbf{y}$$

$$\text{"} \Rightarrow \text{"} : f(x + t(y - x)) \leq (1 - t)f(x) + tf(y) \quad | : t$$

$$f(y) \geq \frac{f(x + t(y - x)) - f(x)}{t} + f(x) \xrightarrow{t \rightarrow 0^+} \nabla f(x)^T (y - x) + f(x)$$

# 1st-Order Condition / Proof

Let  $\text{dom } f = X$  be convex.

$$f : X \rightarrow \mathbb{R} \text{ convex} \Leftrightarrow f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}) \quad \forall \mathbf{x}, \mathbf{y}$$

$$\text{"} \Rightarrow \text{"} : f(x + t(y - x)) \leq (1 - t)f(x) + tf(y) \quad | : t$$

$$f(y) \geq \frac{f(x + t(y - x)) - f(x)}{t} + f(x) \xrightarrow{t \rightarrow 0^+} \nabla f(x)^T (y - x) + f(x)$$

$$\text{"} \Leftarrow \text{"} : \text{Apply twice to } z := \theta x + (1 - \theta)y$$

$$f(x) \geq f(z) + \nabla f(z)^T (x - z)$$

$$f(y) \geq f(z) + \nabla f(z)^T (y - z)$$

# 1st-Order Condition / Proof

Let  $\text{dom } f = X$  be convex.

$$f : X \rightarrow \mathbb{R} \text{ convex} \Leftrightarrow f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}) \quad \forall \mathbf{x}, \mathbf{y}$$

$$\text{"} \Rightarrow \text{"} : f(\mathbf{x} + t(\mathbf{y} - \mathbf{x})) \leq (1 - t)f(\mathbf{x}) + tf(\mathbf{y}) \quad | : t$$

$$f(\mathbf{y}) \geq \frac{f(\mathbf{x} + t(\mathbf{y} - \mathbf{x})) - f(\mathbf{x})}{t} + f(\mathbf{x}) \xrightarrow{t \rightarrow 0^+} \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}) + f(\mathbf{x})$$

$$\text{"} \Leftarrow \text{"} : \text{Apply twice to } \mathbf{z} := \theta \mathbf{x} + (1 - \theta) \mathbf{y}$$

$$f(\mathbf{x}) \geq f(\mathbf{z}) + \nabla f(\mathbf{z})^T (\mathbf{x} - \mathbf{z})$$

$$f(\mathbf{y}) \geq f(\mathbf{z}) + \nabla f(\mathbf{z})^T (\mathbf{y} - \mathbf{z})$$

$$\begin{aligned} \leadsto \theta f(\mathbf{x}) + (1 - \theta)f(\mathbf{y}) &\geq f(\mathbf{z}) + \nabla f(\mathbf{z})^T (\theta \mathbf{x} + (1 - \theta) \mathbf{y}) - \nabla f(\mathbf{z})^T \mathbf{z} \\ &= f(\mathbf{z}) + \nabla f(\mathbf{z})^T \mathbf{z} - \nabla f(\mathbf{z})^T \mathbf{z} \\ &= f(\mathbf{z}) = f(\theta \mathbf{x} + (1 - \theta) \mathbf{y}) \end{aligned}$$

# 1st-Order Condition / Strict Variant

**strict 1st-order condition:** a differentiable function  $f$  is strictly convex iff

- ▶  $\text{dom } f$  is a convex set
- ▶ for all  $\mathbf{x}, \mathbf{y} \in \text{dom } f$

$$f(\mathbf{y}) > f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x})$$

Let  $\text{dom } f = X$  be convex.

$$f : X \rightarrow \mathbb{R} \text{ convex} \Leftrightarrow f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}) \quad \forall \mathbf{x}, \mathbf{y}$$

Q: What does this imply for points  $\mathbf{x}$  with  $\nabla f(\mathbf{x}) = 0$  ?

# Global Minima

Let  $\text{dom } f = X$  be convex.

$$f : X \rightarrow \mathbb{R} \text{ convex} \Leftrightarrow f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}) \quad \forall \mathbf{x}, \mathbf{y}$$

Consequence: Points  $x$  with  $\nabla f(x) = 0$  are (equivalent) global minima.

- ▶ minima form a convex set
- ▶ if  $f$  is strictly convex: there is exactly one global minimum  $x^*$ .



## 2nd-Order Condition

$f$  is **twice differentiable** if  $\text{dom } f$  is open and the Hessian  $\nabla^2 f(\mathbf{x})$

$$\nabla^2 f(\mathbf{x})_{n,m} = \frac{\partial^2 f(\mathbf{x})}{\partial x_n \partial x_m}$$

exists everywhere.

**2nd-order condition:** a twice differentiable function  $f$  is convex iff

## 2nd-Order Condition

$f$  is **twice differentiable** if  $\text{dom } f$  is open and the Hessian  $\nabla^2 f(\mathbf{x})$

$$\nabla^2 f(\mathbf{x})_{n,m} = \frac{\partial^2 f(\mathbf{x})}{\partial x_n \partial x_m}$$

exists everywhere.

**2nd-order condition:** a twice differentiable function  $f$  is convex iff

- $\text{dom } f$  is a convex set

## 2nd-Order Condition

$f$  is **twice differentiable** if  $\text{dom } f$  is open and the Hessian  $\nabla^2 f(\mathbf{x})$

$$\nabla^2 f(\mathbf{x})_{n,m} = \frac{\partial^2 f(\mathbf{x})}{\partial x_n \partial x_m}$$

exists everywhere.

**2nd-order condition:** a twice differentiable function  $f$  is convex iff

- ▶  $\text{dom } f$  is a convex set
- ▶ for all  $\mathbf{x} \in \text{dom } f$

$$\nabla^2 f(\mathbf{x}) \succeq 0 \quad \text{for all } \mathbf{x} \in \text{dom } f$$

## 2nd-Order Condition

$f$  is **twice differentiable** if  $\text{dom } f$  is open and the Hessian  $\nabla^2 f(\mathbf{x})$

$$\nabla^2 f(\mathbf{x})_{n,m} = \frac{\partial^2 f(\mathbf{x})}{\partial x_n \partial x_m}$$

exists everywhere.

**2nd-order condition:** a twice differentiable function  $f$  is convex iff

- ▶  $\text{dom } f$  is a convex set
- ▶ for all  $\mathbf{x} \in \text{dom } f$

$$\nabla^2 f(\mathbf{x}) \succeq 0 \quad \text{for all } \mathbf{x} \in \text{dom } f$$

Furthermore:

- ▶ for functions  $f$  on  $\text{dom } f \subseteq \mathbb{R}$  simply  $f''(x) \geq 0$  for all  $x \in \text{dom } f$
- ▶ if  $\nabla^2 f(\mathbf{x}) \succ 0$  for all  $\mathbf{x} \in \text{dom } f$ , then  $f$  is strictly convex
  - ▶ the converse is not true,  
e.g.,  $f(x) = x^4$  is strictly convex, but has 0 derivative at 0.

# Positive Semidefinite Matrices (Review)

A symmetric matrix  $A \in \mathbb{R}^{n \times n}$  is **positive semidefinite** ( $A \succeq 0$ ):

$$x^T A x \geq 0, \quad \forall x \in \mathbb{R}^N$$

Equivalent:

- (i) all eigenvalues of  $A$  are  $\geq 0$ .
- (ii)  $A = B^T B$  for some matrix  $B$

## Positive Semidefinite Matrices (Review)

A symmetric matrix  $A \in \mathbb{R}^{n \times n}$  is **positive semidefinite** ( $A \succeq 0$ ):

$$x^T A x \geq 0, \quad \forall x \in \mathbb{R}^N$$

Equivalent:

- (i) all eigenvalues of  $A$  are  $\geq 0$ .
- (ii)  $A = B^T B$  for some matrix  $B$

A symmetric matrix  $A \in \mathbb{R}^{N \times N}$  is **positive definite** ( $A \succ 0$ ):

$$x^T A x > 0, \quad \forall x \in \mathbb{R}^N \setminus \{0\}$$

Equivalent:

- (i) all eigenvalues of  $A$  are  $> 0$ .
- (ii)  $A = B^T B$  for some nonsingular matrix  $B$

# Quadratic Functions

$$f(x) := ax^2 + bx + c, \quad x, a, b, c \in \mathbb{R}$$

# Quadratic Functions

univariate / onedimensional:

$$f(x) := ax^2 + bx + c, \quad x, a, b, c \in \mathbb{R}$$

multivariate / multidimensional:

$$f(x) := x^T A x + b^T x + c, \quad x, b, c \in \mathbb{R}^N, \quad A \in \mathbb{R}^{N \times N}$$



# Recognizing Convex Functions

- ▶ There are a number of operations that preserve the convexity of a function.
- ▶ If  $f$  can be obtained by applying those operations to a convex function,  $f$  is also convex.

# Recognizing Convex Functions

- ▶ There are a number of operations that preserve the convexity of a function.
- ▶ If  $f$  can be obtained by applying those operations to a convex function,  $f$  is also convex.

## Nonnegative multiple:

- ▶ if  $f$  is convex and  $a \geq 0$  then  $af$  is convex.
- ▶ Example:  $5x^2$  is convex since  $x^2$  is convex

# Recognizing Convex Functions

- ▶ There are a number of operations that preserve the convexity of a function.
- ▶ If  $f$  can be obtained by applying those operations to a convex function,  $f$  is also convex.

## Nonnegative multiple:

- ▶ if  $f$  is convex and  $a \geq 0$  then  $af$  is convex.
- ▶ Example:  $5x^2$  is convex since  $x^2$  is convex

Q: Is the sum of two convex functions  $g$  and  $h$  convex again?

# Recognizing Convex Functions

- ▶ There are a number of operations that preserve the convexity of a function.
- ▶ If  $f$  can be obtained by applying those operations to a convex function,  $f$  is also convex.

## Nonnegative multiple:

- ▶ if  $f$  is convex and  $a \geq 0$  then  $af$  is convex.
- ▶ Example:  $5x^2$  is convex since  $x^2$  is convex

## Sum:

- ▶ if  $g$  and  $h$  are convex functions, then  $g + h$  is convex.
- ▶ Example:  $f(x) = e^x + x \log x$  with  $\text{dom } f = \mathbb{R}^+$  is convex since  $e^x$  and  $x \log x$  are convex

# Recognizing Convex Functions / Composition

## Composition of two convex functions:

- ▶ let  $g : \mathbb{R}^N \rightarrow \mathbb{R}$ ,  $h : \mathbb{R} \rightarrow \mathbb{R}$  be both convex and

$$f(x) := h(g(x))$$

- ▶ in general  $f$  is **not** convex
- ▶ counter example  $N = 1$ ,  $g(x) = h(x) = e^{-x}$ :

$$\begin{aligned}\left(e^{-e^{-x}}\right)'' &= \left(e^{-e^{-x}}(-e^{-x})(-1)\right)' \\ &= \left(e^{-e^{-x}}e^{-x}\right)' \\ &= e^{-e^{-x}}e^{-x}e^{-x} + e^{-e^{-x}}e^{-x}(-1) \\ &= e^{-e^{-x}}e^{-x}(e^{-x} - 1) < 0 \quad \text{for } x > 0\end{aligned}$$

# Recognizing Convex Functions / Composition

## **Composition with affine functions:**

- ▶ if  $f$  is convex then  $f(\mathbf{Ax} + \mathbf{b})$  is convex.

# Recognizing Convex Functions / Composition

## Composition with affine functions:

- ▶ if  $f$  is convex then  $f(A\mathbf{x} + \mathbf{b})$  is convex.
- ▶ Example: norm of an affine function  $\|A\mathbf{x} + \mathbf{b}\|$

# Recognizing Convex Functions / Composition

## Composition with nondecreasing functions:

► if  $g : \mathbb{R}^N \rightarrow \mathbb{R}$ ,  $h : \mathbb{R} \rightarrow \mathbb{R}$  and

$$f(\mathbf{x}) = h(g(\mathbf{x}))$$



# Recognizing Convex Functions / Composition

## Composition with nondecreasing functions:

- ▶ if  $g : \mathbb{R}^N \rightarrow \mathbb{R}$ ,  $h : \mathbb{R} \rightarrow \mathbb{R}$  and

$$f(\mathbf{x}) = h(g(\mathbf{x}))$$

- ▶  $f$  is convex if:
  - ▶  $g$  is convex,  $h$  is convex and nondecreasing *or*
  - ▶  $g$  is concave,  $h$  is convex and nonincreasing

# Recognizing Convex Functions / Composition

## Composition with nondecreasing functions:

- ▶ if  $g : \mathbb{R}^N \rightarrow \mathbb{R}$ ,  $h : \mathbb{R} \rightarrow \mathbb{R}$  and

$$f(\mathbf{x}) = h(g(\mathbf{x}))$$

- ▶  $f$  is convex if:
  - ▶  $g$  is convex,  $h$  is convex and nondecreasing *or*
  - ▶  $g$  is concave,  $h$  is convex and nonincreasing
- ▶ proof:

$$\begin{aligned}\nabla^2 h(g(\mathbf{x})) &= \nabla (h'(g(\mathbf{x})) \nabla g(\mathbf{x})) \\ &= h''(g(\mathbf{x})) \nabla g(\mathbf{x}) \nabla g(\mathbf{x})^T + h'(g(\mathbf{x})) \nabla^2 g(\mathbf{x})\end{aligned}$$

- ▶ Examples:
  - ▶  $e^{g(\mathbf{x})}$  is convex if  $g$  is convex
  - ▶  $\frac{1}{g(\mathbf{x})}$  is convex if  $g$  is concave and positive

# Recognizing Convex Functions

## Pointwise Maximum:

- ▶ if  $f_1, \dots, f_M$  are convex functions then  
 $f(\mathbf{x}) := \max\{f_1(\mathbf{x}), \dots, f_M(\mathbf{x})\}$  is convex.

# Recognizing Convex Functions

## Pointwise Maximum:

- ▶ if  $f_1, \dots, f_M$  are convex functions then  
 $f(\mathbf{x}) := \max\{f_1(\mathbf{x}), \dots, f_M(\mathbf{x})\}$  is convex.
- ▶ Example:  $f(\mathbf{x}) := \max_{m=1, \dots, M} (a_m^T \mathbf{x} + b_m)$  is convex

# Recognizing Convex Functions

There are many different ways to establish the convexity of a function:

1. Check the definition.

# Recognizing Convex Functions

There are many different ways to establish the convexity of a function:

1. Check the definition.
2. Show that  $\nabla^2 f(\mathbf{x}) \succeq 0$  for twice differentiable functions.

# Recognizing Convex Functions

There are many different ways to establish the convexity of a function:

1. Check the definition.
2. Show that  $\nabla^2 f(\mathbf{x}) \succeq 0$  for twice differentiable functions.
3. Show that  $f$  can be obtained from other convex functions by operations that preserve convexity.

# Summary

- ▶ **Convex sets** are closed under line segments (convex combinations).
- ▶ **Convex functions** are defined on a convex domain and
  - ▶ are below any of their secant segments / chords (definition),
  - ▶ are globally above their tangents (1st-order condition),
  - ▶ have a positive semidefinite Hessian (2nd-order condition).
- ▶ For convex functions, points with vanishing gradients are (equivalent) **global minima**.
- ▶ Operations that preserve convexity:
  - ▶ scaling with a nonnegative constant
  - ▶ sums
  - ▶ pointwise maximum
  - ▶ composition with an affine function
  - ▶ composition with a nondecreasing convex scalar function
  - ▶ composition of a noninc. convex scalar function with a concave funct.
    - ▶ esp.  $-g$  for a concave  $g$



# Further Readings

- ▶ Convex sets:
  - ▶ Boyd and Vandenberghe, 2004, chapter 2, esp. 2.1
  - ▶ see also ch. 2.2 and 2.3
- ▶ Convex functions:
  - ▶ Boyd and Vandenberghe, 2004, chapter 3, esp. 3.1.1–7, 3.2.1–5
- ▶ Convex optimization:
  - ▶ Boyd and Vandenberghe, 2004, chapter 4, esp. 4.1–3
  - ▶ see also ch. 4.4

# References



Boyd, Stephen and Lieven Vandenberghe (2004). *Convex Optimization*. Cambridge University Press.