

Molecular emission around low-mass protostars in the Serpens Main with IRAM 30m



A. Mirocha^{1,2}, A. Karska², L. E. Kristensen³, M. Gronowski⁴, M. Figueira⁵, M. Gładkowski^{2,6}, M. Żółtowski²

¹Astronomical Observatory, Jagiellonian University, ul. Orla 171, 30-244 Kraków, Poland; ²Centre for Astronomy, Faculty of Physics, Astronomy and Informatics, Nicolaus Copernicus University, ul. Grudziądzka 5, 87-100 Toruń, Poland; ³Centre for Star and Planet Formation, Niels Bohr Institute and Natural History Museum of Denmark, University of Copenhagen, Øster Voldgade 5-7, DK-1350 Copenhagen K, Denmark; ⁴Faculty of Physics, University of Warsaw, ul. Pasteura 5, 02-093 Warszawa, Poland; ⁵National Centre for Nuclear Research, ul. Pasteura 7, 02-093 Warszawa, Poland; ⁶Nicolaus Copernicus Astronomical Center, ul. Rabińska 8, 87-100 Toruń, Poland


Questions

- Where are molecules dissociated in molecular outflows?
- What are the typical UV fields in Class 0/I protostars?
- What is the influence of UV radiation on the surroundings of low-mass protostars?

Observations

- Spectra are obtained using IRAM 30m single dish antenna
- Targeted lines: HCN, CN, CS and their isotopologues

Serpens Main		
Molecule	Freq. [GHz]	Beam size [arcsec]
HCN 1-0	88.63	28
CN 1-0	113.49	22
CS 3-2	146.96	16
C ³⁴ S 3-2	144.62	16
H ¹³ CN 2-1	86.34	29
H ¹³ CN 1-0	172.68	14



**CN/HCN:
UV field tracer**


HCN $\xrightarrow{\text{UV}}$ CN + H

Conclusions


- CN/HCN ratio is higher around more evolved low-mass protostars
- N(CN)/N(HCN) ratio does not depend strongly on excitation conditions
- Nahoon astrochemical model shows that an additional UV radiation of a few percent of the average in the ISM is required
- The UV radiation cannot be neglected in models of star formation

Astrochemical model

- Nahoon code calculates abundances of 474 species
- In low temperature regime abundances of CN/HCN not depend strongly on temperature




SED analysis



Sbmm source	T_{bol} (K)	L_{bol} (L_{\odot})	Class
SMM9	46.14	11.69	Early Class 0
SMM1	40.35	108.72	Early Class 0
SMM5	148.24	4.49	Early Class I
SMM10	85.09	5.13	Late Class 0
SMM4	29.54	13.6	Early Class 0
SMM6	526.44	43.39	Late Class I
SMM12	100.87	6.68	Early Class I
SMM3	42.39	27.49	Early Class 0
SMM2	41.6	5.1	Early Class 0
SMM8		0.068 ^a	

CN/HCN column density ratio

- RADEX set of models
- Column density ratio covers the range of 1-10 irrespectively of the gas parameters



CN/HCN spatial distribution

Serpens CN J=1-0 divided by HCN J=1-0

