

Why a combination, nC_k , is always a natural number?
Different way to look at the answer accentuates factors
and prime numbers contained within a particular range of
1 to n .

Amit Kumar

July 25, 2024

1 Abstract

This article discusses why a combination nC_k will always be a natural number and never a fraction or an irrational number. This article separates an nC_k into selective, mutually exclusive and collectively exhaustive cases and then shows for each case that all the terms in a denominator have a corresponding multiple in the numerator. All k for a particular n for a nC_k are visualized through plots to gain better understanding. This different way of looking at a Combination nC_k , gives hindsights into factors and prime numbers contained within a particular range of 1 to n , in the context of the plots.

2 Introduction

From countless ages, it has been desired to know the number of ways an arrangement can be obtained or selected out of many possible ways. The quest has travelled from ancient times and has captured the imagination of the great mathematicians like Bhaskara-II, Blaise Pascal, Pierre-de-Fermat, Leonhard Euler to name a few.

Permutations and Combinations have become fundamental concepts with wide ranging applications in almost every sphere of academic studies or technical endeavors. Combination, nC_k represents number of ways k elements out of n can be chosen, *without any regard to order between the choosen items*. It is defined as ${}^nC_k = \frac{n!}{(n-k)! \times (k)!}$. While it is possible to show that each of the denominator terms $(n-k)!$ and $k!$ individually cancel terms in the numerator, it is not rigorously shown if the product of these two terms ever compete for a single numerator. If this condition is true where two terms in the denominator have a single term in the numerator, then the result will be a fraction and not a natural number. The results have indeed been observed to be integers empirically. However, surprisingly, a rigorous mathematical discussion on this question is relatively scarce. Some indirect proofs are definitely there such as using Pascal's triangle. The n 'th row of the Pascal's triangle is expansion of Binomial Coefficients, where each Coefficient is a combination (nC_k). It should be apparent that a rigorous proof of the observation that a combination will never be a fraction is crucial since many fundamental science concepts depend on it.

This paper aims to fill the gap and demonstrate that the final value of a combination will always collapse to an integer (more correctly a natural number). At the end of it we visualize our results and we find interesting connection of factors and prime numbers contained within the range 1 to n for a combination nC_1 .

3 Methods

3.1 Why permutation nP_k is a natural number?

Permutation is number of arrangements, when the order matters and is given by:

$${}^nP_k = \frac{n!}{(n-k)!} \text{ where } n, k \in \mathbb{N} \text{ and } 1 \leq k \leq n \quad (1)$$

e.g. Number of possibilities to choose 5 items out of 9 choices, we will write as ${}^9P_5 = \frac{9!}{(9-5)!}$. It is easy to show that $n!$ can be broken into multiplication two parts: one going from n to $(n-k+1)$ and other going from $(n-k)$ to 1. The denominator can also be written as multiplication of numbers from $(n-k)$ to 1 as below.

$$\begin{aligned} {}^nP_k &= \frac{n!}{(n-k)!} \\ &= \frac{(n) \times (n-1) \cdots \times (n-k+1) \times (n-k) \times (n-k-1) \cdots \times 3 \times 2 \times 1}{(n-k) \times (n-k-1) \times (n-k-2) \cdots \times 3 \times 2 \times 1} \\ &= (n) \times (n-1) \cdots \times (n-k+1) \end{aligned}$$

Since each participating number in the product is a natural number, the net product is also a natural number. ■

3.2 Why Combination nC_k is a natural number?

Permutation is number of arrangements, when the order *does not* matter and is given by:

$${}^nC_k = \frac{n!}{(n-k)! \times (k)!} \text{ where } n, k \in \mathbb{N} \text{ and } 1 \leq k \leq n \quad (2)$$

This time we can cancel the denominator's $k!$

$$\begin{aligned} {}^nC_k &= \frac{(n) \times (n-1) \cdots \times (k+1) \times (k) \times (k-1) \cdots \times 3 \times 2 \times 1}{(n-k) \times (n-k-1) \cdots \times 3 \times 2 \times 1 \times (k) \times (k-1) \cdots \times 3 \times 2 \times 1} \\ &= \frac{(n) \times (n-1) \cdots \times (k+1)}{(n-k) \times (n-k-1) \cdots \times 3 \times 2 \times 1} \end{aligned}$$

Thus it can be written down as:

$${}^nC_k = \frac{(k+1) \times (k+2) \cdots \times (n-1) \times (n)}{(n-k) \times (n-k-1) \cdots \times 3 \times 2 \times 1} \quad (3)$$

3.2.1 Different possible values of n and k w.r.t each other.

If $n//2$ is defined as integer portion of division of n by 2, then we can see that one of the two below must be true:

$$n = (n//2 \times 2) \text{ where } n \text{ is an even number} \quad (4)$$

$$n = (n//2 \times 2) + 1 \text{ where } n \text{ is an odd number} \quad (5)$$

On a similar note, k can be defined as six and six only possibilities w.r.t n :

$$k = n//2 \text{ where } n \text{ is an even number} \quad (6)$$

$$k = n//2 - l \text{ where } n \text{ is an even number and } l \in \mathbb{N}, 1 \leq l \leq n//2 \quad (7)$$

$$k = n//2 + m \text{ where } n \text{ is an even number and } m \in \mathbb{N}, 1 \leq m \leq n//2 \quad (8)$$

$$k = n//2 \text{ where } n \text{ is an odd number} \quad (9)$$

$$k = n//2 - l \text{ where } n \text{ is an odd number and } l \in \mathbb{N}, 1 \leq l \leq (n-1)/2 \quad (10)$$

$$k = n//2 + m \text{ where } n \text{ is an odd number and } m \in \mathbb{N}, 1 \leq m \leq (n+1)/2 \quad (11)$$

Let us take concrete examples to understand equations from (4) to (11).

$^{10}C_0$ means n is even, $k = 0$ and $l = n//2 = 5$: Thus equations (4) and (7) are realized.

$^{10}C_2$ means n is even, $k = 2$ and $l = 3$: Thus equations (4) and (7) are realized.

$^{10}C_4$ means n is even, $k = 4$ and $l = 1$: Thus equations (4) and (7) are realized.

$^{10}C_5$ means n is even and $k = n//2 = 5$: Thus equations (4) and (6) are realized.

$^{10}C_6$ means n is even, $k = 6$ and $m = 1$: Thus equations (4) and (8) are realized.

$^{10}C_8$ means n is even, $k = 8$ and $m = 3$: Thus equations (4) and (8) are realized.

$^{10}C_{10}$ means n is even, $k = 10$ and $m = n//2 = 5$: Thus equations (4) and (8) are realized.

$^{11}C_0$ means n is odd, $k = 0$ and $l = (n-1)/2 = 5$: Thus equations (5) and (10) are realized.

$^{11}C_2$ means n is odd, $k = 2$ and $l = 3$: Thus equations (5) and (10) are realized.

$^{11}C_4$ means n is odd, $k = 4$ and $l = 1$: Thus equations (5) and (10) are realized.

$^{11}C_5$ means n is odd and $k = n//2 = 5$: Thus equations (5) and (9) are realized.

$^{11}C_6$ means n is odd, $k = 6$ and $m = 1$: Thus equations (5) and (11) are realized.

$^{11}C_8$ means n is odd, $k = 8$ and $m = 3$: Thus equations (5) and (11) are realized.

$^{11}C_{11}$ means n is odd, $k = 11$ and $m = (n+1)/2 = 6$: Thus equations (5) and (11) are realized.

We can clearly see based on equations (6) to (11) that there are six possibilities that we separately need to take care and prove. In the following subsections, 3.2.3 to 3.2.8, we will prove each of the above condition.

3.2.2 Setting up the problem

Looking at equation (3) it is clear that each term in the denominator can be written as $(n-k-x)$, where $x \in \mathbb{W}, 0 \leq x \leq (n-k-1)$ and each term in the numerator can be written as $(k+y)$, where $y \in \mathbb{N}, 1 \leq y \leq (n-k)$. Formalizing in equations as below:

$$\text{denominator} = \prod_{x=0}^{n-k-1} (n-k-x), \text{ where } n \text{ and } k \text{ are constants and } x \text{ is the variable} \quad (12)$$

$$\text{numerator} = \prod_{y=1}^{n-k} (k+y), \text{ where } k \text{ is a constant and } y \text{ is the variable} \quad (13)$$

Now we have to show that for every $(n-k-x)$ and $(k+y)$, there exists a natural number a such that $a \times (n-k-x) = (k+y)$ holds true. i.e.

$$a \times (n-k-x) = (k+y), \text{ where } a \in \mathbb{N} \quad (14)$$

3.2.3 nC_k is always a natural number when equation (4) and (6) hold true

$$\begin{aligned}
a \times (n - k - x) &= (k + y) \\
a \times (2 \times n//2 - n//2 - x) &= (n//2 + y) \\
a \times (n//2 - x) &= (n//2 + y) \\
a &= \frac{(n//2 + y)}{(n//2 - x)} \\
y &= a \times (n//2 - x) - n//2 \\
y &= (a - 1) \times n//2 - a \times x
\end{aligned} \tag{15}$$

Let us take a couple of examples.

For ${}^{10}C_5$ and $x = 1$ we get the solution as below

$$\begin{aligned}
y &= (a - 1) \times 5 - a \\
y &= a \times (5 - 1) - 5 \\
y &= a \times 4 - 5
\end{aligned}$$

Clearly if $a = 2$ we get a valid solution and $y = 3$. Let us see what this means. This means that for denominator $(n - k - x) = (10 - 5 - 1) = 4$, we will get a corresponding numerator which will have its factor as $(k + y) = (5 + 3) = 8$. This is consistent. Let us take another example. For ${}^{10}C_5$ and $x = 3$ we get the solution as below

$$\begin{aligned}
y &= (a - 1) \times 5 - a \times 3 \\
y &= a \times (5 - 3) - 5 \\
y &= a \times 2 - 5
\end{aligned}$$

Clearly if $a = 3$ we get a valid solution and $y = 1$. Let us see what this means. This means that for denominator $(n - k - x) = (10 - 5 - 3) = 2$, we will get a corresponding numerator which will have its factor as $(k + y) = (5 + 1) = 6$. This is also consistent.

For (15), we know that $x \in \mathbb{W}, 0 \leq x \leq (n - k - 1)$ and $y \in \mathbb{N}, 1 \leq y \leq (n - k)$. These two conditions get simplified due to (6) as $0 \leq x \leq (n//2 - 1)$. Let us put extreme conditions of x to see if we can obtain a valid a for the denominator for every y .

$$\begin{aligned}
y &= (a - 1) \times n//2 - a \times 0 \text{ this is due to (15)} \\
y &= (a - 1) \times n//2
\end{aligned}$$

Clearly an a can be chosen such that R.H.S is a natural number. Similarly we can say that,

$$\begin{aligned}
y &= (a - 1) \times n//2 - a \times (n//2 - 1) \text{ this is due to (15)} \\
y &= \cancel{a \times n//2} - n//2 - \cancel{a \times n//2} + a \\
y &= a - n//2
\end{aligned}$$

Clearly an a can be chosen such that R.H.S is a natural number. Thus we saw that an a is available in the range of extreme x values so that we always have a so that every denominator is a factor for some numerator. ■

3.2.4 nC_k is always a natural number when equation (4) and (7) hold true

$$\begin{aligned}
a \times (n - k - x) &= (k + y) \\
a \times (2 \times n/2 - n/2 + l - x) &= (n/2 - l + y) \\
a &= \frac{(n/2 - l + y)}{(n/2 + l - x)} \\
y &= (a - 1) \times n/2 + l \times (a + 1) - a \times x \tag{16}
\end{aligned}$$

How do we interpret this? Let us see an example.

${}^{10}C_4$. Here $n = 10$ and thus $n/2 = 5, k = 4, l = n/2 - k = 5 - 4 = 1$. Furthermore let us assume that $x = 1$, i.e. the denominator is $(n - k - 1)$ for which the factor containing numerator has to be found.

Putting the above values in (16), we get $y = (a - 1)5 + 1(a + 1) - a \times 1 = (a - 1)5 + 1$

Clearly $a = 2$ will solve the equation which is consistent with restrictions on y and the value of y will be $y = (2 - 1) \times 5 + 1 = 6$. That means the corresponding multiples will reside in the numerator $(k + y) = (k + 6)$.

Clearly, putting all the values in equation (14) should get us no contradiction.

$$\begin{aligned}
a \times (n - k - x) &= (k + y) \\
2 \times (10 - 4 - 1) &= (4 + 6) \\
2 \times (5) &= (10) \\
10 &= 10
\end{aligned}$$

Please note that $a = 1$ will also give a consistent solution as $y = (1 - 1) \times 5 + 1 = 1$, which means $(k + 1)$ as the numerator. It can be seen that $(n - k - x) = (10 - 4 - 1) = 5$ will have two numerators where the multiples reside, one in $(k + 1) = 5$ and other in $(k + 6) = 10$. Let us take another example as ${}^{10}C_4$ and $x = 3$. i.e. the denominator is $(n - k - 3)$ and the corresponding numerator, $(k + y)$, has to be found. Using (16), we get $y = (a - 1)5 + 1(a + 1) - a \times 3 = (a - 1)5 + 6$. Again $a = 3$ will solve the equation and y will be $y = (3 - 1) \times 5 + 1 + 3 \times 3 = 2$. That means the corresponding factor will reside in the numerator $(k + y) = (k + 2)$.

. To see if it is consistent, we will have to put the values in (14).

$$\begin{aligned}
a \times (n - k - x) &= (k + y) \\
2 \times (10 - 4 - 3) &= (4 + 2) \\
2 \times (3) &= (6) \\
6 &= 6
\end{aligned}$$

Now according to (12),

$$\begin{aligned}
0 \leq x \leq (n - k - 1) &= 0 \leq x \leq (2 \times n/2 - n/2 + l - 1) \\
&= 0 \leq x \leq (n/2 + l - 1) \tag{17}
\end{aligned}$$

We will show that the equations are consistent at the extremes of x and therefore must be true for other values of x .

Choosing $x = 0$ in the equation (16), we get

$$\begin{aligned}
y &= (a - 1) \times n/2 + l \times (a + 1) - a \times x \\
y &= (a - 1) \times n/2 + l \times (a + 1) - a \times 0 \\
y &= (a - 1) \times n/2 + l \times (a + 1)
\end{aligned}$$

Since l is $a \in \mathbb{N}$, so the minimum value l can have is 1. Thus the second term in the R.H.S. above will be positive. To make y as positive integer, there is always a value of a we can choose to satisfy this restriction.

At the other extreme, choosing $x = (n//2 + l - 1)$, we get the below.

$$\begin{aligned} y &= (a - 1) \times n//2 + l \times (a + 1) - a \times (n//2 + l - 1) \\ y &= a \times n//2 - n//2 + l \times a + l - a \times n//2 - a \times l + a \\ y &= \cancel{a \times n//2} - n//2 + l \times \cancel{a} + l - \cancel{a \times n//2} - \cancel{a \times l} + a \\ y &= a - n//2 + l \end{aligned}$$

Again it is evident that an a can be chosen such that y can be in a form consistent with assumptions. Thus we have shown that the equation (16) is consistent and solvable at extremes of x and thus must be solvable between. ■

3.2.5 nC_k is always a natural number when equation (4) and (8) hold true

$$\begin{aligned} a \times (n - k - x) &= (k + y) \\ a \times (2 \times n//2 - n//2 - m - x) &= (n//2 + m + y) \\ a &= \frac{(n//2 + m + y)}{(n//2 - m - x)} \\ y &= (a - 1) \times n//2 - m \times (a + 1) - a \times x \end{aligned} \quad (18)$$

Interpreting this result is as before through an example.

${}^{10}C_6$. Here $n = 10$ and thus $n//2 = 5, k = 6, m = k - n//2 = 6 - 5 = 1$. Furthermore let us assume that $x = 1$, i.e. the denominator is $(n - k - 1)$ for which the factor containing numerator has to be found.

. Substituting the above values in (18), we get $y = (a - 1)5 - m(a + 1) - a \times 1 = (a - 1)5 - 1(1 + a) - a = (a - 1)5 - 2a - 1$

. It is easy to see that the equation is solved for $a = 3$. In such a case y will be equal to $(3 - 1) \times 5 - 2 \times 3 - 1 = 3$. i.e. for $(n - k - 1)$, there is a multiple in $(k + 3)$, where $n = 10$ and $k = 6$. It is trivial to verify this.

$$\begin{aligned} a \times (n - k - x) &= (k + y) \\ 3 \times (10 - 6 - 1) &= (6 + 3) \\ 3 \times (3) &= (9) \\ 9 &= 9 \end{aligned}$$

As per (12),

$$\begin{aligned} 0 \leq x \leq (n - k - 1) &= 0 \leq x \leq (2 \times n//2 - n//2 - m - 1) \\ &= 0 \leq x \leq (n//2 - m - 1) \end{aligned} \quad (19)$$

We need to show that there is no contradiction at the extremes (18) and thus it will hold true for middle values as well. Choosing $x = 0$ in equation (18) we arrive at,

$$\begin{aligned} y &= (a - 1) \times n//2 - m \times (a + 1) - a \times x \\ y &= (a - 1) \times n//2 - m \times (a + 1) - a \times 0 \\ y &= (a - 1) \times n//2 - m \times (a + 1) \end{aligned}$$

The above should hold for $y \geq 1$. Thus

$$\begin{aligned}(a-1) \times n//2 &\geq 1 + m \times (a+1) \\ a \times n//2 - m \times a &\geq (1 + m + n//2) \\ a &\geq \frac{(1 + m + n//2)}{(n//2 - m)}\end{aligned}$$

The last equation is defined as long as $n//2 \geq m$. This constraint is already in place due to the premise of (8). Thus a value of a can be computed such that (18) is satisfied. Working at the other extreme of $x = (n//2 - m - 1)$ we observe the below.

$$\begin{aligned}y &= (a-1) \times n//2 - m \times (1+a) - a \times (n//2 - m - 1) \\ y &= a \times n//2 - n//2 - m - m \times a - a \times n//2 - a \times m + a \\ y &= \cancel{a \times n//2} - n//2 - m - \cancel{m \times a} - \cancel{a \times n//2} - \cancel{a \times m} + a \\ y &= (a - n//2) - m\end{aligned}$$

As seen, y has allowed solutions at both extremes as there is always an a that can be chosen to satisfy the constraints. ■

3.2.6 nC_k is always a natural number when equation (5) and (9) hold true

$$\begin{aligned}a \times (n - k - x) &= (k + y) \\ a \times (2 \times n//2 + 1 - n//2 - x) &= (n//2 + y) \\ a &= \frac{(n//2 + y)}{(n//2 + 1 - x)} \\ y &= (a-1) \times n//2 + a - a \times x\end{aligned}\tag{20}$$

The best way to understand the above equation, as before, is through an example. In ${}^{11}C_5$, $n = 11$, $n//2 = 5 = k$. Let x be equal to 1, then substituting the values in the above equation, we get,

$$\begin{aligned}y &= (a-1) \times 5 + a - a \times 1 \\ y &= (a-1) \times 5 + \cancel{a} - \cancel{a \times 1} \\ y &= (a-1) \times 5\end{aligned}$$

Clearly $a = 2$ will give a solution and y will be equal to 5. That means for the above setup for $(n - k - 1)$, there will be a factor in $(k + 5)$. It is easy to verify this.

$$\begin{aligned}a \times (n - k - x) &= (k + y) \\ 2 \times (11 - 5 - 1) &= (5 + 5) \\ 2 \times (5) &= (10) \\ 10 &= 10\end{aligned}$$

Just like before we can appreciate that the equation (20) has boundaries for extreme values of x . Thus if we show the lower extreme and upper extreme values of x to have a consistent solution, then the proof must hold for values in between.

We remember that $0 \leq x \leq (n - k - 1) = 0 \leq x \leq (2 \times n//2 + 1 - n//2 - 1) = 0 \leq x \leq n//2$. First we will check for the lower limit of 0. Putting 0 in equation (20), we get,

$$\begin{aligned} y &= (a - 1) \times n//2 + a - a \times x \\ y &= (a - 1) \times n//2 + a - a \times 0 \\ y &= (a - 1) \times n//2 + a \end{aligned}$$

Clearly a value of a can be chosen such that R.H.S satisfies the constraint that $y \in \mathbb{N}$. When we think of the other extreme, we obtain the following.

$$\begin{aligned} y &= (a - 1) \times n//2 + a - a \times x \\ y &= (a - 1) \times n//2 + a - a \times (n - k - 1) \\ y &= (a - 1) \times n//2 + a - a \times (2 \times n//2 + 1 - n//2 - 1) \\ y &= (a - 1) \times n//2 + a - a \times (n//2) \\ y &= a - n//2 \end{aligned}$$

Here also it is clear that a value of a can be chosen such that the R.H.S is greater than or equal to 1.

Thus we have shown that a valid y can be obtained by choosing an appropriate a , for two extreme values of x . ■

3.2.7 nC_k is always a natural number when equation (5) and (10) hold true

$$\begin{aligned} a \times (n - k - x) &= (k + y) \\ a \times (2 \times n//2 + 1 - n//2 + l - x) &= (n//2 - l + y) \\ a &= \frac{(n//2 - l + y)}{(n//2 + 1 + l - x)} \\ y &= (a - 1) \times n//2 + l \times (a + 1) - a \times x + a \quad (21) \end{aligned}$$

Let us find the value for an example. Suppose we are given ${}^{11}C_4$. Here $n = 11$, $n//2 = 5$, $k = 4$, $l = 1$ and let's say $x = 1$. Then putting these values in the above equation, we get:

$$\begin{aligned} y &= (a - 1) \times n//2 + l \times (a + 1) - a \times x + a \\ y &= (a - 1) \times 5 + 1 \times (a + 1) - a \times 1 + a \\ y &= (a - 1) \times 5 + \cancel{1 \times a} + 1 - \cancel{a \times 1} + a \\ y &= (a - 1) \times 5 + 1 + a \end{aligned}$$

Clearly $a = 1$ gives a solution and thus $y = 2$. This argues that for $(n - k - 1)$, there is a factor in $(k + 2)$ for this example. i.e. for $(11 - 4 - 1)$ we have a factor in $(4 + 2)$. It is evident that 6 is on both sides of the equation.

Now we have to show that given an x and l and n , we can get a y such that $y \in \mathbb{N}$. We will show for two extreme values of x , it should hold true for values in between.

Let's recall that $0 \leq x \leq (n - k - 1) = 0 \leq x \leq (2 \times n//2 + 1 - n//2 + l - 1) = 0 \leq x \leq (n//2 + l)$. Starting with $x = 0$ and substituting $x = 0$ in equation (21).

$$\begin{aligned} y &= (a - 1) \times n//2 + l \times (a + 1) - a \times x + a \\ y &= (a - 1) \times n//2 + l \times (a + 1) - a \times 0 + a \\ y &= (a - 1) \times n//2 + l \times (a + 1) + a \end{aligned}$$

In the above each term on the right hand side is a positive integer and therefore the net value of y is also going to be a positive integer and thus the restriction that $1 \leq y, y \in \mathbb{N}$

Putting the other extreme value of x in equation (21)

$$\begin{aligned}
y &= (a-1) \times n//2 + l \times (a+1) - a \times x + a \\
y &= (a-1) \times n//2 + l \times (a+1) - a \times (n//2 + l) + a \\
y &= a \times n//2 - n//2 - l \times a + l - a \times n//2 - a \times l + a \\
y &= \cancel{a \times n//2} - n//2 - \cancel{l \times a} + l - \cancel{a \times n//2} - \cancel{a \times l} + a \\
y &= a - n//2 + l
\end{aligned}$$

Given the limits on l as per equation (10), a value of l can be found such that a consistent value of y can be obtained. ■

3.2.8 nC_k is always a natural number when equation (5) and (11) hold true

$$\begin{aligned}
a \times (n - k - x) &= (k + y) \\
a \times (2 \times n//2 + 1 - n//2 - m - x) &= (n//2 + m + y) \\
a &= \frac{(n//2 + m + y)}{(n//2 + 1 - m - x)} \\
y &= (a-1) \times n//2 - m \times (a+1) - a \times x + a \quad (22)
\end{aligned}$$

Let us see the equation (22) in action.

Suppose it is given that we have to work on ${}^{11}C_6$, then $n = 11, n//2 = 5, k = 6, m = 1$. Furthermore suppose $x = 3$, i.e. we have to find the multiple of $(n - k - 3)$ in the numerator.

$$\begin{aligned}
y &= (a-1) \times n//2 - m \times (a+1) - a \times x + a \\
y &= (a-1) \times 5 - 1 \times (a+1) - a \times 3 + a \\
y &= (a-1) \times 5 - 1 - 3 \times a \\
y &= 5 \times a - 5 - 1 - 3 \times a \\
y &= 2 \times a - 6
\end{aligned}$$

Clearly there is a solution for $a = 4$. Once we have obtained, a then we can plug it in equation (22) to obtain the value of y . We get,

$$\begin{aligned}
y &= (a-1) \times n//2 - m \times (a+1) - a \times x + a \\
y &= (4-1) \times 5 - 1 \times (4+1) - 4 \times 3 + 4 \\
y &= 15 - 5 - 12 + 4 \\
y &= 2
\end{aligned}$$

Thus for ${}^{11}C_6$, for the numerator $(n - k - 3)$, there is a factor in $(k + 2)$. It is easy to verify this using (14). Putting the values:

$$\begin{aligned}
4 \times (11 - 6 - 3) &= (6 + 2) \\
4 \times 2 &= 8
\end{aligned}$$

Thus we arrive at a consistent solution.

What is left to be shown is that given an x, m and n , we can always get a y such that $y \in \mathbb{N}$.

Since x has limits in equation (22), it suffices to show that a consistent y can be obtained always if consistent y can be obtained at the lower and upper extremes.

Now $0 \leq x \leq (n - k - 1) = 0 \leq x \leq (2 \times n//2 + 1 - n//2 - m - 1) = 0 \leq x \leq (n//2 - m)$. Given this premise, we first substitute the lower limit of x which is 0.

$$\begin{aligned} y &= (a - 1) \times n//2 - m \times (a + 1) - a \times x + a \\ y &= (a - 1) \times n//2 - m \times (a + 1) - a \times 0 + a \\ y &= (a - 1) \times n//2 - m \times (a + 1) + a \\ y &= a \times (n//2 - m + 1) - (n//2 + m) \end{aligned}$$

So as long as $a \times (n//2 - m + 1) - (n//2 + m) \geq 1$, the RHS will be a natural number and consistent y can be obtained.

$$\begin{aligned} a \times (n//2 - m + 1) - (n//2 + m) &\geq 1 \\ a \times (n//2 - m + 1) &\geq (n//2 + m + 1) \\ a &\geq \frac{(n//2 + m + 1)}{(n//2 - m + 1)} \end{aligned} \tag{23}$$

Thus for the above to hold true $(1 + n//2 - m) \geq 0$.

Due to constraints imposed by (3.2.8)

$$\begin{aligned} m &\leq (n + 1)/2 \\ m &\leq (2 \times n//2 + 1 + 1)/2 \\ m &\leq (2 \times n//2 + 2)/2 \\ m &\leq (n//2 + 1) \\ 0 &\leq (1 + n//2 - m) \end{aligned}$$

So the requirement is already satisfied for (23) and thus for $x = 0$ it is always possible to have a $y \in \mathbb{N}$. Now let us see the other extreme of x which is $(n//2 - m)$. Putting this value in equation (22).

$$\begin{aligned} y &= (a - 1) \times n//2 - m \times (a + 1) - a \times x + a \\ y &= (a - 1) \times n//2 - m \times (a + 1) - a \times (n//2 - m) + a \\ y &= a \times n//2 - n//2 - m \times a - m - a \times n//2 + a \times m + a \\ y &= \cancel{a \times n//2} - n//2 - \cancel{m \times a} - m - \cancel{a \times n//2} + \cancel{a \times m} + a \\ y &= (a - n//2) - m \end{aligned}$$

If $(a - n//2) - m \geq 1$, the RHS will always be a positive integer and a consistent y can be obtained.

$$\begin{aligned} (a - n//2) - m &\geq 1 \\ a &\geq (n//2 + m + 1) \end{aligned} \tag{24}$$

Since the R.H.S is a positive integer it is always possible to get such a value of a and thus y can be obtained for the other extreme of x , as well. ■

4 Conclusion

This article discusses about nC_k , where if n is even it can be written as $2 * n//2$ (where $n//2$ is the integer part of division of n by 2) while it can be stated as $2 * n//2 + 1$ if it is odd.

Furthermore k , can take only three values: $(n//2)$, $(n//2 - l)$ or $(n//2 + m)$, where l and m are positive integers greater than 1, and they denote *less* and *more* respectively.

The denominator is composed of product of $(n - k - x)$ terms, while the numerator is composed of products of $(k + y)$ terms, with the following constraints.

$$0 \leq x \leq (n - k - 1)$$

$$1 \leq y \leq (n - k)$$

With these premises we obtain the following relationship table between y and x (given n and k). Please note that l and m need not be supplied and can always be calculated if n and k are given. However l and m must be known/calculated before using any of the below equations. Also, in the below table, the equation must be solved for all possible values of a using the following algorithm.

1. Decide n is even or odd.
2. given k and n , decide if l or m is there and if yes what value.
3. Get range of x such that $0 \leq x \leq (n - k - 1)$
4. Solve for all values of a (for each x) using the below table such that $1 \leq y \leq (n - k)$.
5. For each a get a feasible and corresponding y .

n	k	y
Even	$n//2 \pm 0$ (i.e. $k = n//2$)	$y = (a - 1) \times n//2 - a \times x$
Even	$n//2 - l$ (i.e. $k < n//2$)	$y = (a - 1) \times n//2 + l \times (a + 1) - a \times x$
Even	$n//2 + m$ (i.e. $k > n//2$)	$y = (a - 1) \times n//2 - m \times (a + 1) - a \times x$
Odd	$n//2 \pm 0$ (i.e. $k = n//2$)	$y = (a - 1) \times n//2 + a - a \times x$
Odd	$n//2 - l$ (i.e. $k < n//2$)	$y = (a - 1) \times n//2 + a + l \times (a + 1) - a \times x$
Odd	$n//2 + m$ (i.e. $k > n//2$)	$y = (a - 1) \times n//2 + a - m \times (a + 1) - a \times x$

There can be an equivalent table where all the a terms are brought together. The below table can help find a quicker since all other terms are known. We can keep increasing a by 1 so that y stays in the range as per premise.

n	k	y
Even	$n//2$ (i.e. $k = n//2$)	$y = a \times (n//2 - x) - n//2$
Even	$n//2 - l$ (i.e. $k < n//2$)	$y = a \times (n//2 - x + l) - n//2 + l$
Even	$n//2 + m$ (i.e. $k > n//2$)	$y = a \times (n//2 - x - m) - n//2 - m$
Odd	$n//2$ (i.e. $k = n//2$)	$y = a \times (n//2 - x + 1) - n//2$
Odd	$n//2 - l$ (i.e. $k < n//2$)	$y = a \times (n//2 - x + l + 1) - n//2 + l$
Odd	$n//2 + m$ (i.e. $k > n//2$)	$y = a \times (n//2 - x - m + 1) - n//2 - m$

4.1 Visualizing the relationship between x and y as in 4

For (15), (16), (18), (20), (21) and (22), we can have plots, where for each plot x is on the abscissa while y is on ordinate for different values of nC_k (with implicit l or m if applicable).

The code for these plots (Figure 1-12) is given in appendix A.




Figure 1: Plot of 2C_k




Figure 2: Plot of 3C_k

Graph for various nC_k , where n is even




Figure 3: Plot of 4C_k

Graph for various nC_k , where n is odd




Figure 4: Plot of 5C_k

Graph for various nCk, where n is even




Figure 5: Plot of 6C_k

Graph for various nCk, where n is odd




Figure 6: Plot of 7C_k

Graph for various nC_k , where n is even




Figure 7: Plot of 8C_k

Graph for various nC_k , where n is odd




Figure 8: Plot of 9C_k

Graph for various nC_k , where n is even




Figure 9: Plot of $^{10}C_k$

Graph for various nC_k , where n is odd




Figure 10: Plot of $^{11}C_k$

Graph for various nCk, where n is even




Figure 11: Plot of $^{12}C_k$

Graph for various nCk, where n is odd




Figure 12: Plot of $^{13}C_k$

4.2 Delving deeper into the graphs

Based on the images seen before it is clear that the values of a for which an x has a corresponding y have a set of striking patterns if we analyze all k combinations of nC_k for a given n .

1. For a given n , graph is most dense for nC_1 and least dense for ${}^nC_{n-1}$. In other words, maximum number of a (and thus x and y) points are present on the graph in the former case and least points are present on the graph in the later case.
2. Vertical lines can be drawn for each x (which goes into the denominator term $(n - k - x)$).
3. For nC_1 , graph, the position of dots on the graph (a dot is a valid solution with a value of a), for any vertical line can be easily predicted. For the right most vertical line (equal to the value of $(n - 1 - 1)$) for x , the dots start at the value of 2 for the lowest value of $y = 1$ and then keep incrementing by 1 for every 1 position towards the top for a value of $(n - 1)$ corresponding to the $y = (n - 1)$. For the next left line (corresponding to the $(n - 2 - 1)$), the dots start at the value of 1 for the lowest value of $y = 2$ and then keep incrementing at every 2 position by 1. For the preceeding subsequent line at $(n - 3 - 1)$, the dots start at the value of 1 for the lowest value of $y = 3$ and then keep incrementing at every 3 position. This pattern holds for every preceeding line. For a line at position $(n - k - 1)$, the points start at the value of 1 for the lowest value of $y = k$ and then keep incrementing at every k location by 1.
4. Horizontal lines can be drawn for each y (which goes into the numerator term $(k + y)$).
5. For nC_1 , graph, the position of dots on the graph, for any horizontal line can be easily predicted. For a value of y we consider $y + 1$ and we find that the value of a for the dots are nothing but the factors of $y + 1$. Clearly prime numbers have no factors other than themselves and 1.
6. As we go towards higher k (starting from 1 and go towards $(n - 1)$) for the nC_k , the nC_1 graph shifts digonally (left and downwards) and terms outside the boundaries of $1 \leq y \leq (n - k)$ and $0 \leq x \leq (n - k - 1)$ keep 'falling off' the graph and 'disappear'. This we can call as the 'destruction picture' or 'low-to-high picture'. Please see the attached image at this URL https://github.com/amit7urmc/CombinationsAndPrimes/blob/master/25_C_k_264.mp4.
7. As we go from higher to lower k (starting from $(n - 1)$ and towards 1) for the nC_k , the nC_1 graph shifts digonally (right and upwards) and terms within the boundaries of $1 \leq y \leq (n - k)$ and $0 \leq x \leq (n - k - 1)$ keep 'joining' the graph and 'appear'. This we can call as the 'construction picture' or 'high-to-low picture'. Please see the attached image at this URL https://github.com/amit7urmc/CombinationsAndPrimes/blob/master/25_C_k_reverse_264.mp4.
8. In either of the pictures (destruction or construction), we can pause at any nC_k and we can calculate the value of combination as follows: All the numbers on the right most vertical lines go in the numerator and then in each vertical line highest point is taken as the divisor and its corresponding right most number as the dividend. The result of this goes into the numerator for each vertical line (except the right most line). e.g. for ${}^{25}C_{19}$, we can pause the construction/destruction picture there (or straightaway calculate the state using the code given in the appendix A) and we see that the right most vertical line consists of 25, 24, 23, 22, 21 and 20. There are 5 vertical lines (except the right most). For the first vertical line in the left, the highest vertical point is $a = 4$ and its corresponding right most point is $a = 24$. So the first vertical line 'collapses' to $24/4 = 6$. The subsequent line to




Figure 13: Distribution of x, y and a in the graph of ${}^{25}C_1$

```

| Even | n//2 | y = (a-1)\times n//2 - a\times x
| Even | n//2-1 | y = (a-1)\times n//2 + 1 \times (a+1) - a\times x
| Even | n//2+m | y = (a-1)\times n//2 - m \times (a+1) - a\times x
| Odd | n//2 | y = (a-1)\times n//2 + a - a\times x
| Odd | n//2-1 | y = (a-1)\times n//2 + a + 1\times(a+1) - a\times x
| Odd | n//2+m | y = (a-1)\times n//2 + a -m\times(a+1) - a\times x

```

The chart can be made in the following step.

1. Decide n is even or odd.
2. given k and n, decide if l or m is there and if yes what value.
3. Get range of x such that $0 \leq x \leq (n-k-1)$
4. Solve for a(s) and y(s) for each x with restrain that $1 \leq y \leq (n-k)$.
5. Return a list of three lists [x, y, a] # for plotting

```

"""
IS_EVEN = n%2 == 0
l, not_l_m, m = n//2-k if k < n//2 else None, n//2 if k==n//2 else None,
k-n//2 if k > n//2 else None

x = list(range(0,n-k,1))
y = []
solution_params = []
for each_x in x:
    a = []
    starting_guess = 1
    indiv_solution_params = []
    if IS_EVEN:
        if not_l_m:
            while True:
                y_ = (starting_guess-1)*n//2 - (starting_guess*each_x)
                if y_ > (2*(n//2)-k):
                    break
                if y_ >= 1:
                    a.append(starting_guess)
                    indiv_solution_params.append((l,not_l_m,m,
                                                    starting_guess
                                                    ))
                    starting_guess += 1
                y_s = list(map(lambda a_dash: (a_dash-1)*n//2 - (a_dash*
                                                                    each_x),a))
            y.append(y_s)
        elif l:
            while True:
                y_ = (starting_guess-1)*(n//2) + 1*(starting_guess+1) - (
                                                                    starting_guess*
                                                                    each_x)

                if y_ > (2*(n//2)-k):
                    break
                if y_ >= 1:
                    a.append(starting_guess)
                    indiv_solution_params.append((l,not_l_m,m,
                                                    starting_guess
                                                    ))
                    starting_guess += 1
                y_s = list(map(lambda a_dash: (a_dash-1)*(n//2) +1*(a_dash+1)
                                                                    - (a_dash*each_x),a))

```

```

        y.append(y_s)
    elif m:
        while True:
            y_ = (starting_guess-1)*(n//2) -m*(starting_guess+1) - (
                starting_guess*
                each_x)

            if y_ > (2*(n//2)-k):
                break
            if y_ >= 1:
                a.append(starting_guess)
                indiv_solution_params.append((l,not_l_m,m,
                    starting_guess
                ))

            starting_guess += 1
            y_s = list(map(lambda a_dash: (a_dash-1)*(n//2) -m*(a_dash+1)
                - (a_dash*each_x),a))
            y.append(y_s)
    else:
        if not_l_m:
            while True:
                y_ = (starting_guess-1)*(n//2) +starting_guess - (
                    starting_guess*
                    each_x)

                if y_ > ((2*(n//2)+1)-k):
                    break
                if y_ >= 1:
                    a.append(starting_guess)
                    indiv_solution_params.append((l,not_l_m,m,
                        starting_guess
                    ))

                starting_guess += 1
                y_s = list(map(lambda a_dash: (a_dash-1)*(n//2)+a_dash - (
                    a_dash*each_x),a))
                y.append(y_s)
            elif l:
                while True:
                    y_ = (starting_guess-1)*(n//2) +starting_guess + l*(
                        starting_guess+1)
                        - (starting_guess*
                        each_x)

                    if y_ > ((2*(n//2)+1)-k):
                        break
                    if y_ >= 1:
                        a.append(starting_guess)
                        indiv_solution_params.append((l,not_l_m,m,
                            starting_guess
                        ))

                    starting_guess += 1
                    y_s = list(map(lambda a_dash: (a_dash-1)*(n//2)+a_dash +l*(
                        a_dash+1) - (a_dash*
                        each_x),a))
                    y.append(y_s)
            elif m:
                while True:
                    y_ = (starting_guess-1)*(n//2) +starting_guess -m*(
                        starting_guess+1)
                        - (starting_guess*
                        each_x)

                    if y_ > ((2*(n//2)+1)-k):
                        break
                    if y_ >= 1:

```

```

        a.append(starting_guess)
        indiv_solution_params.append((l,not_l_m,m,
                                      starting_guess
                                      ))

        starting_guess += 1
        y_s = list(map(lambda a_dash: (a_dash-1)*(n//2)+a_dash -m*(
                                      a_dash+1) - (a_dash*
                                      each_x),a))

        y.append(y_s)
        solution_params.append(indiv_solution_params)
    returned_x = []
    returned_y = []
    returned_params = []
    for each_y_list_index, each_y_list in enumerate(y):
        solution_params_ = solution_params[each_y_list_index]
        for each_y_index, each_y in enumerate(each_y_list):
            returned_x.append(x[each_y_list_index])
            returned_y.append(each_y)
            returned_params.append(solution_params_[each_y_index])
    returned_results = [returned_x, returned_y, returned_params]
    return returned_results

def create_one_case_graph(n: int, k: int)->None:
    max = n
    fig, ax = plt.subplots(nrows=1, ncols=1, figsize=(12,12))
    ax.set_xlim(right=n+1)
    ax.set_xlim(left=-1)
    ax.set_ylim(top=n+1)
    ax.set_ylim(bottom=0)
    ax.set_xticks(ticks=list(range(1,n-k,1)), labels=[str(i) for i in range(1
                                                    ,n-k,1)])
    ax.yaxis.set_major_locator(MaxNLocator(integer=True, min_n_ticks=1))
    ax.xaxis.set_major_locator(MaxNLocator(integer=True, min_n_ticks=1))
    x, y, params = create_chart_combination(n,k)
    ax.scatter(x=x, y=y, c=[[0,0,0]], marker='o')
    ax.grid(which='both')
    ax.set_title('$ ^{d}C_{d}$'%(n,k))
    for data_index, _ in enumerate(params):
        param = _
        annotation = f"a={param[3]}"
        xy = (x[data_index], y[data_index])
        ax.annotate(annotation, xy=xy, xytext=(10,-20),textcoords='offset
                                                    pixels')
    ax.set_xlabel('x as in denominator (n-k-x)')
    ax.set_ylabel('y as in numerator (k+y)')
    fig.savefig(f'./{n}_{k}_alone.png')
    plt.close()

def create_all_cases_graph(n: list[int, int])->None:
    """
    Supplied one even and one add number. 0th index should be even while 1st
    index should be odd.
    """
    assert n[0]%2==0 and n[1]%2
    markers = ["o","v","1","s","p","P","*","H","D","X"]
    fig, axs1 = plt.subplots(nrows=int(ceil(sqrt(n[0]))),ncols=int(ceil(sqrt(
        n[0]))), sharey=True, sharex=True)

    for axs_row in axs1:
        for axs_col in axs_row:
            axs_col.yaxis.set_major_locator(MaxNLocator(integer=True,
                                                         min_n_ticks=1))

```

```

        axs_col.xaxis.set_major_locator(MaxNLocator(integer=True,
                                                    min_n_ticks=1))

## Draw for even numbers
counter = 0
for k in range(1,n[0],1):
    if(k%int(ceil(sqrt(n[0])))==0):
        counter += 1
    x, y, params = create_chart_combination(n[0],k)
    axs1[counter][k%int(ceil(sqrt(n[0])))].scatter(x=x, y=y, c=[[0,0,0]],
                                                    marker=markers[k%(len(markers)
                                                    -1)])
    axs1[counter][k%int(ceil(sqrt(n[0])))].grid(which='both')
    axs1[counter][k%int(ceil(sqrt(n[0])))].set_title('$ ~{d}C_{d}$'%(n[
0],k))

    for data_index, _ in enumerate(params):
        param = _
        annotation = f"a={param[3]}"
        xy = (x[data_index], y[data_index])
        axs1[counter][k%int(ceil(sqrt(n[0])))].annotate(annotation, xy=xy
                                                    , xytext=(10,-10),
                                                    textcoords='offset pixels'
                                                    )

fig.suptitle("Graph for various nCk, where n is even")
fig.supxlabel("x as in denominator (n-k-x)")
fig.supylabel("y as in numerator (k+y)")

fig, axs2 = plt.subplots(nrows=int(ceil(sqrt(n[1]))),ncols=int(ceil(sqrt(
n[1]))), sharey=True, sharex=True)

for axs_row in axs2:
    for axs_col in axs_row:
        axs_col.yaxis.set_major_locator(MaxNLocator(integer=True,
                                                    min_n_ticks=1))

        axs_col.xaxis.set_major_locator(MaxNLocator(integer=True,
                                                    min_n_ticks=1))

## Draw for odd numbers
counter = 0
for k in range(1,n[1],1):
    if(k%int(ceil(sqrt(n[1])))==0):
        counter += 1
    x, y, params = create_chart_combination(n[1],k)
    axs2[counter][k%int(ceil(sqrt(n[1])))].scatter(x=x, y=y, c=[[0,0,0]],
                                                    marker=markers[k%(len(markers)
                                                    -1)])
    axs2[counter][k%int(ceil(sqrt(n[1])))].grid(which='both')
    axs2[counter][k%int(ceil(sqrt(n[1])))].set_title('$ ~{d}C_{d}$'%(n[
1],k))

    for data_index, _ in enumerate(params):
        param = _
        annotation = f"a={param[3]}"
        xy = (x[data_index], y[data_index])
        axs2[counter][k%int(ceil(sqrt(n[1])))].annotate(annotation, xy=xy
                                                    , xytext=(10,-10),
                                                    textcoords='offset pixels'
                                                    )

fig.suptitle("Graph for various nCk, where n is odd")
fig.supxlabel("x as in denominator (n-k-x)")
fig.supylabel("y as in numerator (k+y)")

plt.show()

```



```

if __name__ == "__main__":
    create_all_cases_graph([2,3])
    create_all_cases_graph([4,5])
    create_all_cases_graph([6,7])
    create_all_cases_graph([8,9])
    create_all_cases_graph([10,11])
    create_all_cases_graph([12,13])

n=25
for k in range(1,n,1):
    create_one_case_graph(n, k)

```

B Code to generate the prime numbers as in 4

```

class generatePrime:

    def __init__(self, fileToWrite: str="./primeListGenerated.txt")-> None:
        """
        A list of prime numbers discovered and their current steps.
        To save computing the length in every iteration the list length is
            also maintained.
        fileToWrite is a file where generated prime numbers are saved.
        """
        self.primeNumbersDiscovered = [2]
        self.runningNumber = 2
        self.primeNumbersDiscoveredListLength = 1
        self.fileToWrite = fileToWrite

    def yield_prime(self, upto_n:int)->None:
        """
        upto_n denotes how many prime numbers need to be written to the file.
        """
        with open(self.fileToWrite,"w") as fileHandle:
            fileHandle.write("primeNumbersGenerated")
            fileHandle.write("\n2")
            counter = 0
            while True:
                skipFlag = False
                self.runningNumber += 1
                for i in range(self.primeNumbersDiscoveredListLength):
                    if not self.runningNumber%self.primeNumbersDiscovered[i]:
                        skipFlag = True
                        break

                if not skipFlag:
                    self.primeNumbersDiscovered.append(self.runningNumber)
                    self.primeNumbersDiscoveredListLength += 1
                    fileHandle.write(f"\n{self.runningNumber}")
                    counter += 1
                if counter >= upto_n-1:
                    break

    def compare_against_known_result(self, know_result_file:str="./
        PrimeNumbersTop1000.txt")->bool:
        """
        A known result file is taken as input.

```

```

Care must be taken that it has at least same number of primes as the
upto_n parameter in the
yield_prime method.

"""
present_absent = None
with open(know_result_file, "r") as knownHandle, open(self.
                                                    fileToWrite,"r") as
                                                    calculateddHandle:

    knownResult = knownHandle.readlines()[1:]
    calculatedResult = calculateddHandle.readlines()[1:]
    assert len(knownResult) == len(calculatedResult)
    present_absent = [True if knownResult[i]==calculatedResult[i]
                      else False for i in range(
len(knownResult))]

if all(present_absent):
    return True
return False

if __name__ == "__main__":
    upto_n = 1000
    full_match_str = "full_match"
    some_mismatch_str = "some_mismatch"
    primeObj = generatePrime()
    primeObj.yield_prime(upto_n)
    full_match = primeObj.compare_against_known_result()
    print(f"We tried to match first known {upto_n} primes with those
generated by this program. The
answer is {full_match_str if
full_match else some_mismatch_str}
")

```

This code is also available in github <https://github.com/amit7urmc/CombinationsAndPrimes>.