

The area between functions $x^{\frac{1}{n}}$ and x^n in the domain $0 \leq x \leq 1$ is equal to $\frac{n-1}{n+1}$

Amit Kumar

April 26, 2025

1 Abstract

When the graphs of functions $y = x^{\frac{1}{n}}$ and $y = x^n$ are plotted [in the domain $0 \leq x \leq 1$, range/co-domain $0 \leq y \leq 1$] then an area is carved between the two functions by the virtue of intersection of the two graphs (at 0 and 1). If we consider the area formed by the unit square [$0 \leq x \leq 1$ and $0 \leq y \leq 1$] to be 100%, then the above mentioned 'carved out area due to intersection' is a function of 'n' and grows as the value of 'n' grows, asymptotically reaching 100% for $n=\infty$. The relationship between the 'carved area' and 'n' is simply: $\Delta = (\frac{n-1}{n+1} \times 100)\%$ of the total area between the unit square formed.

2 Derivation

Let us see three plots to understand the behavior of the area carved out by the intersection of two curves. Below are the curves for $n=2, 3$ and 10 (Figure 1). It is easy to calculate the area formed by the intersection between




Figure 1: The area made by intersection is an increasing function of n

the two functions by going through these 3 steps.

1. Calculate the area under the function $x^{\frac{1}{n}}$.
2. Calculate the area under the function $y = x^n$.
3. Subtract the second step from the first step.

Thus we can calculate the area as below.

$$\begin{aligned}
\Delta &= \int_0^1 x^{\frac{1}{n}} dx - \int_0^1 x^n dx \\
&= \left. \frac{x^{\frac{1}{n}+1}}{\frac{1}{n}+1} \right|_0^1 - \left. \frac{x^{n+1}}{n+1} \right|_0^1 \\
&= \left[\frac{1^{\frac{1}{n}+1}}{\frac{1}{n}+1} - \frac{0^{\frac{1}{n}+1}}{\frac{1}{n}+1} \right] - \left[\frac{1^{n+1}}{n+1} - \frac{0^{n+1}}{n+1} \right] \\
&= \frac{1}{\frac{1}{n}+1} - \frac{1}{n+1} \\
&= \frac{n}{n+1} - \frac{1}{n+1} \\
&= \frac{n-1}{n+1}
\end{aligned}$$

For $n = 1$ there should be no area while for $n = \infty$ the area should cover the whole square intuitively. As a sanity check we can calculate the area for $n = 1$ and $n = \infty$ and we find the results to be 0 and 1 respectively, as expected.

3 Discussion

Below is a table for different values of n (from 1 to 100).

n	fraction
1	0.0000
2	0.3333
3	0.5000
4	0.6000
5	0.6667
6	0.7143
10	0.8182
25	0.9231
50	0.9608
100	0.9802

This observation can be turned into a plot as below (Figure 2).




Figure 2: The fraction asymptotically reaches 100% for large values of n

As we can see this equation $\left(y = \frac{n-1}{n+1}\right)$ is a special case of Möbius-Transformation. A Möbius-Transformation transformation is defined as $f(z) = \frac{az+b}{cz+d}$, where $ad - bc \neq 0$. In this context this function is special since z is replaced by x and $a = c = d = 1 \neq b = -1$. Please see <https://math.stackexchange.com/questions/5059632/does-the-function-y-fracx-1x1-have-a-name> for some very helpful comments on this.