


Kinetic Theory of Gases (K.T.)

Rule of thumb: Every material (solid, liquid, gas, plasma, intermediate phases) are made of atoms. They "may" attract or repel & form molecules of liquid or be restricted in definite shape of solid by huge cohesive force.

Experimental hints in favour of K.T.

1. Diffusion and Solution:  When slowly poured Alcohol over water, it spreads throughout.
2. Expansion of substance with heat: atoms tend to move away.
3. Phenomena of evaporation & vapour pressure.
4. Brownian motion. 1827 R. Brown + incessant motion of pollen on water.


Basic assumptions & postulates of K.T.

1. A gas consists of large number of identical atoms, which are rigid, elastic & equal mass objects.
2. Atoms are in chaos \rightarrow motion is fully irregular & spans in all three directions.
3. Inevitably the gas molecules collide with each other & surface of container (wall, sphere, cylinder). Total K.E. remains constant, but velocity of each atom continuously changes both in magnitude & direction. In evolving state (intermediate) density in a volume element will change but in steady state, collisions do not affect the density.
4. In between two successive collisions, molecules move in straight line following Newton's law.
5. Collisions are perfectly elastic i.e. no force of attraction/repulsion ($P.E. = 0$), energy is fully kinetic.

6. Atoms are "point" mass, meaning, their total volume $\ll \ll \ll$ volume of the container.

Concept of solid angle

Solid angle subtended by an area at a point is defined as the area intercepted by the cone on a unit sphere (radius = 1) with its centre at the apex of the cone.




If dS is an area that makes a solid angle $d\omega$ at origin O at a distance $OO' = r$, then from similar figures

$$\frac{d\omega}{1^2} = \frac{dS \cos \theta}{r^2} \quad \therefore d\omega = \frac{dS \cos \theta}{r^2}$$

unit of solid angle = steradian.

Ques 1. Calculate the solid angle (a) subtended by an infinite plain at a point in front of it, (b) hemisphere and (c) full sphere at its center.



Consider the annular ring,
 r distance apart from P & thick dr .

$$\begin{aligned} \text{Area of this ring} &= \pi(r+dr)^2 - \pi r^2 \\ &= 2\pi r dr \end{aligned}$$

(N.B. we throw $o(dr^2)$ term in limit $dr \rightarrow 0$)

So solid angle subtended by that circular annulus

$$d\omega = \frac{dS \cos \theta}{OG^2} = \frac{2\pi r dr \cos \theta}{r^2 + x^2}$$



Infinite plain meaning θ going from 0 to $\pi/2$.

2

$$\begin{aligned} \therefore w &= \int_{\theta=0}^{\pi/2} \frac{2\pi r dr \cos\theta}{r^2 + x^2} \\ &= 2\pi \int_{\theta=0}^{\pi/2} \frac{x \tan\theta \sec^2\theta d\theta \cos\theta}{x^2 \sec^2\theta} \\ &= 2\pi \int_0^{\pi/2} \sin\theta d\theta = \underline{2\pi}. \end{aligned}$$

$$\begin{aligned} [r &= x \tan\theta \\ dr &= x \sec^2\theta d\theta \\ \& \ x^2 + r^2 &= x^2 \sec^2\theta] \end{aligned}$$

Hemisphere



$$\begin{aligned} LP &= a \\ OP &= r \end{aligned}$$

$$\frac{a}{r} = \sin\theta$$

$$PQ = a d\phi$$

$$PS = r d\theta$$

$$\therefore dS = \text{area } PQRS = a d\phi \times r d\theta = r^2 \sin\theta d\theta d\phi$$

$$\text{So } dw \text{ at point } O = \frac{r^2 \sin\theta d\theta d\phi}{r^2} \times \cos\theta = \sin\theta d\theta d\phi$$

$$\text{So solid angle subtended} = \int dw = \int_{\theta=0}^{\pi/2} \int_{\phi=0}^{2\pi} \sin\theta d\theta d\phi = \underline{2\pi}.$$

Full sphere

$$\text{solid angle subtended} = \int_{\theta=0}^{\pi} \int_{\phi=0}^{2\pi} \sin\theta d\theta d\phi = \underline{4\pi}.$$

We will find out now pressure exerted by a perfect gas from K.T. (a) collisionless atoms in a box moving in 3 directions, (b) collisionless atoms coming from all directions. collision will be dealt in "mean free path".

Method 1

$$AB = AD = AE = l$$

The gas is confined within this cube of volume l^3 . P (say) is a gas atom

with velocity " c " whose components in 3-direction is (u, v, w) .

N = total no. of atoms or molecules.

So each of them have different velocity $c_1, c_2, c_3, c_4, \dots$ etc
so different components $(u_1, v_1, w_1), (u_2, v_2, w_2), (u_3, v_3, w_3), \dots$

$$\begin{aligned} \text{Mean square average } \overline{c^2} &= \frac{c_1^2 + c_2^2 + c_3^2 + \dots}{N} = \frac{u_1^2 + u_2^2 + u_3^2 + \dots}{N} \\ &+ \frac{v_1^2 + v_2^2 + v_3^2 + \dots}{N} + \frac{w_1^2 + w_2^2 + w_3^2 + \dots}{N} \\ &= \overline{u^2} + \overline{v^2} + \overline{w^2} \quad \text{--- (1)} \end{aligned}$$

mean square
velocity in
x-direction

y-direction

z-direction.

Consider particle P with mass m , velocity $\vec{c} = (u, v, w)$.

It travels from ABCD to EFGH, makes collision to exert pressure, rebounds elastically, momentum gets changed, comes back to ABCD to make another collision.

Total distance traveled with velocity u is $2l$.

\therefore Time between collision = $\frac{2l}{u}$, meaning

number of collision per second = $\frac{u}{2l}$.


Momentum imparted in $+x$ direction of on EFGH = mu .

Momentum obtained in $-x$ direction after collision = $-mu$.

\therefore change of momentum = $mu - (-mu) = 2mu$.

Rate of change of momentum for one atom in x direction

$$= 2mu \times \frac{u}{2l} = \frac{mu^2}{l}$$



Similarly in Y and Z direction, rate of change of momentum is $\frac{mv^2}{l}$ & $\frac{mw^2}{l}$ for one atom

∴ Total rate of change of momentum for all atoms per unit area along X direction is

$$P_x = \frac{m(u_1^2 + u_2^2 + u_3^2 + \dots)}{l} \times \frac{1}{l^2} = m \bar{u^2} \frac{N}{l^3} = mn \bar{u^2} \quad (\text{see eq. ①})$$

Similarly $P_y = mn \bar{v^2}$, $P_z = mn \bar{w^2}$.

In steady state, molecules move in all directions, so no preference, meaning $\bar{u^2} = \bar{v^2} = \bar{w^2}$ & $P_x = P_y = P_z$.


meaning $\bar{u^2} = \bar{v^2} = \bar{w^2} = \frac{1}{3} \bar{c^2}$ (see eq. ①)

collecting all pieces together,

$$P_x = P_y = P_z = \frac{1}{3} mn \bar{c^2} \quad \text{or}$$

$$P = \frac{1}{3} mn \bar{c^2}$$

Method 2



N no. of molecules moving in all directions with all possible velocity. How many collide with vessel & exert pressure?

number of vectors per unit area = $\frac{N}{4\pi r^2}$

∴ number of molecules at dA is $\frac{NdA}{4\pi r^2}$


We already learned that $dA = r^2 \sin\theta d\theta d\phi$

$$\frac{NdA}{4\pi r^2} = \frac{N}{4\pi} \sin\theta d\theta d\phi$$

∴ number of molecules per unit volume within velocity range c & $c+dc$ [dn_c], within direction θ & $\theta+d\theta$ & ϕ & $\phi+d\phi$

[$dw = \sin\theta d\theta d\phi$]

$$dn_{c,\theta,\phi} = \frac{dn_c}{4\pi} \sin\theta d\theta d\phi$$



Let's find now, how many of them strike dA of the wall of container. Geometrically, this is the number of molecules within the slanted prism of length cdt with edges in the direction θ & ϕ

$$= \frac{dn_c}{4\pi} \sin\theta d\theta d\phi \times c dA \cos\theta dt$$

∴ Total number of collisions at dA per unit time

$$\int_{c=0}^{\infty} \int_{\theta=0}^{\pi/2} \int_{\phi=0}^{2\pi} \frac{dn_c}{4\pi} \sin\theta d\theta d\phi \times c dA \cos\theta$$


$$= \frac{dA}{4\pi} \int_0^{\infty} c dn_c \underbrace{\int_0^{\pi/2} \sin\theta \cos\theta d\theta}_{1/2} \underbrace{\int_0^{2\pi} d\phi}_{2\pi} = \frac{dA}{4} \int_0^{\infty} c dn_c$$

If n_1 atoms per unit volume moves with velocity c_1 ,
 n_2 " " " " " " c_2 ,
 n_3 " " " " " " c_3, \dots

average velocity $\bar{c} = \frac{n_1 c_1 + n_2 c_2 + n_3 c_3 + \dots}{n_1 + n_2 + n_3 + \dots}$

$$= \frac{\sum n_i c_i}{\sum n_i} = \frac{1}{n} \int_0^{\infty} c \, dn_c$$

\therefore Number of molecules colliding at dA of the container per unit time = $\frac{dA}{4} n \bar{c}$



Now let's compute change in momentum by molecules striking area dA in unit time.

normal component of incident momentum
is $mc \cos \theta$ & reflected momentum $- mc \cos \theta$.