




## ENVC 24 : Energy and Environment

---

### Part-3 : Non-conventional Energy Resources



Kanyakumari Windmills,  
India




Pelamis Wave Energy  
Converter, Scotland



Krafla Geo-thermal  
Energy, Iceland


# Earth & Atmosphere

- World energy usage/year ➡ 500 ExaJoules.




# Earth & Atmosphere

- World energy usage/year ➡ 500 ExaJoules.
- Major constituents of dry air by volume %
  - ➡  $N_2=78.084$ ,  $O_2=20.946$ ,  $Ar=0.934$ ,  $CO_2=0.04$ ,
  - ➡  $Ne=0.001818$ ,  $He=0.000524$ ,  $CH_4=0.000179$ .



# Earth & Atmosphere

- World energy usage/year ➡ 500 ExaJoules.
- Major constituents of dry air by volume %
  - ➡  $N_2=78.084$ ,  $O_2=20.946$ ,  $Ar=0.934$ ,  $CO_2=0.04$ ,
  - ➡  $Ne=0.001818$ ,  $He=0.000524$ ,  $CH_4=0.000179$ .
- Water vapour is  $\sim 0.25\%$  by mass over full atmosphere and  $0.001\text{-}5\%$  by volume, which also abruptly varies locally according to weather/season.





# Earth & Atmosphere

- World energy usage/year → 500 ExaJoules.
- Major constituents of dry air by volume %
  - $N_2 = 78.084$ ,  $O_2 = 20.946$ ,  $Ar = 0.934$ ,  $CO_2 = 0.04$ ,
  - $Ne = 0.001818$ ,  $He = 0.000524$ ,  $CH_4 = 0.000179$ .
- Water vapour is  $\sim 0.25\%$  by mass over full atmosphere and  $0.001\text{--}5\%$  by volume, which also abruptly varies locally according to weather/season.

- Layers of the atmosphere →

**Troposphere** → 0 – 12 km


**Stratosphere\*** → 12 – 50 km

**Mesosphere** → 50 – 80 km

**Thermosphere** → 80 – 700 km

**Exosphere** → 700 –  $10^4$  km

\* → Ozone ( $O_3$ ) layer



# Earth & Atmosphere

- World energy usage/year → 500 ExaJoules.
- Major constituents of dry air by volume %
  - $N_2=78.084$ ,  $O_2=20.946$ ,  $Ar=0.934$ ,  $CO_2=0.04$ ,
  - $Ne=0.001818$ ,  $He=0.000524$ ,  $CH_4=0.000179$ .
- Water vapour is  $\sim 0.25\%$  by mass over full atmosphere and  $0.001\text{--}5\%$  by volume, which also abruptly varies locally according to weather/season.

- Layers of the atmosphere →

**Troposphere** → 0–12 km


**Stratosphere\*** → 12–50 km

**Mesosphere** → 50–80 km

**Thermosphere** → 80–700 km

**Exosphere** → 700– $10^4$  km

Approximately 80% mass of Earth's atmosphere is in the Troposphere.



\*→ Ozone ( $O_3$ ) layer

# Earth & Atmosphere

- World energy usage/year → 500 ExaJoules.
- Major constituents of dry air by volume %  
 →  $N_2 = 78.084$ ,  $O_2 = 20.946$ ,  $Ar = 0.934$ ,  $CO_2 = 0.04$ ,  
 →  $Ne = 0.001818$ ,  $He = 0.000524$ ,  $CH_4 = 0.000179$ .
- Water vapour is ~ 0.25% by mass over full atmosphere and 0.001-5% by volume, which also abruptly varies locally according to weather/season.



- Layers of the atmosphere →

**Troposphere** → 0 – 12 km

**Stratosphere\*** → 12 – 50 km

**Mesosphere** → 50 – 80 km

**Thermosphere** → 80 – 700 km

**Exosphere** → 700 –  $10^4$  km


Earth Radius(R) → 6371 Km

Earth+Troposphere Radius(R') → 6383 Km

Earth Volume →  $\frac{4}{3} \pi R^3 \sim 1.083 \times 10^{21} m^3$

Troposphere Volume →  $\frac{4}{3} \pi R'^3 - \frac{4}{3} \pi R^3$   
 $= 6.133 \times 10^{18} m^3$

Approximately 80% mass of Earth's atmosphere is in the Troposphere.



\* → Ozone ( $O_3$ ) layer

# Atmosphere

- As Volume of Troposphere is  $6.133 \times 10^{18} m^3$ , then 0.04% of  $CO_2$  accounts for  $2.453 \times 10^{15} m^3$ . To moderate on Greenhouse gas, estimate have to add on this number!!  
1 mole of  $CO_2$  corresponds to 22.4 litre or  $22.4 \times 10^{-3} m^3$  at S.T.P.( 1atm P, 0°C T).

$$\frac{0.04}{100} \times 6.133 \times 10^{18} = 2.453 \times 10^{15} m^3.$$





# Atmosphere

- As Volume of Troposphere is  $6.133 \times 10^{18} m^3$ , then 0.04% of  $CO_2$  accounts for  $2.453 \times 10^{15} m^3$ . To moderate on Greenhouse gas, estimate have to add on this number!!  
1 mole of  $CO_2$  corresponds to 22.4 litre or  $22.4 \times 10^{-3} m^3$  at S.T.P.( 1atm P, 0°C T).
- World energy usage/year ➡ 500 ExaJoules. So, heat of formation of  $CO_2$  is  $\Delta H = -394 kJ/mol$  and therefore  $CO_2$  emission amounts to an energy release/year ~  
$$\frac{5 \times 10^{20} \times 22.4 \times 10^{-3}}{3.94 \times 10^5} = 2.843 \times 10^{13} m^3 \text{ of } CO_2 / yr.$$



# Atmosphere

- As Volume of Troposphere is  $6.133 \times 10^{18} m^3$ , then 0.04% of  $CO_2$  accounts for  $2.453 \times 10^{15} m^3$ . To moderate on Greenhouse gas, estimate have to add on this number!!  
1 mole of  $CO_2$  corresponds to 22.4 litre or  $22.4 \times 10^{-3} m^3$  at S.T.P.( 1atm P, 0°C T).
- World energy usage/year  $\Rightarrow$  500 ExaJoules. So, heat of formation of  $CO_2$  is  $\Delta H = -394 kJ/mol$  and therefore  $CO_2$  emission amounts to an energy release/year  $\sim$   
$$\frac{5 \times 10^{20} \times 22.4 \times 10^{-3}}{3.94 \times 10^5} = 2.843 \times 10^{13} m^3 \text{ of } CO_2 / yr.$$
- Time required to double the amount of  $CO_2$  in the atmosphere at the present usage is  $\frac{2.453 \times 10^{15}}{2.843 \times 10^{13}} = 86 \text{ years} !!$  Big reasons to worry, as if we increase more, this time of doubling will be reduced and Greenhouse gas effects (global warming, snowstorms, ice-age?) can initiate towards a severe climate change. So energy usage by humans can significantly alter the composition of atmosphere within a very short period of time!

