Sem-I - Newtonian Mechanics

(Instructor: AKB, Department of Physics, Asutosh College)

Assignment II: Non-inertial Frame of Reference

Submission due date: 04/09/2019

Q.1) Show that for a single particle with constant mass, the equation of motion implies the following differential equation for the kinetic energy:

$$\frac{\mathrm{dE}}{\mathrm{dt}} = \mathbf{F} \cdot \mathbf{v},$$

while if the mass varies with time, the corresponding equation is


$$\frac{d(mE)}{dt} = \mathbf{F} \cdot \mathbf{p.}$$

[2]

Q.2) Calculate the rate of rotation of the plane of oscillation of a Foucault pendulum at latitude 30 degrees and hence determine the time it will take to turn through a complete right angle.

[2]

Q.3)



In figure beside, the polar coordinates (r, θ) are measured relative to an inertial frame \mathcal{F} , while \mathcal{F}' is a frame rotating about the z-axis passing through \mathcal{O} with angular velocity Ω . By using the velocity and acceleration transformation formula between inertial and rotating coordinate system, show that

$$\mathbf{v} = \dot{\mathbf{r}}\hat{\mathbf{r}} + \mathbf{r}\dot{\mathbf{\theta}}\hat{\mathbf{\theta}}, \quad \mathbf{a} = (\ddot{\mathbf{r}} - \mathbf{r}\dot{\mathbf{\theta}}^2)\hat{\mathbf{r}} + (\mathbf{r}\ddot{\mathbf{\theta}} + 2\dot{\mathbf{r}}\dot{\mathbf{\theta}})\hat{\mathbf{\theta}}.$$

[Note here as $\dot{\Omega} \neq 0$, therefore unlike our derivation in class, you also have to consider $\dot{\Omega} \times \mathbf{r}'$ term along with fictitious acceleration terms in the rotating coordinate system.]


[3]

Q.4) An ocean current circulating counter-clockwise when viewed from directly overhead was discovered in a well-isolated layer beneath the surface. The period of rotation was 14 hours ($\omega = \frac{2\pi}{14} \, \text{hr}^{-1}$). In Earth's surface fixed coordinate frame, the ocean current is x-axis pointing south, y-axis pointing east, z-axis pointing vertically upward and $\Omega = -\Omega \sin \lambda \hat{\mathbf{i}} + \Omega \cos \lambda \hat{\mathbf{k}}$ where λ is colatitude ($\Omega = \frac{2\pi}{24} \, \text{hr}^{-1}$). At what latitude and in which hemisphere was the current detected?

[4]

Q.5)

A turn-table marked with three orthogonal axes (x', y', z') is rotating on the Earth, assumed to be an inertial frame (x, y, z), with constant angular velocity $\boldsymbol{\omega}$ about z-axis. At t = 0, both frames coincide with each other. A ball of mass m is rolling outward without slipping along the x'-axis with a constant velocity $\boldsymbol{\nu}$. What is the total force exerted by the turn-table on



the ball? Find out the total force as measured in the Earth frame coordinates (x,y,z).

[4]