

On the Distribution of the Maximum Eigenvalue of Graphs

James B. Shearer

*Department of Mathematical Sciences
IBM Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598*

Dedicated to Alan J. Hoffman on the occasion of his 65th birthday.

Submitted by Richard A. Brualdi

ABSTRACT

Given a graph G , let $\lambda(G)$ denote the largest eigenvalue of the adjacency matrix of G . We prove that for any $\lambda \geq \sqrt{2+\sqrt{5}}$ (≈ 2.058) there exists a sequence of graphs G_1, G_2, \dots such that $\lim_{k \rightarrow \infty} \lambda(G_k) = \lambda$, thus answering a question posed by Hoffman.

In [2] Hoffman defined \mathbf{A} to be the set of all symmetric matrices (of all finite orders), every entry of which is a nonnegative integer. For $A \in \mathbf{A}$ let $\lambda(A)$ be the largest eigenvalue (spectral radius) of A . Let $\mathbf{R} = \{\lambda \mid \lambda = \lambda(A) \text{ for some } A \in \mathbf{A}\}$. Hoffman posed the problem of finding the limit points of \mathbf{R} , and in [2] found all limit points of $\mathbf{R} \leq \sqrt{2+\sqrt{5}}$. Here we complete the solution of the problem posed by Hoffman by showing that every point in $[\sqrt{2+\sqrt{5}}, \infty)$ is a limit point of \mathbf{R} . In fact we show this holds when \mathbf{A} is restricted to be the set of adjacency matrices of finite graphs.

Our result is the following theorem.

THEOREM. *For any $\lambda \geq \sqrt{2+\sqrt{5}}$ there exists a sequence of graphs G_1, G_2, \dots such that $\lim_{k \rightarrow \infty} \lambda(G_k) = \lambda$.*

Proof. Let

$$\phi_1 = \frac{\lambda - \sqrt{\lambda^2 - 4}}{2}, \quad \phi_2 = \frac{\lambda + \sqrt{\lambda^2 - 4}}{2}$$

be the roots of $x^2 - \lambda x + 1 = 0$. Define sequences $a_0, a_1, \dots; n_0, n_1, \dots$ recursively as follows: $a_0 = 1, a_1 = \lambda, n_0 = 0$,

$$n_k = \max \left\{ j \in \mathbf{Z} \left| \lambda - \frac{a_{k-1}}{a_k} - \frac{j}{\lambda} \geq \phi_1 \right. \right\}, \quad k \geq 1, \quad (1)$$

$$a_{k+1} = \left[\lambda - \left(\frac{n_k}{\lambda} \right) \right] a_k - a_{k-1}, \quad k \geq 1. \quad (2)$$

Let G_k be a tree consisting of a path $P_0 - P_1 - \dots - P_k$ with n_j leaves Q_{j1}, \dots, Q_{jn_j} connected to P_j , $j = 1, \dots, k$ (see Figure 1). We claim $\lim_{k \rightarrow \infty} \lambda(G_k) = \lambda$.

We must first show that the $n_k \geq 0$ so that the G_k are well defined. Define $r_k = a_k/a_{k-1}$, $k = 1, \dots$. Then

$$n_k = \max \left\{ j \in \mathbf{Z} \left| \lambda - \frac{1}{r_k} - \frac{j}{\lambda} \geq \phi_1 \right. \right\}, \quad k \geq 1, \quad (3)$$

$$r_{k+1} = \lambda - \frac{n_k}{\lambda} - \frac{1}{r_k}, \quad k \geq 1. \quad (4)$$

Note that $r_1 = \lambda > \phi_1$, and (3) and (4) imply $r_{k+1} \geq \phi_1$. Hence $r_k \geq \phi_1$ for any $k \geq 1$. Hence $\lambda - 1/r_k - \phi_1 \geq \lambda - 1/\phi_1 - \phi_1 = 0$. Therefore j may be assumed to be ≥ 0 so $n_k \geq 0$.

Next note that G_k is a subgraph of G_{k+1} , so the $\lambda(G_k)$ form an increasing sequence. We now assert that $\lambda(G_k) < \lambda$ for all k . To see this consider the sequences a_j and r_j as functions of λ (while holding the sequence n_j fixed). We claim that λ' is an eigenvalue of G_k iff $a_{k+1}(\lambda') = 0$. For let $b_j = a_j/\lambda'$, $j = 1, \dots, k$. Suppose v is an eigenvector of G_k . We may assume $v(P_0) = 1$. Then we may determine $v(P_j) = a_j$ and $v(Q_{ji}) = b_j$, $i = 1, \dots, n_j$, $j = 1, \dots, k$, recursively. For v to be an eigenvector we must have $\lambda'v(P_k) = v(P_{k-1}) + n_k b_k$, or $\lambda'a_k = a_{k-1} + n_k a_k/\lambda'$, or $(\lambda' -$




FIG. 1.

$n_k/\lambda')a_k - a_{k-1} = 0$, or $a_{k+1} = 0$ as claimed. Suppose $\lambda' > \lambda$. Then clearly $r_1(\lambda') > r_1(\lambda) \geq \phi_1 > 0$. Also $r_j(\lambda') > r_j(\lambda) \geq \phi_1 > 0$ together with (4) implies $r_{j+1}(\lambda') > r_{j+1}(\lambda)$, and we have already seen that $r_{j+1}(\lambda) \geq \phi_1 > 0$. Hence by induction we obtain $r_j(\lambda') > r_j(\lambda) \geq \phi_1 > 0$ for all $j = 1, \dots$. Hence for $\lambda' > \lambda$, $a_{k+1}(\lambda') = r_{k+1}(\lambda')r_k(\lambda') \cdots r_1(\lambda') > r_{k+1}(\lambda)r_k(\lambda) \cdots r_1(\lambda) \geq \phi_1^{k+1} > 0$. Hence the largest root of $a_{k+1}(\lambda')$ is $< \lambda$ which means $\lambda(G_k) < \lambda$, as asserted.

We have shown so far that $\lambda(G_1) < \lambda(G_2) < \cdots < \lambda(G_k) < \cdots < \lambda$. To complete the proof it will suffice to show that for any $\varepsilon > 0$ we can find a k so that $\lambda(G_k) > \lambda - \varepsilon$. We need the following theorem. Let M be a symmetric matrix, let $\lambda(M)$ be the largest eigenvalue of M , and let v be any nonzero vector. Then $\lambda(M) \geq v^T M v / v^T v$. For a proof see for example [1]. We apply this theorem as follows. Let M be the adjacency matrix of G_k . Let a vector v be defined on G_k as follows: $v(P_j) = a_j$, $j = 0, \dots, k$; $v(Q_{ji}) = b_j = a_j/\lambda$, $i = 1, \dots, n_j$, $j = 1, \dots, k$. Then we have $v^T M v = \lambda v^T v - a_k a_{k+1}$. Hence $\lambda(G_k) \geq \lambda - a_k a_{k+1} / v^T v$. Now $v^T v \geq a_1^2 + \cdots + a_k^2$. Suppose $a_j \leq B$ for all j . Then $\lim_{k \rightarrow \infty} a_k a_{k+1} / (a_1^2 + \cdots + a_k^2) = 0$ [for either $\lim_{k \rightarrow \infty} a_k = 0$ or $\lim_{k \rightarrow \infty} (a_1^2 + \cdots + a_k^2) = \infty$]. Hence it remains to show that the a_k are bounded. From (3) and (4) we know that r_{k+1} lies in the interval $[\phi_1, \phi_1 + 1/\lambda)$. Hence if $\phi_1 + 1/\lambda \leq 1$ (i.e. $\lambda \geq 2.325 -$), then $r_{k+1} < 1$ for $k > 1$, which means that $a_1 > a_2 > \cdots$ and the theorem follows at once. For $\lambda \in [2.058 + , 2.325 -]$ the argument is more complicated.

Suppose $\lambda \in (\theta_1, \theta_2)$, where $\theta_1 = \sqrt{2 + \sqrt{5}} = 2.058 +$ and θ_2 is the root of $\phi_1 + 1/\lambda = 1$ (or $\lambda^3 - 3\lambda^2 + 2\lambda - 1 = 0$; $\theta_2 = 2.325 -$). Then some of the r_k may be greater than 1, but any run of r_j 's greater than 1 is of bounded length and is followed immediately by a run of r_j 's less than 1 such that the product of the r_j 's in both runs is less than 1. Suppose $1 \leq r_k < \phi_1 + 1/\lambda$ and $n_k > 0$. Then we claim $r_{k+1}r_k < 1$. For

$$\begin{aligned} r_{k+1} &= \lambda - \frac{n_k}{\lambda} - \frac{1}{r_k} \\ \Rightarrow r_k r_{k+1} &\leq \left(\lambda - \frac{1}{\lambda} \right) r_k - 1 < \left(\lambda - \frac{1}{\lambda} \right) \left(\phi_1 + \frac{1}{\lambda} \right) - 1. \end{aligned}$$

Now a simple but tedious calculation, which we omit, shows $\lambda \geq \sqrt{2 + \sqrt{5}} \Rightarrow (\lambda - 1/\lambda)(\phi_1 + 1/\lambda) - 1 \leq 1$, so we have $r_k r_{k+1} < 1$, as claimed. Note since $r_k > 1$, this implies $r_{k+1} < 1$. Hence if $r_j, r_{j+1}, \dots, r_{j+i}, r_{j+i+1}$ is a run of r_j 's > 1 , we must have $n_j = n_{j+1} = \cdots = n_{j+i} = 0$. Now consider the map $f(r) = \lambda - 1/r$. This map has fixed points ϕ_1 and ϕ_2 and is increasing on (ϕ_1, ϕ_2) . Note $\lambda > \theta_1$ implies $\phi_1 + 1/\lambda < \phi_2$ (by another calculation which we

omit). Hence if $1 \leq r_k < \phi_1 + 1/\lambda$, some iterate of f on r_k will exceed $\phi_1 + 1/\lambda$. If $f^{j+1}(r_k)$ is the first iterate $\geq \phi_1 + 1/\lambda$, then it is clear that $r_{k+i} = f^i(r_k)$, $i = 0, \dots, j$; $n_{k+i} = 0$, $i = 0, \dots, j-1$; $n_{k+j} > 0$. Hence $r_{k+j+1} < 1$ by the preceding claim. Hence runs of r_j 's > 1 must be finite (and in fact can be bounded by the number of iterates of f on 1 which it takes to exceed $\phi_1 + 1/\lambda$; this number is clearly finite, as the iterates are increasing to ϕ_2 and we have $\phi_1 + 1/\lambda < \phi_2$). Suppose r_j, \dots, r_k is such a run (i.e. $r_{j-1} \leq 1 < r_j < r_{j+1} < \dots < r_k$; $n_j = \dots = n_{k-1} = 0$; $n_k > 0$, so $r_{k+1} < 1$). We have shown $r_k r_{k+1} < 1$. We claim $r_{k-1} r_{k+2} < 1$. For note that $r_k = \lambda - 1/r_{k-1}$, $r_{k+2} \leq \lambda - 1/r_{k+1}$. Therefore $r_{k-1} = 1/(\lambda - r_k)$, so

$$r_{k-1} r_{k+2} \leq \frac{1}{\lambda - r_k} \left(\lambda - \frac{1}{r_{k+1}} \right) = \frac{\lambda r_{k+1} - 1}{\lambda r_{k+1} - r_k r_{k+1}} < 1,$$

since $r_k r_{k+1} < 1$. Similarly, $r_{k-1} r_{k+2} < 1 \Rightarrow r_{k-2} r_{k+3} < 1$, etc. Now $1 < r_j < \dots < r_k$, so $r_{k-l} r_{k+1+l} < 1$ implies $r_{k+1+l} < 1$, $0 \leq l \leq k-j$. Also, $r_j r_{j+1} \dots r_k r_{k+1} \dots r_{k+1+k-j} < 1$, which establishes our claim that any run of r_j 's > 1 is balanced by a following run of r_j 's < 1 . Thus the a_j remain bounded (in fact, slightly more careful estimates would show $a_j < C(1-\varepsilon)^j$ for some $\varepsilon > 0$.) This completes the proof of the theorem for $\lambda > \sqrt{2+\sqrt{5}}$. When $\lambda = \sqrt{2+\sqrt{5}}$ this construction fails, but the theorem still holds, since for any $\varepsilon > 0$ we may find a graph G by the above construction such that $\lambda < \lambda(G) < \lambda + \varepsilon$. Letting $\varepsilon \rightarrow 0$, we obtain a sequence of such graphs, so that $\lim_{k \rightarrow \infty} \lambda(G_k) = \lambda$ as desired. ■

The author would like to thank Alan Hoffman for calling this problem to his attention and for encouraging him to write this note. The author would also like to acknowledge helpful conversations with Don Coppersmith and Bruce Kitchens concerning the construction given in this paper and the proof of its validity.

REFERENCES

- 1 D. K. Faddeer and R. N. Foddeera, *Computational Methods of Linear Algebra*, Freeman, San Francisco, 1963, pp. 85–86.
- 2 A. J. Hoffman, On limit points of spectral radii of non-negative symmetric integral matrices, in *Lecture Notes in Math.* 303, Springer, 1972, pp. 165–172.

Received 11 August 1987; final manuscript accepted 19 August 1987