

DISTRICT NUTRITION PROFILE

Led by IFPRI 🖔

BONGAIGAON | ASSAM

MARCH 2022

About District Nutrition Profiles:

District Nutrition Profiles (DNPs) are available for 707 districts in India. They present trends for key nutrition and health outcomes and their cross-sectoral determinants in a district. The DNPs are based on data from the National Family Health Survey (NFHS)-4 (2015-2016) and NFHS-5 (2019-2020). They are aimed primarily at district administrators, state functionaries, local leaders, and development actors working at



Figure 1: Map highlights district Bongaigaon in the state/UT of Assam

Optimum fetal and child nutrition and development

IMMEDIATE DETERMINANTS

Breastfeeding, nutrient-rich foods, caregiving practices, low burden of infectious diseases

UNDERLYING AND BASIC DETERMINANTS

Women's status, sanitation and hygiene, food security, socioeconomic conditions

Source: Adapted from Black et al. (2008)

NUTRITION-SPECIFIC INTERVENTIONS

Service delivery to mothers and infants along the continuum of care, access to health facilities

INTERVENTIONS THAT AFFECT UNDERLYING AND BASIC **DETERMINANTS**

Women's empowerment, sanitation, agriculture, and social safety net programs

What factors lead to child undernutrition?

Given the focus of India's national nutrition mission on child undernutrition, the DNPs focus on the determinants of child undernutrition (Figure on the left). Multiple determinants of suboptimal child nutrition and development contribute to the outcomes seen at the district-level. Different types of interventions can influence these determinants. Immediate determinants include inadequacies in food, health, and care for infants and young children, especially in the first two years of life. Nutrition-specific interventions such as health service delivery at the right time during pregnancy and early childhood can affect immediate determinants. Underlying and basic determinants include women's status, household food security, hygiene, and socio-economic conditions. Nutrition-sensitive interventions such as social safety nets, sanitation programs, women's empowerment, and agriculture programs can affect underlying and basic determinants.

District demographic profile, 2019-20

Bongaigaon



Sex ratio (females per 1,000 males) of the total population



227,195

Number of women in reproductive age (15-49 vrs)



Number of pregnant women



Number of live births



9,699

Total number of children under 5 vrs



13,595

Children under 5 yrs whose births were registered

- 1. IFPRI estimates The headcount was calculated as the product of the undernutrition prevalence and the total eligible projected population for each district in 2019. Projected population for 2019 was estimated using Census 2011
- 2. NFHS-4 (2015-16) & NFHS-5 district & state factsheets (2019-20).

Citation: Singh. N., P.H. Nguyen, M. Jangid, S.K. Singh, R. Sarwal, N. Bhatia, R. Johnston, W. Joe, and P. Menon. 2022. District Nutrition Profile: Bongaigaon, Assam. New Delhi,-India: International Food Policy Research Institute.

Acknowledgement: Financial support was provided by the Bill & Melinda Gates Foundation through POSHAN, led by the International Food Policy Research Institute. We thank Amit Jena (Independent Researcher) for design and programming support.














Assam





Burden on nutrition outcomes (2020)

Indicators	No. of children (<5 yrs)
Low-birth weight	NA
Stunted	36,821
Wasted	16,099
Severely wasted	5,914
Underweight	28,150
Overweight/obesity	7,021
Anemia	55,256
Total children	79,699


Note: NA refers to data are unavailable for a given round of NFHS data.

Points of discussion:

- · What are the trends in undernutrition among children under five years of age (stunting, wasting, underweight, and anemia)?
- What are the trends in overweight/obesity among children under five years of age in the district?

The state of nutrition outcomes among women (15-49 years)

Bongaigaon



Assam






Burden on nutrition outcomes (2020)

Indicators	No. of women (15-49 yrs)
Underweight	29,672
Overweight/obesity	38,464
Hypertension	30,899
Diabetes	28,672
Anemia (non-preg)	162,285
Anemia (preg)	7,866
Total women (preg)	15,378
Total women	227,195

Note: NA refers to data are unavailable for a given round of NFHS data.

Points of discussion:

- What are the trends in underweight and anemia among women (15-49 yrs) in the district?
- · What are the trends in overweight/obesity and other nutrition-related non-communicable diseases in the district?



Points of discussion:

• What are the trends in infant and young child feeding (timely initiation of breastfeeding, exclusive breastfeeding, timely initiation of complementary feeding, and adequate diet)? What can be done to improve infant and young child feeding?

40%

60%

80%

100%


Note: NA refers to data are unavailable for a given round of NFHS data.

• What are the trends in IFA consumption among pregnant women in the district? How can the consumption be improved?

20%

0%

· What additional data are needed to understand diets and/or other determinants?



Points of discussion:

- · How can the district increase women's literacy, and reduce early marriage, if needed?
- How does the district perform on providing drinking water and sanitation to its residents? Since sanitation and hygiene play an important role in improving nutrition outcomes, how can all aspects of sanitation be improved?
- How can programs that address underlying and basic determinants (education, poverty, gender) be strengthened?
- · What additional data are needed on food systems, poverty or other underlying determinants?

Demand for FP satisfied lodized salt Pregnancy registered (MPC card) ANC first trimester > 4 ANC visits Weighing Birth preparedness counselling Breastfeeding counselling Tetanus injection Received IFA tab/syrup Deworming Food supplementation Institutional birth

Financial assistance (JSY)

Skilled birth attendant

Postnatal care for mothers

Postnatal care for babies

Health & nutrition education

Full immunization

Pediatric IFA

Food supplementation (6-35 months)

Counselling on child growth


ORS during diarrhea

Zinc during diarrhea

Careseeking for ARI

Preschool at AWC







80%

Points of discussion:

· How does the district perform on health and nutrition interventions along the continuum of care? Does it adequately provide both prenatal and postnatal services to women of reproductive age, pregnant women, new mothers and newborns?

NA

0%

NA

40%

20%

· How has access to health and ICDS services changed over time (food supplementation, health and nutrition education and health checkups)?