

# **DISTRICT NUTRITION PROFILE**

Led by IFPRI 🖔

**BEGUSARAI | BIHAR** 

**MARCH 2022** 

# **About District Nutrition Profiles:**

District Nutrition Profiles (DNPs) are available for 707 districts in India. They present trends for key nutrition and health outcomes and their cross-sectoral determinants in a district. The DNPs are based on data from the National Family Health Survey (NFHS)-4 (2015-2016) and NFHS-5 (2019-2020). They are aimed primarily at district administrators, state functionaries, local leaders, and development actors working at



Figure 1: Map highlights district Begusarai in the state/UT of Bihar

What factors lead to child undernutrition?

Optimum fetal and child nutrition and development

#### **IMMEDIATE DETERMINANTS**

Breastfeeding, nutrient-rich foods, caregiving practices, low burden of infectious diseases

#### **UNDERLYING AND BASIC DETERMINANTS**

Women's status, sanitation and hygiene, food security, socioeconomic conditions

Source: Adapted from Black et al. (2008)

#### **NUTRITION-SPECIFIC INTERVENTIONS**

Service delivery to mothers and infants along the continuum of care, access to health facilities

#### INTERVENTIONS THAT AFFECT UNDERLYING AND BASIC **DETERMINANTS**

Women's empowerment, sanitation, agriculture, and social safety net programs

Given the focus of India's national nutrition mission on child undernutrition, the DNPs focus on the determinants of child undernutrition (Figure on the left). Multiple determinants of suboptimal child nutrition and development contribute to the outcomes seen at the district-level. Different types of interventions can influence these determinants. Immediate determinants include inadequacies in food, health, and care for infants and young children, especially in the first two years of life. Nutrition-specific interventions such as health service delivery at the right time during pregnancy and early childhood can affect immediate determinants. Underlying and basic determinants include women's status, household food security, hygiene, and socio-economic conditions. Nutrition-sensitive interventions such as social safety nets, sanitation programs, women's empowerment, and agriculture programs can affect underlying and basic determinants.

District demographic profile, 2019-20

Begusarai



Sex ratio (females per 1,000 males) of the total population



862,224

Number of women in reproductive age (15-49 vrs)



109,594

Number of pregnant women



82,127

Number of live births



Total number of children under 5 vrs



63,431

Children under 5 yrs whose births were registered

- 1. IFPRI estimates The headcount was calculated as the product of the undernutrition prevalence and the total eligible projected population for each district in 2019. Projected population for 2019 was estimated using Census 2011
- 2. NFHS-4 (2015-16) & NFHS-5 district & state factsheets (2019-20).

Citation: Singh. N., P.H. Nguyen, M. Jangid, S.K. Singh, R. Sarwal, N. Bhatia, R. Johnston, W. Joe, and P. Menon. 2022. District Nutrition Profile: Begusarai, Bihar. New Delhi, -India: International Food Policy Research Institute.

Acknowledgement: Financial support was provided by the Bill & Melinda Gates Foundation through POSHAN, led by the International Food Policy Research Institute. We thank Amit Jena (Independent Researcher) for design and programming support.














Bihar

# Burden on nutrition outcomes (2020)

Indicators	No. of children (<5 yrs)
Low-birth weight	NA
Stunted	159,915
Wasted	89,021
Severely wasted	37,353
Underweight	150,937
Overweight/obesity	17,067
Anemia	283,539
Total children	423,504


Note: NA refers to data are unavailable for a given round of NFHS data.

## Points of discussion:

- · What are the trends in undernutrition among children under five years of age (stunting, wasting, underweight, and anemia)?
- What are the trends in overweight/obesity among children under five years of age in the district?

# The state of nutrition outcomes among women (15-49 years)

Begusarai



Bihai


Burden on nutrition outcomes (2020)

Indicators	No. of women (15-49 yrs)
Underweight	224,092
Overweight/obesity	151,838
Hypertension	119,849
Diabetes	114,158
Anemia (non-preg)	544,839
Anemia (preg)	64,310
Total women (preg)	109,594
Total women	862,224

Note: NA refers to data are unavailable for a given round of NFHS data.


### Points of discussion:

- What are the trends in underweight and anemia among women (15-49 yrs) in the district?
- · What are the trends in overweight/obesity and other nutrition-related non-communicable diseases in the district?



### Points of discussion:

- What are the trends in infant and young child feeding (timely initiation of breastfeeding, exclusive breastfeeding, timely initiation of complementary feeding, and adequate diet)? What can be done to improve infant and young child feeding?
- What are the trends in IFA consumption among pregnant women in the district? How can the consumption be improved?
- · What additional data are needed to understand diets and/or other determinants?



## Points of discussion:

- · How can the district increase women's literacy, and reduce early marriage, if needed?
- How does the district perform on providing drinking water and sanitation to its residents? Since sanitation and hygiene play an important role in improving nutrition outcomes, how can all aspects of sanitation be improved?
- How can programs that address underlying and basic determinants (education, poverty, gender) be strengthened?
- · What additional data are needed on food systems, poverty or other underlying determinants?

Demand for FP satisfied lodized salt Pregnancy registered (MPC card) ANC first trimester > 4 ANC visits Weighing Birth preparedness counselling Breastfeeding counselling Tetanus injection Received IFA tab/syrup Deworming Food supplementation Institutional birth Financial assistance (JSY) Skilled birth attendant Postnatal care for mothers Postnatal care for babies Food supplementation Health & nutrition education Health checkup (ICDS) Full immunization Vitamin A Pediatric IFA Deworming Food supplementation (6-35 months) Weighing Counselling on child growth ORS during diarrhea Zinc during diarrhea







# Points of discussion:

· How does the district perform on health and nutrition interventions along the continuum of care? Does it adequately provide both prenatal and postnatal services to women of reproductive age, pregnant women, new mothers and newborns?

0%

20%

40%

80%

100% Note: NA refers to data are unavailable for a given round of NFHS data.

Careseeking for ARI

Preschool at AWC

Health checkup from AWC

· How has access to health and ICDS services changed over time (food supplementation, health and nutrition education and health checkups)?