



NORCET 9 MAINS (2025)



1. A 65-year-old patient with a history of diabetes mellitus and coronary artery disease presents with dizziness and near syncope. ECG shows the following, what is the most appropriate management. (AHN)



- A. Amiodarone
b. Transcutaneous pacemaker insertion
c. Anticoagulant therapy with continuous observation
d. Atropine
2. Identify the type of anesthesia shown in the image. (AHN)



- A. Subdural anesthesia b. Spinal anesthesia
c. Epidural anesthesia d. General anesthesia
3. A case of road traffic accident (RTA) develops neuropathic pain. Which of the following drugs is most appropriate for the management of neuropathic pain? (AHN)
- a. Acetaminophen b. Ibuprofen
c. Tramadol d. Gabapentin
4. A patient with rheumatoid arthritis is presented with a history of gastric ulcer. Which of the following drugs is most appropriate for this patient? (AHN)
- a. Celecoxib b. Naloxone
c. Acetaminophen d. Ibuprofen

5. Which of the following is not typically seen in a patient with heat stroke? (AHN)
- a. Excessive sweating b. Temperature 103°F
c. Rapid pulse d. Hot dry skin
6. A patient on ECMO continues to have difficulty breathing despite adequate settings. What is the most appropriate next step? (AHN)
- a. Increase tidal volume
b. Check for circuit obstruction
c. Increase ECMO flow rate
d. Increase Fio2 100%
7. In a patient with a shockable cardiac rhythm after how many shocks should adrenaline be administered? (AHN)
- a. 0.1 mL after the first shock
b. 0.1 mL after the second shock
c. 1 mg after the third shock
d. 1 mg after the second shock
8. Stenosis or obstruction of the Foramina of Luschka and Magendie is most commonly associated with which condition? (AHN)
- a. Dandy Walker malformation
b. Arnold Chiari malformation
c. Hydrocephalus due to aqueductal stenosis
d. Spina bifida
9. Which of the following anti-tuberculosis drugs is the most bactericidal? (AHN)
- a. Rifampicin b. Ethambutol
c. Isoniazid d. Pyrazinamide
10. A patient is prescribed an inhaler with 200 puffs the order is 2 puffs every 6 hours, for how many days will one inhaler last? (AHN)
- a. 20 days b. 25 days
c. 28 days d. 30 days
11. Calcium gluconate is not given by which method? (AHN)
- a. Fast IV b. Slow IV
c. IV infusion d. Oral route
12. What is the recommended breathing rate during CPR in a patient with an advanced airway? (AHN)
- a. 10 breaths per minute b. 12 breaths per minute
c. 20 breaths per minute d. 6 breaths per minute

Answer Key

1. b 2. c 3. d 4. a 5. a 6. b 7. c 8. a 9. a 10. b 11. a
12. a



13. A burn patient requires 4800 mL fluid in the first 24 hours as per the Parkland formula. How much fluid should be given in the first 8 hours? (AHN)
 a. 1200 mL b. 2400 mL
 c. 1800 mL d. 3000 mL
14. Elevated jugular venous pressure is most commonly caused by: (AHN)
 a. Hypervolemia b. Fluid overload
 c. Hypovolemia d. Dehydration
15. A patient receiving heparin has an aPTT of 85 seconds, what is the most appropriate next action? (AHN)
 a. Hold the next dose of heparin
 b. Increase the heparin dose
 c. Continue the same heparin dose
 d. Administer vitamin K
16. What is the primary purpose of a bougie during airway management? (AHN)
 a. To assist endotracheal tube placement in a difficult airway
 b. To measure airway resistance during mechanical ventilation
 c. To facilitate suctioning of tracheal secretions
 d. To maintain airway patency during mask ventilation
17. During central venous catheter (CVC) insertion, a patient develops sudden dyspnea and diminished breath sounds on one side, what is the most likely complication? (AHN)
 a. Pneumothorax b. Hemothorax
 c. Cardiac tamponade d. Air embolism
18. Which of the following is an early sign of increased intracranial pressure (ICP)? (AHN)
 A. Bradycardia b. Irregular respiration
 c. Headache and vomiting d. Hypertension
19. A COPD patient develops respiratory acidosis. Which intervention is most appropriate? (AHN)
 a. Administer oxygen via BiPAP
 b. Start mechanical ventilation
 c. Give sodium bicarbonate IV
 d. Restrict fluids
20. During dialysis a patient's potassium level rises to 6.8 mEq/L. What is the immediate life-saving intervention? (AHN)
 a. Administer calcium gluconate IV
 b. Start potassium-restricted diet
 c. Administer furosemide
 d. Restrict IV fluids
21. Atropine is given before abdominal surgery to: (AHN)
 a. Reduce gastric secretions b. Increase peristalsis
 c. Prevent infection d. Increase urine output
22. A patient is presented with pelvic trauma associated with testicular injury. What is the priority nursing action? (AHN)
 a. Insert Foley catheter immediately
 b. Avoid urethral catheterization
 c. Apply ice packs and reassurance only
 d. Start antibiotics and send urine culture
23. Which finding in a patient receiving opioid analgesics indicates the most serious concern and requires immediate intervention? (AHN)
 a. Constipation b. Nausea and vomiting
 c. Respiratory rate 10/min d. Urinary retention
24. A patient on mechanical ventilation shows sudden desaturation and increased peak airway pressure on the ventilator. What is the immediate nursing action? (AHN)
 a. Increase the oxygen flow
 b. Perform suctioning
 c. Administer bronchodilator
 d. Notify the physician
25. A patient with a history of peptic ulcer requires pain management, which analgesic approach is safest? (AHN)
 a. NSAIDS b. Avoid NSAIDS
 c. Aspirin d. NSAIDS with PPI
26. A patient develops headache and dizziness after a lumbar puncture, what is the most appropriate immediate nursing action? (AHN)
 a. Send sample to laboratory
 b. Position patient in prone
 c. Notify physician
 d. Encourage deep breathing
27. Which of the following is the most common side effect of nitroglycerin? (AHN)
 a. Bradycardia b. Headache
 c. Nausea d. Palpitations
28. A patient is suspected of having acute pancreatitis, which laboratory values are most commonly elevated? (AHN)
 a. Amylase and lipase b. Triglycerides
 c. ALT and AST d. Creatinine
29. A patient's arterial blood gas shows pH 7.50, PaCO₂ 28 mmHg, HCO₃⁻ 24 mEq/L. Which of the following is the most likely cause? (AHN)
 a. Chronic obstructive pulmonary disease
 b. Pulmonary embolism
 c. Air embolism
 d. Hypoxia
30. A patient is unconscious after trauma with suspected cervical spine injury. What is the priority nursing intervention? (AHN)
 a. Insert oropharyngeal airway
 b. Start IV fluids
 c. Perform neuro assessment
 d. Apply cervical immobilization
31. A patient with chronic kidney disease presents with hyponatremia what is the most appropriate initial nursing intervention? (AHN)
 a. Administer 0.9% NS
 b. Restrict fluids
 c. Administer 3% hypertonic saline
 d. Start diuretics

Answer Key

13. b 14. a 15. a 16. a 17. a 18. c 19. a 20. a 21. a 22. b 23. c
 24. b 25. b 26. b 27. b 28. a 29. b 30. d 31. a



32. A patient with central diabetes insipidus receives vasopressin. What is the expected effect on urine? (AHN)
- Urine volume remains high
 - Urine specific gravity rises
 - Urine volume does not change
 - Urine specific gravity lowers
33. Patients on monoamine oxidase inhibitors should avoid which of the following? (AHN)
- Tyramine
 - Glucose
 - Fructose
 - Lactose
34. A patient is presents with dilated pupils, elevated blood pressure, and tachycardia. What type of pain is this? (FON)
- Acute pain
 - Chronic pain
 - Neuropathic pain
 - Visceral pain
35. The passage of bright red blood per rectum is referred to as: (FON)
- Hematochezia
 - Melena
 - Hematemesis
 - Occult blood
36. A patient's chest tube becomes accidentally dislodged. What is the most appropriate immediate nursing action? (FON)
- Immerse the tube end in sterile saline
 - Apply an airtight dressing
 - Clamp the chest tube
 - Encourage deep breathing
37. During suctioning a patient's oxygen saturation drops from 94% to 88%. What should be the immediate nursing action? (FON)
- Continue suctioning until airway clears
 - Stop suction and administer oxygen
 - Increase suction pressure and continue
 - Call the physician immediately
38. The patients urine output is 100 mL in 24 hours, how is this classified? (FON)
- Oliguria
 - Anuria
 - Polyuria
 - Normal urine output
39. A nurse is instructed to infuse 1000 mL of IV fluid over 8 hours using an IV set with a drop factor of 15 gtt/ML. What should be the correct infusion rate in drops per minute? (FON)
- 42 drops/min
 - 31 drops/min
 - 25 drops/min
 - 18 drops/min
40. A woman undergoing evaluation for infertility is advised a hysterosalpingogram. On which day of the menstrual cycle should the nurse schedule this procedure? (OBGY)
- Day 21
 - Day 14
 - Day 3
 - Day 8

41. Identify the given instrument: (OBGY)



- Green Armytage Forceps
- Babcock Forceps
- Allis Tissue Forceps
- Kocher's Forceps

42. Identify the given instrument: (OBGY)



- Doyen's Retractor
- Deaver Retractor
- Langenbeck Retractor
- Morris Retractor

43. Identify the item shown in the image given below: (OBGY)




- Obstetric V-Drape
- Episiotomy Pack
- Perineal Sheet
- Cesarean Section Drape

Answer Key

32. b 33. a 34. a 35. a 36. b 37. b 38. b 39. b 40. d 41. a 42. b 43. a



44. A cardiotocography tracing shows cord compression. Which fetal heart rate pattern is most expected? (OBGY)
 a. Variable decelerations b. Late decelerations
 c. Early decelerations d. Hypoxia
45. A patient develops uterine hyperstimulation after administration of prostaglandins for induction of labour, what is the most appropriate management? (OBGY)
 a. Calcium gluconate
 b. Magnesium sulphate
 c. Hydralazine
 d. Terbutaline
46. A 28-year-old woman presents with a history of 3 consecutive first-trimester abortions. She has no structural uterine abnormality and her blood sugar levels are normal on investigation she is found to be positive for antiphospholipid antibody. What is the most likely diagnosis? (OBGY)
 a. Antiphospholipid syndrome
 b. Recurrent pregnancy loss due to uterine anomaly
 c. Gestational diabetes mellitus
 d. Pre-eclampsia
47. Visual Inspection with Acetic Acid (VIA) is primarily used for the screening of which condition? (OBGY)
 a. Cervical cancer b. Endometrial cancer
 c. Ovarian cancer d. Vaginal prolapse
48. A patient develops postpartum hemorrhage at a Primary Health Centre. What is the first-line management? (OBGY)
 a. Start IV tranexamic acid immediately
 b. Perform immediate hysterectomy
 c. Administer uterotonic (Oxytocin)
 d. Begin blood transfusion
49. Which of the following is true about the biophysical profile in obstetrical assessment? (OBGY)
 a. It assesses fetal well-being using fetal heart rate patterns only.
 b. It includes fetal breathing movements, gross body movements, fetal tone, amniotic fluid volume, and non-stress test results.
 c. It is solely based on ultrasound assessment without fetal monitoring.
 d. It is performed only during the first trimester of pregnancy.
50. A patient is in labour, when does the second stage of labour starts? (OBGY)
 a. At full cervical dilation (10 cm)
 b. At 8 cm cervical dilation with strong expulsive contractions
 c. At the onset of regular uterine contractions
 d. At rupture of membranes
51. Which of the following maneuvers is specifically used for head delivery during labour? (OBGY)
 a. Ritgen maneuver
 b. McRoberts maneuver
 c. Pinard's maneuver
 d. Woods screw maneuver
52. In cervicography crossing, which line indicates abnormal labour? (OBGY)
- 
- a. Alert line b. Action line
 c. Slow progress d. No descent
53. A patient in labour has a cervical dilatation of 6 cm with contractions every 5 minutes. What is the appropriate nursing action? (OBGY)
 a. Immediate augmentation
 b. Monitor contractions and FHR
 c. Prepare for delivery
 d. Administer oxytocin
54. Identify the pattern shown in the CTG strip. (OBGY)
 a. Early deceleration
 b. Fetal tachycardia
 c. Fetal bradycardia
 d. Late deceleration
55. Among the following, which food provides the highest amount of protein per 100 grams? (B&N)
 a. Egg b. Soybean
 c. Mutton d. Peanuts
56. In which year was the first National Family Planning Programme launched in India? (CHN)
 a. 1947 b. 1952
 c. 1961 d. 1971
57. Which was the first National Family Planning Programme launched in India? (CHN)
 a. National Population Policy
 b. Family Welfare Programme
 c. Reproductive and Child Health Programme
 d. National Rural Health Mission
58. A 45-year-old patient on isoniazid therapy for tuberculosis presents with complaints of numbness and tingling in the extremities, suggestive of peripheral neuropathy. The nurse suspects; this is related to a deficiency caused by the medication. Which intervention should the nurse prioritize? (CHN)
 a. Administer vitamin B12 supplements
 b. Increase the dose of isoniazid
 c. Administer vitamin B6 (pyridoxine) supplements
 d. Discontinue isoniazid immediately

Answer Key

44. a 45. d 46. a 47. a 48. c 49. b 50. a 51. a 52. a 53. b 54. a
 55. b 56. b 57. b 58. c



59. A 30-year-old patient presents with fever, weight loss, and skin lesions after traveling to a rural area. The nurse suspects leishmaniasis based on the patient's history and symptoms. Which vector is most likely responsible for transmitting this disease? (CHN)
- Mosquito
 - Tsetse fly
 - Sandfly
 - Tick
60. A community health nurse is educating parents about preventive measures for gastroenteritis in their 6-week-old infant. The nurse discusses the administration of a vaccine to protect against rotavirus. Which of the following is the correct route of administration for the rotavirus vaccine? (CHN)
- Intramuscular
 - Subcutaneous
 - Oral
 - Intravenous
61. A community health nurse is conducting an awareness session on vaccine-preventable diseases. A participant asks whether herd immunity can provide protection against tetanus in unvaccinated individuals. What is the most accurate response the nurse should provide? (CHN)
- Herd immunity fully protects against tetanus in unvaccinated individuals
 - Herd immunity partially protects against tetanus due to community vaccination
 - Herd immunity does not protect against tetanus as it is not a contagious disease
 - Herd immunity protects only vaccinated individuals against tetanus
62. A community health nurse is caring for a 4-year-old child presenting with prolonged fever, rash, and conjunctivitis. The physician suspects Kawasaki disease. Which type of blood vessels is primarily affected in this condition? (CHN)
- Large arteries
 - Medium-sized arteries
 - Small capillaries
 - Veins
63. A public health nurse is reviewing data on leading causes of mortality in India during a community health assessment. Among the infectious diseases, which condition is a significant contributor to mortality rates in the country? (CHN)
- Malaria
 - Tuberculosis
 - Dengue fever
 - Cholera
64. A 60-year-old female beneficiary under the Pradhan Mantri Jan Arogya Yojana (PM-JAY) presents to the emergency department with severe abdominal pain requiring hospitalization for surgery. The nurse verifies the patient's eligibility for cashless treatment. What is the annual coverage limit provided by PM-JAY for this beneficiary's family? (CHN)
- Rs. 1 lakh per family per year
 - Rs. 3 lakhs per family per year
 - Rs. 5 lakhs per family per year
 - Rs. 10 lakhs per family per year
65. A community health nurse is organizing a nutrition awareness camp under the Poshan Abhiyaan to address malnutrition in children and women. Which ministry is primarily responsible for implementing this national program? (CHN)
- Ministry of Health and Family Welfare
 - Ministry of Women and Child Development
 - Ministry of Education
 - Ministry of Rural Development
66. A community health nurse is counseling a pregnant woman in a rural area about the benefits of institutional delivery under the Janani Shishu Suraksha Yojana (JSSY). What is the primary objective of this program? (CHN)
- Provide free postnatal care only
 - Promote institutional delivery and ensure free maternal and newborn care
 - Offer financial assistance for home deliveries
 - Provide nutritional supplements to pregnant women
67. A community health nurse is educating parents about the BCG vaccine administered to newborn to prevent tuberculosis. The nurse explains when the immune response to the vaccine is typically assessed. What is the recommended time frame to check for a response to the BCG vaccine? (CHN)
- 2-4 weeks
 - 4-6 weeks
 - 8-10 weeks
 - 12-14 weeks
68. A community health nurse is conducting a survey to assess the burden of illness in a rural community for planning health interventions. Which epidemiological measure is most appropriate to determine the proportion of the population affected by a specific disease at a given time? (CHN)
- Incidence rate
 - Prevalence
 - Mortality rate
 - Attack rate
69. A public health nurse is educating a group of nursing students about the characteristics of killed vaccines during a vaccination drive. One student asks whether killed vaccines require booster doses. Which statement is true regarding killed vaccines? (CHN)
- Killed vaccines never require booster doses
 - Killed vaccines typically require booster doses to maintain immunity
 - Killed vaccines provide lifelong immunity without boosters
 - Killed vaccines are only effective with a single dose
70. A community health nurse is analyzing the demographic profile of a rural population during a health planning session. The age-sex pyramid shows a narrow top and a wide base. What does this demographic pattern indicate about the population? (CHN)
- Low birth rate and low death rate
 - High birth rate and high death rate
 - Low birth rate and high death rate
 - High birth rate and low death rate

Answer Key

59. c 60. c 61. c 62. b 63. b 64. c 65. b 66. b 67. c 68. b 69. b 70. b



71. A community health nurse is counseling a group of women on incorporating high-protein foods into their diet to address malnutrition. Which of the following foods contains the highest amount of protein per serving? (CHN)
- Lentils
 - Soybean
 - Eggs
 - Rice
72. A public health nurse is explaining the features of the Ayushman Bharat Digital Mission to a group of healthcare workers during a training session. What is the primary purpose of the unique health ID provided under this mission? (CHN)
- To provide financial assistance for hospital admissions
 - To create a universal health identity for seamless healthcare access
 - To track vaccination records only
 - To monitor disease outbreaks in real-time
73. A community health nurse is developing a plan to improve maternal and child health outcomes in a rural area under the Reproductive and Child Health (RCH) program. Which framework integrates adolescent health with reproductive and child health services? (CHN)
- RMNCH
 - RMNCH+A
 - NRHM
 - ASHA
74. A pediatric nurse is educating parents about the inactivated polio vaccine (IPV) during a routine immunization visit for their 2-month-old child. Which of the following statements about IPV is true? (CHN)
- IPV is administered orally to prevent polio
 - IPV is a live attenuated vaccine
 - IPV is given as an injectable vaccine to provide immunity against polio
 - IPV requires only a single dose for lifelong immunity
75. A public health nurse is conducting a training session for nursing students on the components of vaccines. One student asks about the role of adjuvants in vaccines. Which of the following statements is true about adjuvants? (CHN)
- Adjuvants weaken the immune response to vaccines
 - Adjuvants are used to enhance the immune response to vaccines
 - Adjuvants are live pathogens included in vaccines
 - Adjuvants prevent the vaccine from being effective
76. A community health nurse is preparing for the annual nutrition awareness activities in September, focusing on malnutrition prevention among children and women. She plans to organize events. Under which national program does it primarily fall? (CHN)
- Ayushman Bharat
 - Swachh Bharat Abhiyan
 - Poshan Abhiyaan
 - National Health Mission
77. A nurse is reviewing the treatment plan for a 35-year-old patient diagnosed with pulmonary tuberculosis. The physician emphasizes the use of bactericidal drugs to effectively kill Mycobacterium tuberculosis. Which of the following anti-tubercular drugs is primarily bactericidal? (CHN)
- Ethambutol
 - Isoniazid
 - Pyrazinamide
 - Both Isoniazid and Pyrazinamide
78. A community health nurse is educating a group of pregnant women in a rural area about a national program aimed at reducing maternal and newborn mortality rates through safe delivery and comprehensive care. Under which program are these efforts primarily implemented? (CHN)
- Poshan Abhiyaan
 - Janani Shishu Suraksha Yojana (JSSY)
 - Ayushman Bharat
 - National Rural Health Mission (NRHM)
79. A community health nurse is managing a 28-year-old woman who develops postpartum hemorrhage (PPH) after delivery and is referred to the Primary Health Center (PHC). Which medication should the nurse prioritize to manage the PPH effectively? (CHN)
- Misoprostol
 - Oxytocin
 - Tranexamic acid
 - Ergometrine
80. A nurse is educating a community group about initiatives aimed at reducing newborn and maternal mortality rates. Which national programme in India is specifically designed to address these issues through comprehensive maternal and child health services? (CHN)
- National Rural Health Mission (NRHM)
 - Janani Shishu Suraksha Karyakram (JSSK)
 - Integrated Child Development Services (ICDS)
 - National Immunization Programme
81. A psychiatric nurse is assessing a 25-year-old male patient who expresses a persistent urge to wear female clothing and identifies with a gender different from his assigned sex at birth. The nurse suspects a gender-related condition. What is the most appropriate term for this presentation? (MHN)
- Gender dysphoria
 - Transvestism
 - Homosexuality
 - Gender conformity
82. A psychiatric nurse is assessing a 30-year-old patient admitted with suspected schizophrenia. The patient exhibits disorganized thinking and reports hearing voices commenting on his actions. Which of the following is considered a first-rank symptom of schizophrenia? (MHN)
- Social withdrawal
 - Thought disorder (e.g., thought broadcasting)
 - Flat affect
 - Visual hallucinations

Answer Key

71. b 72. b 73. b 74. c 75. b 76. c 77. d 78. b 79. b 80. b 81. b 82. b



83. A psychiatric nurse is monitoring a 35-year-old patient on antipsychotic medication who reports a constant urge to move and an inability to stay seated during ward activities. The nurse suspects a medication side effect. Which condition is most likely causing these symptoms? (MHN)
a. Tardive dyskinesia b. Akathisia
c. Dystonia d. Parkinsonism
84. A nurse is caring for a 50-year-old patient diagnosed with neuropathic pain following a nerve injury. The physician prescribes a medication to manage the pain. Which of the following medications is most commonly used for neuropathic pain in this scenario? (MHN)
a. Ibuprofen b. Gabapentin
c. Paracetamol d. Morphine
85. A psychiatric nurse is counseling a 40-year-old patient undergoing disulfiram therapy for alcohol dependence. What is the most critical advice the nurse should provide to ensure the patient's safety during treatment? (MHN)
a. Avoid caffeine-containing beverages
b. Avoid alcohol and alcohol-containing products
c. Increase fluid intake to prevent dehydration
d. Take the medication with meals to reduce side effects
86. A psychiatric nurse is developing a care plan for a 28-year-old patient diagnosed with borderline personality disorder who exhibits emotional instability and impulsive behavior. Which therapy is most commonly recommended as the first-line treatment for this condition? (MHN)
a. Cognitive Behavioral Therapy (CBT)
b. Dialectical Behavior Therapy (DBT)
c. Psychoanalytic Therapy
d. Family Therapy
87. A psychiatric nurse is assessing a patient who reports a low mood for three months followed by a recent period of excessive energy, talkativeness, and reckless behavior. The physician suspects Bipolar I disorder. What is the most likely current diagnosis for this patient? (MHN)
a. Major depressive disorder
b. Bipolar I disorder with current manic episode
c. Generalized anxiety disorder
d. Bipolar II disorder with hypomania
88. A mental health nurse is evaluating a patient who reports persistent low mood, low energy, and feelings of hopelessness for the past few years, without episodes of major depression. The nurse suspects a chronic mood disorder. What is the most likely diagnosis for this patient? (MHN)
a. Major depressive disorder
b. Dysthymia
c. Bipolar II disorder
d. Cyclothymic disorder
89. A nurse is monitoring a 45-year-old patient receiving opioid therapy for post-surgical pain. The nurse observes that the patient's respiratory rate has dropped to 10 breaths per minute. What is the primary concern associated with this finding? (MHN)
a. Hypertension
b. Respiratory depression
c. Tachycardia
d. Hyperglycemia
90. A psychiatric nurse is assessing a patient with suspected schizophrenia who speaks in a way that frequently drifts off-topic, making it difficult to follow the conversation. The nurse identifies this as a disorder of thought form. Which term best describes this presentation? (MHN)
a. Flight of ideas b. Tangentiality
c. Thought blocking d. Loose associations
91. A mental health nurse is caring for a 12-year-old patient diagnosed with conduct disorder, exhibiting aggressive behavior and disregard for rules. What is a key nursing role in managing this patient's condition in a structured environment? (MHN)
a. Encourage free expression without boundaries
b. Implement strict rules and regulations with consistent enforcement
c. Avoid setting limits to reduce confrontations
d. Focus solely on medication administration
92. A nurse is assessing a patient newly diagnosed with schizophrenia. Which of the following is not considered a negative symptom of schizophrenia? (MHN)
a. Apathy b. Alogia
c. Ambivalence d. Anhedonia
93. The Anemia Mukht Bharat (AMB) program uses specific hemoglobin (Hb) cut-offs to define anemia for various population groups. For a pregnant woman in her second trimester, which of the following Hb levels represents the cut-off below which she is classified as anemic? (MHN)
a. Hb<11.0 g/dL b. Hb<12.0 g/dL
c. Hb<10.0 g/dL d. Hb<13.0 g/dL
94. A 45-year-old client with a history of schizophrenia, whose symptoms have been well-controlled for 5 years on a stable dose of a typical antipsychotic, suddenly begins to exhibit involuntary, repetitive movements, including lip smacking and tongue protrusion. The nurse correctly identifies this as Tardive Dyskinesia (TD). What is the priority management strategy? (MHN)
a. Initiate treatment with Propranolol to control the motor movements.
b. Immediately increase the dose of the current antipsychotic medication.
c. Substitute the current antipsychotic with Trihexyphenidyl for symptom control.
d. Discuss a plan to decrease the dose of the current antipsychotic or switch to an atypical agent.

Answer Key

83. b 84. b 85. b 86. b 87. b 88. b 89. b 90. b 91. b 92. c 93. a
94. d



95. A psychiatric nurse is assessing a 38-year-old patient who believes that his neighbors are plotting to harm him and are constantly monitoring his activities. The nurse identifies this as a specific type of delusion. Which term best describes this patient's belief? (MHN)
- Delusion of grandeur
 - Delusion of persecution
 - Delusion of reference
 - Delusion of control
96. Identify the below given image: (Pediatrics)
- 
- Haberman feeder
 - Pigeon bottle
 - Cleft lip/palate nurser
 - Both a and b
97. What is the full form of INAP? (Pediatrics)
- India Newborn Action Plan
 - India Neonatal Action Plan
 - India Newborn Active Plan
 - India Neonatal Active Plan
98. Boot shaped heart is seen in: (Pediatrics)
- Tricuspid atresia
 - Tetralogy of Fallot
 - Truncus arteriosus
 - Total anomalous pulmonary venous return
99. A baby with eczema is prescribed a topical corticosteroid by the physician. What advice should be given to the parent? (Pediatrics)
- Apply at the night time to reduce the systemic absorption
 - Gastric irritation occurs when the hands are put in the mouth
 - It causes the hair growth at the site of application
 - Apply a thin layer only on the affected area and avoid prolonged use
100. In NICU which respiratory distress assessment score is used? (Pediatrics)
- Silverman Anderson score
 - Downs score
 - Both a and b
 - Only a
101. A nurse is assessing a child suspected of having increased intracranial pressure (ICP). Which of the following is an early sign of increased ICP in children? (Pediatrics)
- Altered level of consciousness
 - Headache
 - Nausea and vomiting
 - Seizures
102. A nurse is caring for a pediatric client receiving respiratory support via a High-Flow Nasal Cannula (HFNC). During assessment, the nurse notes the client's oxygen saturation (SpO₂) is 50%. What is the most appropriate and immediate action for the nurse to take? (Pediatrics)
- Increase the flow rate of the High-Flow Nasal Cannula (HFNC) by 5 L/min.
 - Call the Respiratory Therapist to adjust the FiO₂.
 - Immediately discontinue the HFNC and switch the child to bag-valve-mask (BVM) ventilation with 100% oxygen.
 - Reposition the child to prone and reassess the SpO₂ after 15 minutes.
103. A nurse is conducting a health assessment under the ICDS program. A mother reports that her 4-year-old child is not receiving supplementary nutrition. What should the nurse do first? (Pediatrics)
- Refer the child to a pediatrician for evaluation.
 - Verify the child's age and enrollment status in the ICDS program.
 - Educate the mother about the importance of nutrition.
 - Provide immediate supplementary food to the child.
104. A 2-month-old infant is brought to the pediatric OPD by the parents with complaints of increasing head circumference since birth, vomiting, and downward deviation of the eyes. On examination, the nurse notes a tense anterior fontanelle and sunset sign. MRI reveals cystic dilatation of the fourth ventricle with hypoplasia of the cerebellar vermis. The nurse suspects Dandy-Walker syndrome. What is the underlying pathophysiological mechanism? (Pediatrics)
- Aqueductal stenosis leading to supratentorial hydrocephalus.
 - Atresia of the foramina of Luschka and Magendie causing non-communicating hydrocephalus.
 - Overproduction of CSF by the choroid plexus in the lateral ventricles.
 - Malabsorption of CSF due to defective arachnoid granulations.
105. A nurse is managing a 5-year-old child with acute diarrhea. As part of the treatment plan under the ICDS program, the nurse is required to provide zinc supplementation. What is the appropriate dose and duration of zinc supplementation for this child? (Pediatrics)
- 10 mg daily for 7 days
 - 20 mg daily for 14 days
 - 15 mg daily for 10 days
 - 30 mg daily for 7 days
106. A nurse is assessing a newborn in the delivery room 1 minute after birth. The newborn's heart rate is 120 beats per minute, with irregular respirations, some flexion of extremities, and a weak cry. The pulse oximetry reading shows an SpO₂ of 62%. What should the nurse do next? (Pediatrics)
- Initiate positive pressure ventilation immediately.
 - Administer supplemental oxygen and monitor SpO₂.
 - Perform chest compressions and intubate the newborn.
 - Dry the newborn, provide warmth, and reassess after 30 seconds.

Answer Key

95. b 96. a 97. a 98. b 99. d 100. c 101. a 102. c 103. b 104. b 105. b 106. b



107. During a neonatal workshop while discussing the role of the ductus venosus, a student asks about its anatomical function. What is the correct description of the ductus venosus in fetal circulation? (Pediatrics)
- Connects the pulmonary artery to the aorta to bypass the lungs.
 - Shunts oxygenated blood from the umbilical vein to the inferior vena cava.
 - Links the umbilical artery to the portal vein for nutrient delivery.
 - Allows blood flow from the right atrium to the left atrium through the foramen ovale.
108. A 6-week-old infant presents with projectile vomiting after feeds, weight loss, and a palpable olive-shaped mass in the right upper quadrant of the abdomen. On examination, visible peristaltic waves are noted. What is the most likely diagnosis? (Pediatrics)
- Gastroesophageal reflux disease (GERD)
 - Intussusception
 - Pyloric stenosis
 - Duodenal atresia
109. A newborn is diagnosed with clubfoot, presenting with a foot that is turned inward and downward. The parents are concerned about the best initial management for this condition. What is the first-line treatment for this infant's clubfoot? (Pediatrics)
- Immediate surgical correction
 - Observation and follow-up
 - Serial plaster casting (Ponseti method)
 - Physical therapy alone
110. A 3-year-old child presents with a history of fever for 3 days, chills, and no other specific symptoms. According to the Integrated Management of Neonatal and Childhood Illness (IMNCI) guidelines, what is the most appropriate initial management for suspected malaria in this child? (Pediatrics)
- Administer oral antibiotics immediately
 - Refer to a higher center without treatment
 - Perform a rapid diagnostic test for malaria and treat if positive
 - Start antipyretics and observe for 48 hours
111. A newborn is diagnosed with neonatal diabetes mellitus type I, presenting with hyperglycemia and weight loss. The pediatrician discusses the management plan with the parents. According to standard protocols, what is the most appropriate initial treatment for this condition? (Pediatrics)
- Oral hypoglycemic agents
 - Insulin therapy
 - Dietary management alone
 - Observation without medication
112. A preterm newborn in the NICU is being evaluated for respiratory distress syndrome (RDS). The nurse uses a standardized scale to assess the severity of the condition, noting signs such as grunting, nasal flaring, and retractions. Which scale is most commonly used to assess the severity of RDS in this scenario? (Pediatrics)
- Apgar score
 - Silverman-Anderson score
 - Ballard score
 - Dubowitz score
113. The nurse observes that a 3-year-old child is playing with clay and notes hold and manipulate the clay using fingers. According to developmental norms, which fine motor skill is this child demonstrating? (Pediatrics)
- Scribbling with a crayon
 - Building a tower of 6–8 blocks
 - Using fingers to mold clay
 - Copying a circle
114. A newborn is diagnosed with a meningocele, and the pediatric nurse is preparing to assist with cerebrospinal fluid (CSF) analysis to evaluate for associated complications. During the assessment, the nurse observes a sac-like protrusion on the infant's lower back. What is the most likely finding in the CSF analysis if the meningocele is not associated with infection or hemorrhage? (Pediatrics)
- Elevated white blood cell count
 - Normal CSF composition
 - Increased red blood cell count
 - Decreased glucose levels
115. A newborn is diagnosed with myelomeningocele and is being monitored in the NICU. The nurse is instructed to observe the anterior fontanelle as part of the neurological assessment. What is the most likely finding related to the anterior fontanelle if there is increased intracranial pressure? (Pediatrics)
- Sunken fontanelle
 - Normal fontanelle
 - Bulging fontanelle
 - Closed fontanelle
116. A 5-year-old child with asthma is prescribed 2 puffs of Asthalin (salbutamol) every 6 hours for 25 days by the pediatrician. During a follow-up visit, the nurse educates the parents on the proper use of the inhaler and potential side effects. What is the most important instruction the nurse should provide regarding the administration of Asthalin? (Pediatrics)
- Administer only during acute asthma attacks
 - Use a spacer device for effective delivery
 - Increase to 4 puffs if symptoms worsen
 - Administer before meals only

Answer Key

107. b 108. c 109. c 110. c 111. b 112. b 113. c 114. b 115. c 116. b



117. A 14-day-old neonate presents to the pediatric clinic with fast breathing, irritability, and poor feeding. The nurse suspects possible jaundice or infection. What is the most appropriate initial action to assess the neonate's condition? **(Pediatrics)**
- Administer oral antibiotics immediately
 - Perform a complete clinical assessment and check for jaundice or signs of infection
 - Start phototherapy without further evaluation
 - Advise the mother to increase breastfeeding frequency
118. A 4-year-old child in the pediatric ward is receiving high-flow oxygen at 50% due to respiratory distress. The nurse is tasked with ensuring proper oxygen therapy management. What is the most appropriate method to provide humidification for this child? **(Pediatrics)**
- Use a dry oxygen delivery system
 - Provide intermittent humidification with a heated humidifier
 - Administer saline nebulization every hour
 - Use a cold mist humidifier continuously
119. A 3-day-old neonate presents with jaundice, and the serum bilirubin level is 15 mg/dL. The pediatrician initiates phototherapy. Based on the expected bilirubin reduction rate of 1-2 mg/dL every 4-6 hours, what is the anticipated bilirubin level after 12 hours of effective phototherapy? **(Pediatrics)**
- 12-13 mg/dL
 - 9-12 mg/dL
 - 13-14 mg/dL
 - 10-11 mg/dL
120. A nurse is monitoring a preterm neonate in the NICU for signs of hypoxia. Which of the following clinical manifestations is a late sign of hypoxia in this patient? **(Pediatrics)**
- Tachypnea
 - Nasal flaring
 - Cyanosis
 - Restlessness
121. A nurse is providing discharge instructions to the parents of an infant with a jejunostomy who will be managed at home. Which of the following instructions should the nurse emphasize to ensure proper care of the jejunostomy site? **(Pediatrics)**
- Administer high-calorie formula orally to promote weight gain
 - Monitor the jejunostomy site daily for signs of infection or leakage
 - Clean the jejunostomy tube with alcohol-based solution twice daily
 - Restrict fluid intake to prevent excessive output from the jejunostomy

TARGET HIGH
DIGITAL

Answer Key

117. b 118. b 119. b 120. c 121. b

