
(a)
λ from the WS fit
λ sim
SNR 5
SNR 0.5
(b)
k0 from the WS fit (a.u.)
λ from the WS fit
λ sim = 1
λ sim = 2
λ sim = 4
(c)
estimated λ·k0 (a.u.)
true λ·k0 (a.u.)
λ·k0 from the WS fit
mean burst size
(d)
λ sim 1, SNR 0.5, n = 397
median 2.9% vs 2.3%
turnover beyond noise
3.9 [2.8, 5.1] %/h
Frequency
granules
burst-free copies
Burst size (% of granule)
(e)
λ sim 2, SNR 0.5, n = 474
median 3.5% vs 2.2%
turnover beyond noise
8.6 [6.7, 10.6] %/h
Burst size (% of granule)
(f)
λ sim 4, SNR 0.5, n = 632
median 5.1% vs 2.1%
turnover beyond noise
22.3 [17.9, 28.2] %/h
Burst size (% of granule)
(g)
Mean burst size, λ·k0 (a.u.)
SNR 5
SNR 0.5
true value
λ sim = 1
λ sim = 2
λ sim = 4
(h)
Turnover beyond noise (%/h)
λ sim = 1
λ sim = 2
λ sim = 4
(i)
Net change beyond bleaching (%/h)
λ sim = 1
λ sim = 2
λ sim = 4