
Rows: 4x RNA (top), 30x RNA (bottom). p: movie-level permutation (movies are the replicates). Largest 10% of bursts removed
(a) brightness: p = 0.0016
noise per frame 16% vs 22% (p = 0.002)
granule brightness (spot mean)
PP7, 4x RNA
PP7-FUS, 4x RNA
(b) burst size: granules p < 0.001
(naive burst-level test: p = 1e-11)
granules
burst-free copies (noise only)
median burst size (% of granule shed)
PP7, 4x RNA
PP7-FUS, 4x RNA
(c) where the gap comes from
total gap +5.9 points
FUS minus PP7, median burst size (% points)
set by brightness / noise (p = 0.0034)
beyond noise (p = 0.073)
(d) dynamics beyond noise: turnover p = 0.66
fluctuation p = 0.17, net change p = 0.14
turnover beyond noise (%/h, 95% CI)
PP7, 4x RNA
PP7-FUS, 4x RNA
(a) brightness: p = 0.55
noise per frame 21% vs 23% (p = 0.23)
granule brightness (spot mean)
PP7, 30x RNA
PP7-FUS, 30x RNA
(b) burst size: granules p = 0.056
(naive burst-level test: p = 0.007)
granules
burst-free copies (noise only)
median burst size (% of granule shed)
PP7, 30x RNA
PP7-FUS, 30x RNA
(c) where the gap comes from
total gap +2.2 points
FUS minus PP7, median burst size (% points)
set by brightness / noise (p = 0.071)
beyond noise (p = 0.88)
(d) dynamics beyond noise: turnover p = 0.91
fluctuation p = 0.67, net change p = 0.26
turnover beyond noise (%/h, 95% CI)
PP7, 30x RNA
PP7-FUS, 30x RNA