
Rows: 4x RNA (top), 30x RNA (bottom). p: movie-level permutation (movies are the replicates)
(a) brightness: p = 0.0016
noise per frame 16% vs 22% (p = 0.002)
granule brightness (spot mean)
PP7, 4x RNA
PP7-FUS, 4x RNA
(b) burst size: granules p < 0.001
(naive burst-level test: p = 8e-11)
granules
burst-free copies (noise only)
median burst size (% of granule shed)
PP7, 4x RNA
PP7-FUS, 4x RNA
(c) where the gap comes from
total gap +6.1 points
FUS minus PP7, median burst size (% points)
set by brightness / noise (p = 0.01)
beyond noise (p = 0.062)
(d) dynamics beyond noise: turnover p = 0.73
fluctuation p = 0.17, net change p = 0.14
turnover beyond noise (%/h, 95% CI)
PP7, 4x RNA
PP7-FUS, 4x RNA
(a) brightness: p = 0.55
noise per frame 21% vs 23% (p = 0.23)
granule brightness (spot mean)
PP7, 30x RNA
PP7-FUS, 30x RNA
(b) burst size: granules p = 0.1
(naive burst-level test: p = 0.02)
granules
burst-free copies (noise only)
median burst size (% of granule shed)
PP7, 30x RNA
PP7-FUS, 30x RNA
(c) where the gap comes from
total gap +1.9 points
FUS minus PP7, median burst size (% points)
set by brightness / noise (p = 0.19)
beyond noise (p = 0.78)
(d) dynamics beyond noise: turnover p = 0.84
fluctuation p = 0.67, net change p = 0.26
turnover beyond noise (%/h, 95% CI)
PP7, 30x RNA
PP7-FUS, 30x RNA