
CS 779 : TUTORIAL SOLUTIONS

Amit Rajaraman

Last updated January 28, 2023

Contents

1	Tutorial 1	2
2	Tutorial 2	3

§1. Tutorial 1

Exercise 1.1. Prove that the maximum number of subsets of $[n]$ with pairwise non-empty intersection is 2^{n-1} .

Solution

2^{n-1} is clearly attainable by taking $\{S \subseteq [n] : 1 \in S\}$. Furthermore, this is an upper bound since if \mathcal{S} is a family of subsets with pairwise non-empty intersection, then $\mathcal{S}' = \{S^c : S \in \mathcal{S}\}$ has zero intersection with \mathcal{S} and is of the same size, so $2|\mathcal{S}| = |\mathcal{S}'| + |\mathcal{S}| \leq 2^n$.

Exercise 1.2. Suppose you have a set system with m sets $(A_i)_{i=1}^m$ such that $|A_i|$ is odd for each i and $|A_i \cap A_j|$ is even for any $i \neq j$. Prove that $m \leq n$.

Solution

Consider the $m \times n$ matrix M where M_{ij} is 1 if $j \in A_i$ and is 0 otherwise. Then,

$$(MM^T)_{ij} = \sum_{k \in [n]} M_{ik}M_{jk} = |A_i \cap A_j|.$$

In particular, all the diagonal entries of MM^T are odd and all off-diagonal entries are even. Using this, it is not too difficult to show that $\det(MM^T) \neq 0$ (for an easy solution* of this, note that modulo 2, MM^T is congruent to the identity, which has nonzero determinant). Therefore, $m = \text{rank}(MM^T) = \text{rank}(M)$, so $m \leq n$.

Exercise 1.3. Prove that for matrices A, B , $\text{rank}(A + B) \leq \text{rank}(A) + \text{rank}(B)$.

Solution

It suffices to show that any column of $A + B$ is present in the space spanned by the column of A and B . This is straightforward since any column of $A + B$ is just the sum of the two corresponding columns in A and B .

Exercise 1.4. Suppose you have $A + A^T = J - I$, where J is the all ones matrix. Prove that $\text{rank}(A) \geq n/2$.

Solution

Using the previous exercise, we have $n = \text{rank}(J - I) = \text{rank}(A + A^T) \leq \text{rank}(A) + \text{rank}(A^T) = 2 \text{rank}(A)$.

Exercise 1.5. Suppose you have $A + A^T = J - I$, where J is the all ones matrix. Show that if $\text{rank}(A) < n - 1$, there is a vector x such that $Ax = 0$, $x \neq 0$, and $\mathbf{1}^T x = 0$. Using this, prove that $\text{rank}(A) \geq n - 1$.

Solution

Suppose $\text{rank}(A) < n - 1$. Then, $\dim \ker A \geq 2$. We also have $\dim \mathbf{1}^\perp = n - 1$. Therefore, $\ker A$ and $\mathbf{1}^\perp$ have nonzero intersection, and say $x \neq 0$ is in both. x satisfies the conditions mentioned in the question. Now,

$$\begin{aligned} 0 &= x^T(Ax) + (x^T A^T)x \\ &= x^T(J - I)x \\ &= \left(\sum_i x_i \right)^2 - \left(\sum_i x_i^2 \right) = - \sum_i x_i^2, \end{aligned}$$

so $x = 0$, a contradiction. Therefore, $\text{rank}(A) \geq n - 1$.

Exercise 1.6. Suppose B_1, \dots, B_m are complete bipartite graphs whose edge disjoint union yields the complete graph K_n . Show that $m \geq n - 1$.

Solution

Suppose that B_i corresponds to the complete bipartite graph between sets $X_i, Y_i \subseteq [n]$, where $X_i \cap Y_i = \emptyset$. As a graph on vertex set $[n]$, on setting $M_i = \mathbb{1}_X \mathbb{1}_Y^\top$, B_i has adjacency matrix $M_i + M_i^\top$. Note that $\text{rank}(M_i) = 1$ for all i , since $\mathbb{1}_Y^\perp \subseteq \ker M_i$. Because the edge disjoint union of the B_i is K_n , we have $(\sum_i M_i) + (\sum_i M_i)^\top = J - I$. Using the previous exercise, $\text{rank}(\sum_i M_i) \geq n - 1$. Using Exercise 1.3 and the observation that $\text{rank}(M_i) = 1$ for all i , this implies that $m = \sum_{i=1}^m \text{rank}(M_i) \geq n - 1$, completing the proof.

Exercise 1.7. Suppose you have a set system of m sets such that for every pair of sets, the intersection size is fixed as $\lambda \geq 1$. Prove that $m \leq n$.

Solution

Let the set system be $(A_i)_{i=1}^m$. The size of at most one set is equal to λ . Furthermore, if $|A_1| = \lambda$, then $A_i \setminus A_1$ are disjoint for distinct i , so $m - 1 \leq n - \lambda$. Thus, we may assume that the size of every set is greater than λ . Define the matrix M exactly as in Exercise 1.2. We have that the off-diagonal entries of M are equal to λ . Now, $MM^\top = \lambda J + D$, for some diagonal matrix D with all positive diagonal entries. We wish to show that $\text{rank}(\lambda J + D) = m$. Let $x \neq 0$ in \mathbb{R}^n , and let u, v be the components of x along and orthogonal to $\mathbf{1}$ respectively, such that $x = t\mathbf{1}$. Then,

$$\begin{aligned} (\lambda J + D)x &= (\lambda J + D)(u + v) \\ &= n\lambda u + D(u + v) \\ &= D(D^{-1}n\lambda u + u + v). \end{aligned}$$

When $t = 0$, this is clearly nonzero as $v \neq 0$. Otherwise, to conclude, note that

$$\sum_i (D^{-1}n\lambda u + u + v)_i = \sum_i (D_{ii}^{-1}n\lambda + 1)u_i + v_i = \sum_i t(D_{ii}^{-1}n\lambda + 1),$$

which is nonzero as $d_{ii}, \lambda > 0$ and $t \neq 0$.

§2. Tutorial 2

Exercise 2.1. Find the dimension of the space spanned by the following polynomials over the given field.

- (a) $x_1, x_2, x_1x_2, x_1^2x_2, 1, (x_1 + x_2)^2, x_1^2 + x_2^2$ over \mathbb{R} and over \mathbb{F}_2 .
 (b) $x_1^{i_1}x_2^{i_2} \cdots x_n^{i_n}$, where $i_1 + \cdots + i_n = m$ over \mathbb{R} and over \mathbb{F}_2 .

Solution

- (a) Over \mathbb{R} , it is clear that $(x_1 + x_2)^2 = (x_1^2 + x_2^2) + 2(x_1x_2)$, and the collection formed by removing $(x_1 + x_2)^2$ is linearly independent, so the dimension of the space is 6.
 We can again remove $(x_1 + x_2)^2$ in the \mathbb{F}_2 case. Note that $x^2 = x$ in \mathbb{F}_2 . Therefore, over \mathbb{F}_2 , the set of polynomials is $x_1, x_2, x_1x_2, x_1x_2, 1, x_1 + x_2$; it is clear a maximal linearly independent subset of these vectors is $x_1, x_2, x_1x_2, 1$, so the dimension is 4.
- (b) Over \mathbb{R} , all these monomials are linearly independent, so the dimension is the number of ways of choosing n non-negative numbers that sum to m . This is a routine exercise in combinatorics, with the answer being $\binom{m+n-1}{m}$.
 Over \mathbb{F}_2 , since $x_i = x_i^2$ for any i , we only care whether each i_r is zero or not. Let j_r be 1 if $i_r \geq 1$ and 0 otherwise. We then want to find the number of ways of choosing $(j_r)_{r=1}^n$ such that $0 < \sum_{r=1}^n j_r \leq m$. This is precisely $\binom{n}{1} + \binom{n}{2} + \cdots + \binom{n}{d}$, where $d = \min\{n, m\}$.

Exercise 2.2. Given m sets with sizes greater than d and pairwise intersection d , prove that $m \leq (n + 1)$.

Hint. Associate a polynomial to each set so that the polynomials are linearly independent. Give an upper bound on the space spanned by these polynomials.

Solution**Exercise 2.3.**

- (a) How do we define the distance between a pair of points in \mathbb{R}^n ?
- (b) Construct as many points as you can so that the distance between a pair is one of two distances, either d_1 or d_2 . You may also choose d_1 and d_2 to maximize the number.
- (c) Consider m points with exactly two pairwise distances. Associate polynomials $p_i(x)$ to each point such that the polynomials are linearly independent.
- (d) Deduce an upper bound on the dimension of the span of your polynomials. What does this imply about the number of points with exactly two pairwise distances?

Solution

Exercise 2.4. A polynomial is called multilinear if the degree of each variable is at most one. What is the dimension of the space of multilinear polynomials of degree at most d over n variables?

Solution

The solution to this is near-identical to the second part of Exercise 2.1(b), with the answer being $\binom{n}{0} + \binom{n}{1} + \dots + \binom{n}{d}$.

Exercise 2.5. Consider m sets A_1, \dots, A_m such that $|A_i| \equiv k \pmod{p}$ for some prime p . Assume that $|A_i \cap A_j| \in L \pmod{p}$ for some set L , such that $k \notin L$ and $|L| = \ell$. Show that $m \leq \binom{n}{0} + \dots + \binom{n}{\ell}$.

Solution