

---

# MA 5109: EXTREMAL GRAPH THEORY

---

Amit Rajaraman

Last updated October 8, 2021

## Contents

<b>0</b>	<b>Notation</b>	<b>2</b>
<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Basic Definitions	3
1.2	$K_{r+1}$ -free graphs	4
1.3	The Zarankiewicz Problem	7
1.3.1	The Zarankiewicz problem and the extremal function for complete bipartite graphs	9
1.3.2	The case where $s = t = 2$	9
1.3.3	The case where $s = t = 3$	10
1.4	$P_{k+1}$ -free graphs	12
<b>2</b>	<b>Fundamental Results in Extremal Graph Theory</b>	<b>14</b>
2.1	The Erdős-Stone-Simonovits Theorem	14
2.1.1	Motivation	14
2.1.2	The result	14
2.2	An Introduction to Random Graphs	19
2.2.1	A motivating extremal problem (bounding $\text{ex}(n; C_{2k})$ )	19
2.2.2	Digression: A coloring result of Erdős	20
2.3	Szemerédi's Regularity Lemma	22
2.3.1	Motivation	22
2.3.2	The result	23
2.4	Some Corollaries of Szemerédi's Regularity Lemma	26
2.4.1	The graph counting and removal lemmas	26
2.4.2	Roth's Theorem and Corners	28
2.4.3	A weaker version of the Erdős-Stone-Simonovits Theorem	30
2.4.4	An application in computer science	31
2.4.5	The Frankl-Pach Theorem	31
2.4.6	A look at $\text{ex}(n; C_{2k+1})$	33
2.5	Pseudorandomness	35
2.5.1	Notions of randomness	35
2.5.2	Strongly regular graphs	39
2.5.3	The sparse regime	40
<b>3</b>	<b>Matching Theory</b>	<b>44</b>
3.1	The Bipartite Setting	44
3.1.1	Hall's Marriage Theorem	44
3.1.2	Some applications	45

## §0. Notation

We use  $[n]$  to represent  $\{1, 2, \dots, n\}$ .

For integers  $a$  and  $b$ ,  $[a, b]$  means  $\{a, a + 1, \dots, b\}$ .

A graph  $G_n$  is a graph with  $n$  vertices.

Given a graph  $G$ ,  $e(G)$  is the number of edges  $G$  has.

For a vertex  $v$ , denote by  $N(v)$  or  $\Gamma(v)$  the set of *neighbours* of  $v$  – all the vertices that have an edge to  $v$ .

For a vertex  $v$ , denote by  $d_G(v) = |\Gamma(v)|$  the *degree* of  $v$  – the number of edges incident on it. If the graph  $G$  is clear from context, we write simply  $d(v)$ .

For  $v \in V$  and  $K \subseteq V$ ,  $d(v, K)$  is the number of edges

$$|\{u \in K : uv \in E\}|$$

from  $v$  into  $K$ .

For vertices  $v, w$ , write  $d(v, w) = |\Gamma(v) \cap \Gamma(w)|$ .

Given a graph  $G = (V, E)$ , denote by  $\delta(G)$  and  $\Delta(G)$  the minimum and maximum degree in  $G$  respectively. That is,

$$\delta(G) = \min_{v \in V} d(v) \text{ and } \Delta(G) = \max_{v \in V} d(v).$$

We shorten “there exists  $n_0$  such that for all  $n > n_0$ ” to “for all  $n$  sufficiently large”, which in turn is shortened to “for all  $n \gg 0$ ”.

We say that something occurs “with high probability” if the probability of the complement occurring converges to 0 as the input size grows to  $\infty$ .

## §1. Introduction

### 1.1. Basic Definitions

**Definition 1.1.** A (simple undirected) **graph**  $G$  is an ordered pair  $(V, E)$  where  $V$  is a finite set called the *vertex set* and  $E$ , called the *edge set*, is a subset of  $\binom{V}{2}$ , where  $\binom{S}{k}$  represents the set of all  $k$ -element subsets of  $S$ .

We typically represent graphs pictorially, showing vertices as dots and edges as arcs joining the vertices present in the corresponding subset.

A few important graphs are:

- the *null graph* with vertex set  $V$ , where  $E = \emptyset$ .
- the *complete graph*  $K_n$ , where  $V = [n]$  and  $E = \binom{[n]}{2}$ .
- the *complete bipartite graph*  $K_{m,n}$ , where  $V = A \cup B$  with  $|A| = m$ ,  $|B| = n$ , and  $A, B$  are disjoint, and  $E = \{\{a, b\} : a \in A, b \in B\}$ .
- the *path graph*  $P_{n+1}$  of length  $n$ , where  $V = [n+1]$  and  $E = \{\{m, m+1\} : m \in [n]\}$ .
- the *cycle* of length  $n$ , where  $V = [n]$  and  $E = \{\{l, m\} : l, m \in [n], (m-l) \equiv 1 \pmod{n}\}$ .

Now, consider the graph  $G$  with vertex set  $[4]$  and edge set  $\{\{1, 3\}, \{3, 2\}, \{2, 4\}\}$ . This graph appears to be the same as the path graph of length 3, but how do we make this correspondence more concrete?

Relabeling vertices doesn't create a "new" graph.

**Definition 1.2** (Graph Isomorphism). Two graphs  $G = (V, E)$  and  $G' = (V', E')$  are said to be **isomorphic** and we write  $G \simeq G'$  if there exists a bijection  $f : V \rightarrow V'$  such that there is an edge between two vertices  $u$  and  $v$  in  $G$  if and only if there is an edge between  $f(u)$  and  $f(v)$  in  $G'$ .

If two graphs are isomorphic, they are identical for our purposes (we only care about graphs up to isomorphism). We now give a few more definitions that are useful.

**Definition 1.3** (Subgraph). Given a graph  $G = (V, E)$ , a **subgraph**  $H = (V', E')$  is a graph such that  $V' \subseteq V$  and  $E' \subseteq E$ . Given  $V' \subseteq V$ , the subgraph *induced* by  $V'$  on  $G$  is that with vertex set  $V'$  and edge set  $\binom{V'}{2} \cap E$ .

**Definition 1.4** ( $r$ -partite Graph). A graph  $G = (V, E)$  is said to be  **$r$ -partite** if there exists a partition  $V_1, V_2, \dots, V_r$  of  $V$  such that for any edge  $e = uv \in E$ ,  $u$  and  $v$  are in distinct  $V_i$ . That is, there are no edges within any of the  $V_i$ . In particular, a 2-partite graph is said to be **bipartite**.

**Definition 1.5** (Independent Set). Given a graph  $G = (V, E)$ ,  $I \subseteq V$  is said to be **independent** if no two vertices of  $I$  are adjacent (the subgraph induced by  $I$  is null).

$\alpha(G)$ , the *independence number* of  $G$ , denotes the size of the largest independent set in  $G$ .

**Definition 1.6** (Clique). Given a graph  $G = (V, E)$ ,  $K \subseteq V$  is said to be a **clique** if any two vertices of  $K$  are adjacent (the subgraph induced by  $K$  is complete).  $\omega(G)$ , the *clique number* of  $G$ , denotes the size of the largest clique in  $G$ .

**Definition 1.7** (Complement Graph). Given a graph  $G = (V, E)$ , the **complement graph** of  $G$  is  $\bar{G} = (V, \binom{V}{2} \setminus E)$ .

Observe that  $S \subseteq V$  is independent in  $G$  if and only if  $S$  is a clique in  $\bar{G}$ . In particular,  $\alpha(G) = \omega(\bar{G})$ .

**Definition 1.8** (Connectedness). A graph  $G$  is said to be **connected** if for any pair of vertices  $u, v$ , there is a sequence  $u = v_0, v_1, \dots, v_r = v$  for some  $r$  such that  $v_{i-1}v_i$  is an edge for each  $i \in [r]$ .

**Definition 1.9** (Girth). The **girth** of a graph  $G$  is the smallest  $k (> 2)$  for which  $C_k$  is isomorphic to a subgraph of  $G$ .

If  $G$  has no cycles, it is said to have infinite girth.

## 1.2. $K_{r+1}$ -free graphs

Extremal graph theory is motivated by the following simple problem:

At most how many edges can a graph  $G_n$  have if it contains no triangles?

More precisely, what is

$$\max_{\substack{\text{no subgraph of } G_n \\ \text{is isomorphic to } K_3}} e(G_n)?$$

Clearly, this number is well-defined since a graph on  $n$  vertices cannot have more than  $\binom{n}{2}$  edges.

A simple observation is that any complete bipartite graph has no triangles: if there were a triangle, then two vertices would be in the same “part”, which contradicts the existence of edges only between the two parts.

As a consequence, for any  $1 \leq m \leq n$ , it is possible to construct  $m \times (n - m)$  edges (with this bound being attained for  $K_{m, n-m}$ ). In particular, it is possible to construct a graph with  $\lfloor n^2/4 \rfloor$  edges.

**Theorem 1.1** (Mantel’s Theorem). If  $G_n$  has no triangle, then

$$e(G_n) \leq \left\lfloor \frac{n^2}{4} \right\rfloor.$$

Further, equality is attained iff  $G_n \simeq K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$ .

*Proof.* Suppose  $G_n$  has no triangles. Saying that  $G_n$  has no triangles is equivalent to saying that for distinct adjacent  $u, v$ ,  $\Gamma(u) \cap \Gamma(v) = \emptyset$ .

So,  $d(u) + d(v) \leq n$ . Therefore,

$$\begin{aligned}
 ne(G_n) &\stackrel{(1)}{\geq} \sum_{uv \in E} d(u) + d(v) \\
 &= \sum_{uv \in E} |\Gamma(u) \cup \Gamma(v)| \\
 &= |\{(e, w) : e = uv \in E, w \in \Gamma(u) \cup \Gamma(v)\}| \\
 &= \sum_{u \in V} |\{(e, w) : w \in \Gamma(u), e = uv \in E\}| \\
 &= \sum_{u \in V} |\{(v, w) : v, w \in \Gamma(u)\}| \\
 &\stackrel{(2)}{=} \sum_{u \in V} d(u)^2 \\
 &\stackrel{(3)}{\geq} \frac{1}{n} \left( \sum_{u \in V} d(u) \right)^2 \\
 &\stackrel{(4)}{=} \frac{4e(G_n)^2}{n},
 \end{aligned}$$

where (2) follows from the changing the main thing being summed over to  $u$ , the “middle” vertex in the  $L$ -like structure, (3) follows from the **Cauchy-Schwarz inequality**, and (4) follows from the **handshaking lemma**.

What happens when equality is attained? Let us look at the case where  $n$  is even.

(1) is only tight when  $d(u) + d(v) = n$  for all edges  $uv$  and (3) is only tight when  $d(u)$  is a constant (independent of  $u$ ). This implies that  $d(u) = \frac{n}{2}$  for every  $u \in V$ . Now, if  $uv$  is an edge,  $\Gamma(u) \cap \Gamma(v) = \emptyset$  implies that  $\Gamma(u) \cup \Gamma(v) = V$ , and so  $G_n = K_{\frac{n}{2}, \frac{n}{2}}$ .

The case where  $n$  is odd is analyzed similarly, with slight nuances in (3) since exact equality is not attained. ■

While the above is one of the early results in extremal graph theory, the subject was only really born due to Turán in the following result.

**Theorem 1.2 (Turán’s Theorem).** If  $G_n$  has no  $K_{r+1}$  ( $r \geq 2$ ), then  $e(G_n) \leq t_r(n)$ , with equality attained iff  $G \simeq T_r(n)$ .

The version for  $r = 2$  is just a triangle-free graph and is the same as **Mantel’s Theorem**. In the proof of this, we split the vertex set into two parts and dumped all the edges between these parts.

If we want to avoid  $K_4$  ( $r = 3$ ), then perhaps we could split the vertex set into three parts and dump all the edges between these parts.

In general, we want to partition  $V$  of size  $n$  into  $r$  “almost equal” parts and set only those edges between vertices in distinct parts – such a graph is known as the **Turán graph**  $T_r(n)$  and the number of edges  $e(T_r(n))$  is the **Turán number**  $t_r(n)$ .

In particular, when  $r \mid n$ ,

$$t_r(n) = \binom{r}{2} \left( \frac{n}{r} \right)^2 = \frac{n^2}{2} \left( 1 - \frac{1}{r} \right).$$

Here, we give three proofs of Turán’s Theorem.

*Proof of Turán’s Theorem.* We perform strong induction on  $n + r$ . We have already proved the result for  $r = 2$ .

Suppose  $e(G_n) \geq t_r(n)$  and  $G_n$  is  $K_{r+1}$ -free, where  $r > 2$ . We wish to prove that  $G \simeq T_r(n)$ .

Since  $t_r(n) > t_{r-1}(n)$ <sup>1</sup>, the inductive hypothesis implies that  $G$  has a copy  $K \subseteq V$  of  $K_r$ . Observe that for  $v \notin K$ ,

<sup>1</sup>To show this, use the fact that any  $(r - 1)$ -partite graph can be thought of as  $r$ -partite graph with one of the pieces being empty, and then that the Turán graph has the most edges among  $r$ -partite graphs.

$d(v, K) \leq r - 1$  – otherwise, there would be a copy of  $K_{r+1}$  in  $G$ .

As a result,  $e(V \setminus K, K) \leq (r - 1)(n - r)$ . By the induction hypothesis,  $e(V \setminus K, V \setminus K) \leq t_r(n - r)$ . Therefore,

$$t_r(n) \leq e(G_n) \leq t_r(n - r) + (r - 1)(n - r) + \binom{r}{2}.$$

However, as can be checked manually,  $t_r(n - r) + (r - 1)(n - r) + \binom{r}{2} = t_r(n)$ !

It follows that equality holds everywhere –  $e(G_n) = t_r(n)$ ,  $e(V \setminus K) = t_r(n - r)$ , and  $d(v, K) = r - 1$  for all  $v \in V \setminus K$ . This graph is then isomorphic to  $T_r(n)$  – for each  $v \in V \setminus K$ , we can put the vertex in  $K$  that is not adjacent to  $v$  in the same bucket as  $v$ . Then, the only edges are those between distinct buckets (Why?), so  $G_n \simeq T_r(n)$ . ■

*Erdős' Proof of Turán's Theorem.* Erdős proves a slightly more general claim: given a  $K_{r+1}$ -free graph  $G_n$ , there exists an  $r$ -partite graph  $H$  on  $V$  such that  $d_G(v) \leq d_H(v)$  for all  $v \in V$ .

It is then a simple task to check that among the  $r$ -partite graphs on  $n$  vertices, the Turán graph  $T_r(n)$  has the most edges.

To prove our claim, we perform induction on  $r$ .

The claim is trivial for the base case  $r = 1$ .

Now, suppose the claim holds for values less than  $r$ . Let  $v_0 \in V$  such that  $d_G(v_0) = \max_{v \in V} d_G(v)$  (the vertex of maximum degree in  $G$ ) and  $W = \Gamma(v_0)$ . Since  $G$  is  $K_{r+1}$ -free,  $W$  is  $K_r$ -free. Inductively, there is an  $(r - 1)$ -partite graph  $H'$  on  $W$  such that for all  $v \in W$ ,  $d_{H'}(v) \geq d_W(v)$ .

Let  $U = V \setminus W$ . For each  $u \in U$ , remove all its edges in  $G$  and set its new neighbour set as  $W$ .

Our desired graph  $H$  is that with these edges along with those in  $H'$  and the edges from  $v_0$  to  $W$ . That is, the  $r$ th part is  $U \cup \{v_0\}$  and the remaining  $(r - 1)$  parts are those formed by  $H'$ . The graph is clearly  $r$ -partite by definition. What about the degree inequality?

- $d_G(v_0) = d_H(v_0)$  trivially.
- For  $u \in U$ ,  $d_H(u) = d_G(v_0) \geq d_G(u)$ .
- For  $w \in W$ ,

$$d_H(w) = |U| + 1 + d_{H'}(w) \geq |U| + 1 + d_W(w) \geq d_U(w) + 1 + d_W(w) = d_G(w). \quad \blacksquare$$

(Why does equality imply that the graph is isomorphic to  $T_r(n)$ ?)

**Theorem 1.3** (Turán's Theorem, reformulation). If  $d = e(G_n)/n$  is the average degree of the vertices of  $G_n$ , then  $G_n$  has an independent set of size at least  $n/(d + 1)$ .

*Proof.* Why is this equivalent to Turán's Theorem?

If  $G_n$  has no  $K_{r+1}$ , then  $\alpha(\bar{G}) \leq r$ . If  $G_n$  has average degree  $d$ , the above result would imply that  $r \geq n/(d + 1)$ , that is,  $d \leq (n/r) - 1$ . The total number of edges in  $G_n$  is then

$$\binom{n}{2} - \frac{nd}{2} \leq \binom{n}{2} - \frac{n}{2} \left( \frac{n}{r} - 1 \right) = \frac{n^2}{2} \left( 1 - \frac{1}{r} \right),$$

which gives Turán's bound!

Let us now get to the proof of the above reformulation. First, consider the following algorithm to come up with *some* independent set in  $G$ :

1. Order  $V$  to get  $\{v_1, \dots, v_n\}$  and initialize  $S = \emptyset$ .
2. Add  $v_1$  to  $S$ .

3. Having processed vertices  $v_1$  through  $v_i$ , add  $v_{i+1}$  to  $S$  iff there is no vertex in  $S$  that is adjacent to  $v_{i+1}$ .

It is clear that this always produces an independent set, but the size of the independent set depends on the ordering we choose at the beginning.

For a given ordering  $\sigma$ , denote by  $\mathcal{A}(\sigma)$  the independent set produced by the algorithm.

How do we choose a “good” ordering?

Enter the probabilistic method. Define the random variable  $\pi$  to be uniformly random on the set of all permutations of  $V$ . Then,

$$\begin{aligned} \mathbf{E}[|\mathcal{A}(\pi)|] &= \mathbf{E}\left[\sum_{v \in V} \mathbb{1}_{v \in \mathcal{A}(\pi)}\right] \\ &= \sum_{v \in V} \mathbf{E}\left[\mathbb{1}_{v \in \mathcal{A}(\pi)}\right] \\ &= \sum_{v \in V} \Pr[v \in \mathcal{A}(\pi)]. \end{aligned}$$

Fix some  $v \in V$  and permutation  $\sigma$ . What is the probability that  $v \in \mathcal{A}(\sigma)$ ?

If at the time of processing  $v$  for the ordering  $\sigma$ ,  $\Gamma(v) \cap S \neq \emptyset$ , then  $v$  is not picked. In particular, if  $v$  is the first element of  $\Gamma(v) \cup \{v\}$  in the ordering  $\sigma$ , then it is definitely chosen by the algorithm. The probability of this occurring is  $\frac{1}{d(v)+1}$ . So,

$$\begin{aligned} \mathbf{E}[|\mathcal{A}(\pi)|] &= \sum_{v \in V} \Pr[v \in \mathcal{A}(\pi)] \\ &\geq \sum_{v \in V} \frac{1}{d(v)+1} \\ &\stackrel{(*)}{\geq} \frac{n^2}{\sum_{v \in V} (d(v)+1)} = \frac{n}{d+1}, \end{aligned}$$

where  $(*)$  follows from the **AM-HM inequality**.

Since the expectation of  $|\mathcal{A}(\pi)|$  is at least  $n/(d+1)$ , there must exist some permutation  $\sigma$  such that  $|\mathcal{A}(\sigma)| \geq n/(d+1)$ , proving the result. ■

### 1.3. The Zarankiewicz Problem

Turán's Theorem is the primary result that birthed Extremal Graph Theory. To generalize the problem studied in the previous section, define the following.

**Definition 1.10** (Extremal Function). Given a graph  $H$ , define the **extremal function**

$$\text{ex}(n; H) = \max_{\substack{\text{no subgraph of } G_n \\ \text{is isomorphic to } H}} e(G_n) \tag{1.1}$$

as the maximum number of edges in a graph on  $n$  vertices without  $H$  as a subgraph.

More generally, if  $\mathcal{F}$  is a family of graphs, define

$$\text{ex}(n; \mathcal{F}) = \max_{\substack{\text{no subgraph of } G_n \\ \text{is isomorphic to a } H \in \mathcal{F}}} e(G_n).$$

In particular, given graphs  $H_1, \dots, H_m$ , we denote

$$\text{ex}(n; H_1, \dots, H_m) = \text{ex}(n; \{H_1, \dots, H_m\}).$$

With this notation, Turán's Theorem then says that  $\text{ex}(n; K_{r+1}) = t_r(n)$ , with the corresponding maximum in (1.1) being attained iff  $G_n \simeq T_r(n)$ .

**Definition 1.11** (Zarankiewicz Function). Fix  $s, t \in \mathbb{N}$  with  $t \geq s \geq 2$  and  $m, n \in \mathbb{N}$ . The **Zarankiewicz function**  $z(m, n; s, t)$  is the maximum number of edges in a bipartite graph  $G = (A \sqcup B, E)$  such that

- the two components  $A$  and  $B$  of the graph are of cardinality  $m$  and  $n$  respectively<sup>2</sup>, and
- there exist no  $S \subseteq A, T \subseteq B$  with  $|S| = s, |T| = t$ , and all edges between  $S$  and  $T$  present in  $E$ .<sup>3</sup>

For ease of writing, we refer to the above described criterion as the *Zarankiewicz condition*.

That is, we “forbid” the subgraph  $K_{s,t}$  with the components of size  $s$  and  $t$  on the side of  $A$  and  $B$  respectively.

The **Zarankiewicz problem** asks for a closed form representation of  $z(m, n; s, t)$ . Failing this, for fixed  $t$ , it asks for a tight asymptotic bound on  $z(n, n; t, t)$  as  $n$  grows large.

Perhaps surprisingly, this problem remains unsolved! (as of the time of writing these notes)

**Theorem 1.4** (Kővári-Sós-Turán Theorem). For  $t \geq s \geq 2$  and  $m, n \in \mathbb{N}$ ,

$$z(m, n; s, t) \leq (s-1)^{1/t}(n-t+1)m^{1-1/t} + (t-1)m.$$

*Proof.* Let  $G$  be bipartite with vertex set  $A \sqcup B$  and satisfy the Zarankiewicz condition.

Fix  $a \in A$ . By definition,  $|\Gamma(a)| = d(a)$ . Now,  $\binom{d(a)}{t}$  is the number of  $t$ -element subsets of  $\Gamma(a)$ . We may assume that for all  $a \in A$ ,  $d(a) \geq t-1$ . Indeed, otherwise, we may add arbitrary edges to  $a$  to make its degree  $t-1$ ; any vertex from  $A$  in a subgraph isomorphic to  $K_{s,t}$  must have at least degree  $t$  so  $a$  cannot be part of it.

We have that

$$\begin{aligned} \sum_{x \in A} \binom{d(x)}{t} &= |\{(x, T) : x \in A, T \subseteq \Gamma(x), |T| = t\}| \\ &= \sum_{\substack{T \subseteq B \\ |T|=t}} |\{x \in A : T \subseteq \Gamma(x)\}|. \end{aligned}$$

If we fix a  $T$ , then the number of such  $x$  for that  $T$  is at most  $s-1$ , due to our assumption. As a result,

$$\sum_{x \in A} \binom{d(x)}{t} \leq \binom{n}{t}(s-1).$$

Now, observe that the function

$$f(x) = \frac{x(x-1) \cdots (x-t+1)}{t!}$$

is convex on  $[t-1, \infty)$ . Using **Jensen's inequality** together with our assumption that  $d(x) \geq t-1$  for all  $x \in A$ ,

$$\begin{aligned} \binom{n}{t}(s-1) &\geq \sum_{x \in A} \binom{d(x)}{t} \\ &\geq m \binom{\frac{1}{m} \sum_{x \in A} d(x)}{t} \\ &= m \binom{e(G)/m}{t}. \end{aligned} \tag{1.2}$$

<sup>1</sup>by “components” of the bipartite graph we mean that for any edge  $uv$  in the graph,  $u \in A$  and  $v \in B$  or  $u \in B$  and  $v \in A$ .

<sup>2</sup>we make the tacit assumption that  $s \leq m$  and  $t \leq n$ .



Let  $d = e(G)/m$ , the average degree of the vertices in  $A$ . Simplifying the above expression,

$$\frac{s-1}{m} \geq \frac{d(d-1) \cdots (d-t+1)}{n(n-1) \cdots (n-t+1)} \geq \left( \frac{d-t+1}{n-t+1} \right)^t,$$

where the second inequality follows from the fact that  $d \leq n$ . Therefore,

$$e(G) \leq m \left( \left( \frac{s-1}{m} \right)^{1/t} (n-t+1) + (t-1) \right) = (s-1)^{1/t} (n-t+1) m^{1-1/t} + (t-1)m,$$

completing the proof. ■

Next, let us look at some consequences of the above bound.

### 1.3.1. The Zarankiewicz problem and the extremal function for complete bipartite graphs

We get a bound on  $\text{ex}(n; K_{s,t})$ . We claim that for  $n \in \mathbb{N}$  and  $s, t \geq 2$ ,

$$\text{ex}(n; K_{s,t}) \leq \frac{1}{2} z(n, n; s, t). \quad (1.3)$$

Indeed, if  $G_n = (V, E)$  has no  $K_{s,t}$ , make a bipartite graph  $G'$  that has vertex set  $V \times \{0, 1\}$  such that  $uv$  is an edge in  $G$  iff  $\{(u, 0), (v, 1)\}$  is an edge in  $G'$ .

$G'$  satisfies the Zarankiewicz condition. If there do exist  $S \subseteq V \times \{0\}$  and  $T' \subseteq V \times \{1\}$  such that all  $S$ - $T'$  edges are in  $G$ , then  $\pi_1(S) \cap \pi_1(T') = \emptyset$  (otherwise, a vertex would be adjacent to itself in  $G$ , which is clearly false). This implies that  $K_{s,t} \subseteq G$ , which is a contradiction.

Since  $e(G') = 2e(G)$ , the claim follows.

### 1.3.2. The case where $s = t = 2$

When  $s = t = 2$ , we get

$$z(m, n; 2, 2) \leq (n-1)m^{1/2} + m \text{ and } z(n, n; 2, 2) \leq (n-1)n^{1/2} + n.$$

Therefore,

$$\text{ex}(n; K_{2,2}) \leq \frac{1}{2} (n + (n-1)\sqrt{n}).$$

Note that  $K_{2,2} \simeq C_4$ . Therefore, a square-free graph on  $n$  vertices has  $\mathcal{O}(n^{3/2})$  edges.

In fact, this bound is tight! We give an algebraic construction of a suitable graph with no  $K_{2,2}$ , known as the *Levi graph for the projective plane*.

Let  $q$  be a prime power and consider the 3-dimensional vector space  $\mathcal{V} = \mathbb{F}_q^3$  (over  $\mathbb{F}_q$ ). Let  $\mathcal{P}$  and  $\mathcal{L}$  be the set of all 1- and 2-dimensional subspaces of  $\mathcal{V}$  respectively.

Define the graph  $G = (\mathcal{P} \sqcup \mathcal{L}, E)$  as follows. For  $x \in \mathcal{P}$  and  $L \in \mathcal{L}$ , let  $x$  be adjacent to  $L$  in  $G$  iff  $x \subseteq L$ .

We claim that  $G$  has no  $K_{2,2}$ . Suppose otherwise and let  $x_1, x_2 \in \mathcal{P}$  and  $L_1, L_2 \in \mathcal{L}$  such that the  $x_i$  are adjacent to the  $L_j$ . If  $x_1 = \langle u \rangle$  and  $x_2 = \langle v \rangle$ , then  $u$  and  $v$  are linearly independent, which implies that  $L_i = \langle u, v \rangle$ . This contradicts the fact that the  $L_i$  are distinct!

What are the cardinalities of  $\mathcal{P}$  and  $\mathcal{L}$ ?

- To get a 1-dimensional subspace, we pick a non-zero  $u$  and consider  $\langle u \rangle$ . In  $\mathcal{V}$ , there are  $q^3 - 1$  non-zero  $u$ . We must now divide by  $q-1$  to account for the fact that linearly dependent vectors generate the same 1-dimensional subspace. Any non-zero  $u$  has precisely  $q-1$  non-zero multiples. Therefore,

$$|\mathcal{P}| = \frac{q^3 - 1}{q - 1} = q^2 + q + 1.$$

- It turns out that the number of 2-dimensional subspace is equal to the number of 1-dimensional subspaces (more generally, the number of  $d$ -dimensional subspaces is equal to the number of  $(n - d)$ -dimensional subspaces of  $\mathbb{F}_q^n$ ), so

$$|\mathcal{L}| = q^2 + q + 1.$$

How many edges does  $G$  have?

Fix  $x = \langle u \rangle \in \mathcal{P}$ . We wish to determine how many  $L \in \mathcal{L}$  are adjacent to  $x$  in  $G$ . Such an  $L$  can be created by choosing  $v \notin x$  and letting  $L = \langle u, v \rangle$ .

The number of choices of  $v$  is  $q^3 - q$ , but each such subspace is repeated for  $q^2 - q$  choices of  $v$  since the cardinality of  $\langle u, v \rangle$  is  $q^2$ . Therefore,

$$d(x) = \frac{q^3 - q}{q^2 - q} = q + 1.$$

The total number of edges is

$$|\mathcal{P}|(q + 1) = (q + 1)(q^2 + q + 1) = q^3 + 2q^2 + 2q + 1,$$

which is  $\mathcal{O}(|\mathcal{P}|^{3/2})$ ! Therefore, our  $\mathcal{O}(n^{3/2})$  bound is tight.

In fact, the Levi graph is optimal in the case where  $n = 2(q^2 + q + 1)$ , as seen in Corollary 1.5.

**Corollary 1.5.** For  $n \in \mathbb{N}$ ,

$$z(n, n; 2, 2) \leq \frac{n(1 + \sqrt{4n - 3})}{2}. \quad (1.4)$$

Consequently,

$$\text{ex}(n; C_4) \leq \frac{1}{4}n(1 + \sqrt{4n - 3}). \quad (1.5)$$

*Proof.* Equation (1.5) clearly follows from Equations (1.3) and (1.4), so it suffices to show the first equation. Equation (1.2) in the proof of the **Kővári-Sós-Turán Theorem** for the case where  $s = t = 2$  just says that

$$\binom{n}{2} \geq n \binom{d}{2},$$

where  $d = e(G)/m$ . That is,  $d^2 - d - (n - 1) \leq 0$ . Then,

$$d \leq \frac{1 + \sqrt{1 + 4(n - 1)}}{2} = \frac{1 + \sqrt{4n - 3}}{2},$$

which is exactly the bound we want. This bound is tight in the case where  $n = 2(q^2 + q + 1)$ , as seen in the Levi graph. ■

Before we move on to the next section, let us build a tiny bit of intuition for why the construction of the Levi graph works.

The projective plane is chosen to ensure that any two distinct points determine a unique line (which holds even in the non-projective plane setting), and any two distinct lines determine a unique point. This corresponds to the absence of  $K_{2,2}$  – if it *was* present as a subgraph, then there would be two lines (points) that determine two points (lines), which cannot happen!

### 1.3.3. The case where $s = t = 3$

We next look at  $\text{ex}(n; K_{3,3})$ .

The **Kővári-Sós-Turán Theorem** applied here gives

$$z(n, n; 3, 3) \leq (2)^{1/3}(n - 2)n^{2/3} + 2n,$$

which is  $\mathcal{O}(n^{5/3})$ .

Similar to the Levi graph, we construct an (algebraic) extremal example. Let  $p$  be a prime and fix some  $r \in \mathbb{F}_p$ . Consider the graph  $G$  that has vertex set  $\mathcal{V} = \mathbb{F}_p^3$  where  $(x, y, z)$  is adjacent to  $(u, v, w)$  iff

$$(x - u)^2 + (y - v)^2 + (z - w)^2 = r. \quad (1.6)$$

Before moving on to explaining why this works, let us try to impart some intuition. (1.6) resembles the equation of a sphere in  $\mathbb{R}^3$ . If we take three spheres of the same radius, the points of intersection of all three must lie on two circles, corresponding to the circles of intersection of two pairs of spheres. Since any two circles meet at at most two points, the absence of  $K_{3,3}$  follows.

It may be shown that even over  $\mathbb{F}_p$ , two “spheres” intersect on a “circle” and any two circles meet at at most 2 points (Check this!).

So, if we have a  $K_{3,3}$  in the described graph, we have three spheres (centered at each of the three points) that intersect at three points, which is not possible.

It remains to count the number of edges in this graph. To do so, let us estimate the degree of  $(0, 0, 0)$ , since all vertices have the same degree (Why?). That is, we want to determine

$$|\{(x, y, z) \in \mathbb{F}_p^3 : x^2 + y^2 + z^2 = r\}|.$$

Letting  $z$  be arbitrary, we want to find

$$N(\xi) = |\{(x, y) \in \mathbb{F}_p^2 : x^2 + y^2 = \xi\}|$$

for any arbitrary  $\xi \in \mathbb{F}_p$ .

**Definition 1.12** (Legendre Symbol). For  $a \in \mathbb{F}_p$ , define the **Legendre symbol**

$$\left(\frac{a}{p}\right) = \begin{cases} 1, & a \in \mathbb{F}_p^\times \text{ is a square,} \\ -1, & a \in \mathbb{F}_p^\times \text{ is not a square,} \\ 0, & a = 0. \end{cases}$$

With the above, it is not too difficult to see that

$$N(\xi) = \sum_{(a,b): a+b=\xi} \left(1 + \left(\frac{a}{p}\right)\right) \left(1 + \left(\frac{b}{p}\right)\right).$$

Let us now compute the above quantity.

- First of all,

$$\sum_{(a,b): a+b=\xi} 1 = p.$$

- Observe that

$$\sum_{a \in \mathbb{F}_p} \left(\frac{a}{p}\right) = 0.$$

Indeed, the number of squares and non-squares in  $\mathbb{F}_p^\times$  is the same. As a result,

$$\sum_{(a,b): a+b=\xi} \left(\frac{a}{p}\right) = \sum_{a \in \mathbb{F}_p} \left(\frac{a}{p}\right) = 0.$$

Therefore,

$$N(\xi) = p + \sum_{(a,b): a+b=\xi} \left(\frac{a}{p}\right) \left(\frac{b}{p}\right). \quad (1.7)$$

Notice that the map  $\mathbb{F}_p^\times \rightarrow \{-1, 1\}$  given by  $x \mapsto \left(\frac{x}{p}\right)$  is a group homomorphism.

**Lemma 1.6.** If  $\xi \neq 0$ ,  $N(\xi) = N(1)$ .

*Proof.* Using (1.7) and letting  $a' = a/\xi$  and  $b' = b/\xi$ ,

$$\begin{aligned}
 N(\xi) - p &= \sum_{a+b=\xi} \binom{a}{p} \binom{b}{p} \\
 &= \sum_{a'+b'=1} \binom{a'\xi}{p} \binom{b'\xi}{p} \\
 &= \sum_{a'+b'=1} \binom{a'}{p} \binom{b'}{p} \left( \frac{\xi^2}{p} \right) \\
 &= \sum_{a'+b'=1} \binom{a'}{p} \binom{b'}{p} = N(1) - p.
 \end{aligned}$$

So, we have

$$\begin{aligned}
 (p-1)(N(1) - p) &= \sum_{\xi \in \mathbb{F}_p^\times} \sum_{(a,b): a+b=\xi} \binom{a}{p} \binom{b}{p} \\
 &= \sum_{\xi \in \mathbb{F}_p^\times} \sum_{a \in \mathbb{F}_p} \binom{a}{p} \binom{\xi - a}{p} \\
 &= \sum_{a \in \mathbb{F}_p} \left( \binom{a}{p} \sum_{\xi \in \mathbb{F}_p^\times} \binom{\xi - a}{p} \right) \\
 &= \sum_{a \in \mathbb{F}_p} \binom{a}{p} \left( 0 - \binom{-a}{p} \right) \\
 &= - \left( \frac{-1}{p} \right) \sum_{a \in \mathbb{F}_p} \binom{a^2}{p} \quad \text{(using the group homomorphism property)} \\
 &= -(p-1) \left( \frac{-1}{p} \right)
 \end{aligned}$$

and therefore,  $N(1) = p - \left( \frac{-1}{p} \right)$ .

If we choose  $r$  to be a non-square, then  $r - z^2 \neq 0$  for any  $z \in \mathbb{F}_p$  and  $N(r - z^2) = p - \left( \frac{-1}{p} \right)$ . In this case, the degree of  $(0, 0, 0)$  is  $\Theta(p^2)$ . The size  $n$  of the vertex set  $\mathcal{V}$  is  $p^3$ , so the number of edges is  $\Theta(p^5)$  which is  $\Theta(n^{5/3})$  and thus, the bound given by the **Kővári-Sós-Turán Theorem** is (asymptotically) tight.

For  $s \leq t$ , the **Kővári-Sós-Turán Theorem** gives

$$\text{ex}(n; K_{s,t}) < C_{s,t} n^{2-1/s}$$

for some  $c_{s,t}$  depending only on  $s$  and  $t$ . However, it is not known if the above bound is tight for most cases. Lower bounds of the form

$$\text{ex}(n; K_{s,t}) \geq c_{s,t} n^{2-1/t}$$

are known only if  $t > (s-1)!$ . See [KRS96] for more details.

#### 1.4. $P_{k+1}$ -free graphs

Next, we study  $\text{ex}(n; P_{k+1})$ .

First of all, note that  $K_k$  is  $P_{k+1}$ -free. So, if we split the  $n$  vertices up into blocks of  $k$  vertices and add all edges within each block, the resulting graph will be  $P_{k+1}$ -free as well. That is,  $G_n$  is a disjoint union of  $K_k$ s (and possibly one  $K_r$  for  $r < k$ ). For this particular graph  $G_n$ ,

$$e(G_n) \leq \frac{(k-1)n}{2}.$$

It turns out that we cannot do better than this.

**Theorem 1.7.**  $\text{ex}(n; P_{k+1}) \leq (k-1)n/2$  with equality if and only if  $G_n$  is a disjoint union of  $K_k$ s.

The above theorem just says that equality is not attained for connected graphs without  $P_{k+1}$ . To prove this, we prove the following (seemingly) more general statement.

**Lemma 1.8.** Let  $G_n = (V, E)$  be connected and suppose  $d(v) \geq k$  for all  $v \in V$ . If  $n \geq 2k$ ,  $G_n$  contains a path of length  $2k$ . Otherwise,  $G_n$  contains an  $n$ -cycle.

*Proof.* We prove the result for the case where  $n \geq 2k$  first. Consider the longest path  $u = v_1, v_2, \dots, v_r = v$  in  $G_n$  and let  $U = \{v_i : i \in [r]\}$ . We must show that  $r \geq 2k$ . Suppose otherwise and let  $r < 2k$ .

First of all, we must have  $\Gamma(u) \subseteq U$  – otherwise, the path can be extended by adding another vertex from  $\Gamma(u) \setminus U$ . Similarly,  $\Gamma(v) \subseteq U$ .

Next,  $v_1$  and  $v_r$  cannot be adjacent. If they are, then we can obtain a longer path by cycling through and choosing some edge from a vertex in  $U$  to one outside of  $U$  (such a vertex must exist since the graph is connected and has at least  $2k$  vertices). More generally, if there exists  $i$  such that  $v_1, v_{i+1}$  and  $v_i, v_r$  are adjacent, then we arrive at a contradiction (for the same reason).

Let  $S = \{i : v_i v_r \in E\}$  and  $T = \{i : v_1 v_{i+1} \in E\}$ . By the above observation,  $S \cap T = \emptyset$ . However, by our first observation,  $|S| = d(v_r) \geq k$  and  $|T| = d(v_1) \geq k$ . Therefore,  $r \geq |S \cup T| \geq 2k$ .

The result for  $n < 2k$  is shown using nearly the same proof. ■

*Proof of Theorem 1.7.* We perform strong induction on  $n$ . We may assume that  $n > k$ . Suppose  $G_n$  has no  $P_{k+1}$ . If  $G_n$  is not connected, then it consists of a disjoint union of connected subgraphs. We may then apply the inductive hypothesis to each of these smaller pieces.

So, let  $G_n$  be connected. By Lemma 1.8, there is some vertex  $v$  such that  $d(v) \leq (k-1)/2$  (otherwise, there must be a path of length  $k$ ). Additionally, observe that  $G_n$  does not have any subgraph isomorphic to  $K_k$  – using the connectedness assumption gives a path of length  $k$  otherwise.

The graph  $G \setminus \{x\}$ <sup>4</sup> has no  $P_{k+1}$  and further, it has no  $K_k$  either. Therefore,  $G \setminus \{x\}$  is not extremal and  $e(G \setminus \{x\}) < (k-1)(n-1)/2$ . Therefore,

$$\begin{aligned} e(G) &= d(x) + e(G \setminus \{x\}) \\ &< \frac{k-1}{2} + \frac{(k-1)(n-1)}{2} = \frac{(k-1)n}{2}. \end{aligned}$$

■

<sup>4</sup>This is the subgraph induced by  $G$  on the vertex set  $V \setminus \{x\}$

## §2. Fundamental Results in Extremal Graph Theory

### 2.1. The Erdős-Stone-Simonovits Theorem

#### 2.1.1. Motivation

Using **the folklore result** that a graph is bipartite iff it has no odd cycle, we have

$$\text{ex}(n; C_{2k+1}) \geq \left\lfloor \frac{n^2}{4} \right\rfloor.$$

*Remark.* It in fact turns out that for  $n \gg 0$ ,  $\text{ex}(n; C_{2k+1}) = \lfloor n^2/4 \rfloor$ , as we shall see later..

Is there any more general relationship between forbidden subgraphs and  $r$ -partite graphs?

**Definition 2.1** (Chromatic Number). Given a graph  $G$ , its **chromatic number** is given by

$$\chi(G) = \min\{r : G \text{ is } r\text{-partite}\}.$$

Alternatively, we can define the above using the following.

**Definition 2.2.** Given a graph  $G = (V, E)$ , an  $r$ -**coloring** of  $G$  is a function  $f : V \rightarrow [r]$  such that for any  $uv \in E$ ,  $f(u) \neq f(v)$ .

The chromatic number of a graph is the least  $r$  such that it is  $r$ -colorable.

Suppose  $H$  is an arbitrary graph such that  $\chi(H) = r + 1$ . Then, no  $r$ -partite graph contains  $H$ . As a result,

$$\text{ex}(n; H) > t_r(n).$$

Our earlier observation on  $\text{ex}(n; C_{2k+1})$  is then just a consequence of the fact that  $C_{2k+1}$  is 3-colorable.

Let us give a more concrete example. Suppose we want to find  $\text{ex}(n; \text{Petersen})$ , where Petersen is the **Petersen graph**. It may be checked that Petersen has chromatic number 3. So,

$$\text{ex}(n; \text{Petersen}) > t_2(n) = \left\lfloor \frac{n^2}{4} \right\rfloor.$$

Can we do better?

This is answered by the Erdős-Stone-Simonovits Theorem.

#### 2.1.2. The result

**Theorem 2.1** (Erdős-Stone-Simonovits Theorem, Version 1). Let  $r \geq 1$  and  $0 < \varepsilon < 1/2$ . Then, for  $n \gg 0$  and any graph  $G_n$ , if

$$\delta(G_n) \geq \left(1 - \frac{1}{r} + \varepsilon\right)n, \tag{2.1}$$

there exist pairwise disjoint subsets  $V_1, \dots, V_{r+1}$  of  $V$  such that for each  $i$ ,

$$|V_i| = t \geq \frac{\varepsilon \log n}{2^{r-1}(r-1)!}$$

and the complete  $(r+1)$ -partite graph on these subsets is contained in  $G_n$ .

**Theorem 2.2** (Erdős-Stone-Simonovits Theorem, Version 2). Let  $r \geq 1$  and  $0 < \varepsilon < 1/2$ . Then, for  $n \gg 0$  and any graph  $G_n$ , if

$$e(G_n) \geq \left(1 - \frac{1}{r} + \varepsilon\right) \binom{n}{2}, \quad (2.2)$$

there exist pairwise disjoint subsets  $V_1, \dots, V_{r+1}$  of  $V$  such that for each  $i$ ,

$$|V_i| = t \geq \frac{\varepsilon \log n}{2^{r+1}(r-1)!}$$

and the complete  $(r+1)$ -partite graph on these subsets is contained in  $G_n$ .

Observe the difference in the exponent of 2 in the denominator of  $t$  in the two versions.

For example, if  $n \geq e^{32/\varepsilon}$  and Equation (2.2) is satisfied, then  $T_3(12)$ , and thus Petersen, is a subgraph of  $G_n$ . Therefore, (replacing  $\varepsilon$  with  $2\varepsilon$ )

$$\text{ex}(n; \text{Petersen}) \leq \left(\frac{1}{2} + 2\varepsilon\right) \binom{n}{2} \leq \left(\frac{1}{4} + \varepsilon\right) n^2.$$

for all  $\varepsilon > 0$ .

**Corollary 2.3.** If  $\chi(H) = r + 1$ , then for any  $\varepsilon > 0$ , for  $n \gg 0$ ,

$$\left(1 - \frac{1}{r}\right) \frac{n^2}{2} \leq \text{ex}(n; H) \leq \left(1 - \frac{1}{r} + \varepsilon\right) n^2 - \mathcal{O}(n).$$

This implies that if  $\chi(H) = r + 1$ , then for  $n \gg 0$

$$\left(1 - \frac{1}{r}\right) \frac{n^2}{2} \leq \text{ex}(n; H) \leq \left(1 - \frac{1}{r} + o(1)\right) \frac{n^2}{2}$$

*Proof of Erdős-Stone-Simonovits Theorem, Version 1.* We first show the result for  $r = 1$ . Let  $G_n$  be a graph such that  $\delta(G_n) \geq \varepsilon n$ . We want a “large”  $t$  such that  $K_{t,t}$  is isomorphic to a subgraph  $G_n$ .

Suppose that  $G_n$  is  $K_{t,t}$ -free (we shall fix  $t$  later). Now, we have that

$$t \binom{n}{t} \stackrel{(1)}{>} |\{(v, T) : v \in V, T \subseteq \Gamma(v), |T| = t\}| = \sum_{v \in V} \binom{d(v)}{t} \stackrel{(2)}{\geq} n \binom{\varepsilon n}{t},$$

where (2) follows from the hypothesis on  $\delta(G_n)$  and (1) follows due to our assumption that  $G_n$  is  $K_{t,t}$ -free. Therefore,

$$n \binom{\varepsilon n}{t} < t \binom{n}{t}.$$

So,

$$\begin{aligned} 1 &< \frac{t}{n} \cdot \frac{n(n-1) \cdot (n-t+1)}{\varepsilon n(\varepsilon n-1) \cdot (\varepsilon n-t+1)} \\ &< \frac{t}{n} \cdot \left(\frac{n}{\varepsilon n-t+1}\right)^t \\ &= \frac{t}{n} \frac{1}{\varepsilon^t} \left(\frac{1}{1 - \frac{t-1}{\varepsilon n}}\right)^t. \end{aligned}$$

We desire a large  $t$  such that the above results in a contradiction. Let us look at the last term. Suppose we want

$$\left(1 - \frac{t-1}{\varepsilon n}\right)^t \rightarrow 1.$$

Equivalently,

$$t \log \left(1 - \frac{t-1}{\varepsilon n}\right) \rightarrow 0$$

$$t \left( \frac{t-1}{\varepsilon n} + \frac{1}{2} \left(\frac{t-1}{\varepsilon n}\right)^2 + \frac{1}{3} \left(\frac{t-1}{\varepsilon n}\right)^3 + \dots \right) \rightarrow 0.$$

If  $\varepsilon n > t^4$ , the above holds (Why?).

That is, if  $t < (\varepsilon n)^{1/4}$ , then for sufficiently large  $n, t$ ,

$$\left(1 - \frac{t-1}{\varepsilon n}\right)^{-t} < 2$$

and

$$1 < \frac{t}{n} \cdot \frac{1}{\varepsilon^t} \left(1 - \frac{t-1}{\varepsilon n}\right)^{-t} < \frac{2t}{\varepsilon^t n}.$$

If the above is  $\leq 1$ , we shall arrive at a contradiction. That is, if

$$\log(2t) + t \log \left(\frac{1}{\varepsilon}\right) < \log n,$$

then we arrive at a contradiction. In particular,

$$t = \lceil \varepsilon \log n \rceil$$

is a suitable choice (Why?), completing the proof of the theorem for  $r = 1$ .

Now, let us prove the general case by performing induction on  $r$ . Let  $G_n$  be a graph with

$$\delta(G_n) \geq \left(1 - \frac{1}{r} + \varepsilon\right) n.$$

Now,

$$1 - \frac{1}{r} + \varepsilon > 1 - \frac{1}{r-1} + \frac{1}{r(r-1)}.$$

Let  $\varepsilon' = 1/r(r-1)$ . By induction, there are  $V'_1, \dots, V'_r$  such that

$$|V'_i| \geq t' = \frac{\log n}{r(r-1)2^{r-2}(r-2)!} = \frac{\log n}{2^{r-2}r!} \quad (2.3)$$

for each  $i$  and the complete  $r$ -partite graph on these sets is a subgraph of  $G_n$ . Let  $K = \bigcup V'_i$ .

We obviously have  $\varepsilon < 1/r$ , since the claim is vacuously true otherwise.

We shall find  $V_i \subseteq V'_i$  for each  $1 \leq i \leq r$  and some  $V_{r+1} \subseteq V \setminus K$  such that the complete bipartite graph on  $(V_1, \dots, V_{r+1})$  is the required subgraph of  $G_n$ .

Now, define

$$U = \left\{ x \in V \setminus K : d(x, K) \geq \left(1 - \frac{1}{r} + \lambda\right) |K| \right\}$$

for some  $\lambda$  we shall fix later. We shall bound  $e(K, V \setminus K)$  in two different ways. From the perspective of  $K$ ,

$$e(K, V \setminus K) \geq |K| \left( \left(1 - \frac{1}{r} + \varepsilon\right) n - |K| \right).$$



From the perspective of  $V \setminus K$ ,

$$\begin{aligned} e(K, V \setminus K) &= e(V \setminus (K \sqcup U), K) + e(U, K) \\ &\leq (n - |U| - |K|) \left(1 - \frac{1}{r} + \lambda\right) |K| + |U||K|. \end{aligned}$$

Putting the two together,

$$\begin{aligned} |K| \left( \left(1 - \frac{1}{r} + \varepsilon\right) n - |K| \right) &\leq (n - |U| - |K|) \left(1 - \frac{1}{r} + \lambda\right) |K| + |U||K| \\ \left(1 - \frac{1}{r} + \varepsilon\right) n - |K| &\leq (n - |U| - |K|) \left(1 - \frac{1}{r} + \lambda\right) + |U|. \end{aligned}$$

Set  $\lambda = \varepsilon/2$ . We then have

$$\frac{r}{2}\varepsilon n \leq \left(1 - \frac{r\varepsilon}{2}\right) (|U| + |K|).$$

So,

$$|U| \geq \frac{r\varepsilon}{2 - r\varepsilon} n - |K| \stackrel{(*)}{\geq} \frac{\varepsilon}{1 - \varepsilon} n - rt',$$

where  $(*)$  follows from the fact that  $r\varepsilon/(2 - r\varepsilon)$  is increasing in  $r$  and  $r \geq 2$ .

Since  $t'$ , and thus  $rt'$ , is of the order of  $\log n$ , the first term in the expression dominates for sufficiently large  $n$ . So, for  $n \gg 0$ ,  $|U| \geq \varepsilon n$ .

Now,

$$\begin{aligned} \left(1 - \frac{1}{r} + \frac{\varepsilon}{2}\right) |K| &\geq \left(1 - \frac{1}{r} + \frac{\varepsilon}{2}\right) rt' \\ &\geq (r - 1)t' + \frac{\varepsilon r}{2} t'. \end{aligned}$$

This implies that each  $u \in U$  has at least  $(\varepsilon r/2)t'$  neighbours in each  $V'_i$ .

We can now use a pigeonhole argument to choose a subset of  $U$  whose vertices are all adjacent to some common set of vertices in each  $V'_i$ . To do so, consider

$$|\{(u, W_1, \dots, W_r) : W_i \subseteq V'_i, |W_i| = (\varepsilon r/2)t', \text{ and } u \text{ is adjacent to all the vertices of each } W_i\}|.$$

By our earlier observation, this must be at least  $|U| \geq \varepsilon n$ .

On the other hand, it is at most the number of ways of choosing the  $W_i$ , which is  $\binom{t'}{(\varepsilon r/2)t'}^r$ .

In particular, using a pigeonhole argument, there exist  $V_1, \dots, V_r$  and a  $V_{r+1} \subseteq U$  such that  $V_i \subseteq V'_i$  for each  $1 \leq i \leq r$  and for all  $u \in V_{r+1}$ ,  $(u, V_1, \dots, V_r)$  is in the set whose cardinality we just considered, and  $|V_{r+1}| \geq \varepsilon n / \binom{t'}{(\varepsilon r/2)t'}^r$ . Let us now bound this expression.

$$\begin{aligned} |V_{r+1}| &\geq \frac{\varepsilon n}{\left(\frac{t'}{(\varepsilon r/2)t'}\right)^r} \\ &\geq \frac{\varepsilon n}{(2e/\varepsilon r)^{t'\varepsilon r^2/2}} && \text{(see [here](#) for the bound used)} \\ &\geq \varepsilon n \cdot \left(\frac{\varepsilon}{e}\right)^{t'\varepsilon r^2/2} && \text{(since } r \geq 2, 2/r \leq 1). \end{aligned}$$

Setting

$$t = \frac{\varepsilon \log n}{2^{r-1}(r-1)!},$$

we see that  $t = \varepsilon r t' / 2$ .

Keeping in mind that the bound we want is  $|V_{r+1}| \geq t$ ,

$$\begin{aligned}
 \log \left( n \left( \frac{\varepsilon}{e} \right)^{t' \varepsilon r^2 / 2} \right) &\geq \log n + \frac{\varepsilon r^2 t'}{2} \log \left( \frac{\varepsilon}{e} \right) \\
 &\geq t' \left( 2^{r-2} r! - \log \left( \frac{e}{\varepsilon} \right) \cdot \frac{\varepsilon r^2}{2} \right) && \text{(using the expression for } t' \text{ in (2.3))} \\
 &\geq t' \left( 2^{r-2} r! - \log(2e) \cdot \frac{r^2}{4} \right) && (-\varepsilon \log(e/\varepsilon) \text{ is decreasing in } \varepsilon) \\
 &\geq \log(r t' / 2).
 \end{aligned}$$

Therefore,  $V_{r+1} \geq t$ . Since  $|V_i| = \varepsilon r t' / 2 = t$  by definition, the proof is complete. ■

Whew. Let us now give a simple corollary of the above result.

**Porism 2.4.** Suppose  $H_1, \dots, H_m$  are graphs. Then, for any  $\varepsilon > 0$  and  $n \gg 0$ ,

$$\text{ex}(n; H_1, \dots, H_m) \leq \left( 1 - \frac{1}{r} + \varepsilon \right) \frac{n^2}{2} - \mathcal{O}(n^2),$$

where  $r + 1 = \max\{\chi(H_1), \dots, \chi(H_m)\}$ .

Before we move onto the proof of the second version of the Erdős-Stone-Simonovits Theorem, we give a lemma that will assist in its proof.

**Lemma 2.5.** For  $n \gg 0$ , if  $e(G_n) \geq (c + \varepsilon) \binom{n}{2}$  for some  $c > 0$  and  $\varepsilon > 0$ , then there exists  $H \subseteq G_n$  such that

1.  $|H| \geq \sqrt{\varepsilon} n$ .
2.  $\delta(H) \geq c|H|$ .

*Proof.* Assume that  $G_n$  itself does not satisfy the conclusions. Then, there exists some  $x_n \in V(G_n)$  such that  $d(x_n) < cn$ .

Let  $H_{n-1} = G \setminus \{x_n\}$ . If  $H_{n-1}$  fails the second conclusion of the theorem, there exists  $x_{n-1} \in V(H_{n-1})$  such that  $d(x_{n-1}) < c(n-1)$ .

Repeating the above, we get a sequence of graphs  $G_n = H_n \supsetneq H_{n-1} \supsetneq \dots \supsetneq H_\ell$ , where  $V(H_{n-r}) \setminus V(H_{n-r-1}) = \{x_{n-r}\}$  for each  $r$  and  $\ell \geq \sqrt{\varepsilon} n$ . Further, for each  $i$ ,  $d_{H_i}(x_i) < ci$  for  $i = \ell, \dots, n$ .

Now,

$$\begin{aligned}
 e(H_\ell) &> e(H_n) - (cn + c(n-1) + \dots + c(\ell+1)) \\
 &= (c + \varepsilon) \binom{n}{2} - c \left( \frac{n(n+1)}{2} - \frac{\ell(\ell+1)}{2} \right) \\
 &= (c + \varepsilon) \binom{n}{2} - c \left( \binom{n}{2} + n - \binom{\ell+1}{2} \right) \\
 &= \varepsilon \binom{n}{2} - cn + c \binom{\ell+1}{2}.
 \end{aligned}$$

Now, the final expression must be at most  $\binom{\ell}{2}$  (since  $H_\ell$  has  $\ell$  vertices).

As a result, it would suffice to show that the above expression on taking  $\ell = \lfloor \varepsilon n \rfloor$  is greater than  $\binom{\ell}{2}$  for sufficiently large  $n$  (this implies that the sequence must stop before reaching this  $\ell$  due to one of the graphs satisfying the conclusions). Indeed, this is seen to be true as

$$\begin{aligned} \varepsilon \binom{n}{2} - cn + c \binom{\ell+1}{2} &\geq \varepsilon \binom{n}{2} + c \left( \frac{(\lfloor \sqrt{\varepsilon n} \rfloor + 1) \lfloor \sqrt{\varepsilon n} \rfloor}{2} - n \right) \\ &\geq \varepsilon \binom{n}{2} \\ &\geq \binom{\ell}{2}, \end{aligned} \quad (\text{for sufficiently large } n)$$

completing the proof. ■

## 2.2. An Introduction to Random Graphs

**Definition 2.3** (Erdős-Rényi Model). Fix  $0 \leq p \leq 1$ . The **Erdős-Rényi random graph model**, denoted  $G_{n,p}$  is the random variable which is a graph with vertex set  $[n]$ , such that for each  $\{i, j\} \in \binom{[n]}{2}$ ,  $\{i, j\}$  is an edge with probability  $p$ , independently across distinct pairs.

So, for any graph  $H$  on vertex set  $[n]$ ,

$$\Pr(G_{n,p} = H) = p^{e(H)} (1-p)^{\binom{n}{2} - e(H)}.$$

*Remark.*  $G_{n,1/2}$  is the uniform distribution on the set of graphs on  $[n]$ .

A recurring theme in probability theory is that random objects tend to behave very nicely given a large number of samples (along the lines of the laws of large numbers and the central limit theorem).

### 2.2.1. A motivating extremal problem (bounding $\text{ex}(n; C_{2k})$ )

To understand why random graphs are important, let us look at  $\text{ex}(n; C_{2k})$ .

**Theorem 2.6.** For any  $k$ , there exists a constant  $c$  such that for  $n \gg 0$ , there is a  $C_k$ -free  $G_n$  with

$$e(G_n) \geq c \cdot n^{1+1/(k-1)}.$$

The above result does not yield anything useful for  $k$  odd.

*Proof.* Consider  $G_{n,p}$  for some  $p$  we shall fix later. Let  $N(G)$  be the number of copies of  $C_k$  in a given graph  $G$ . Given a cycle  $(v_1, \dots, v_k)$ , observe that the sequences  $(v_2, v_3, \dots, v_k, v_1)$  and  $(v_k, v_{k-1}, \dots, v_1)$  determine the same cycle. That is, performing cyclic shifts of a sequence of vertices or reversing their order around gives the same cycle. Let  $\mathcal{C}$  be the set of all these cycles.<sup>5</sup> By our observation,

$$|\mathcal{C}| = \frac{n(n-1) \cdots (n-k+1)}{2k} = \frac{n!}{2k \cdot k!}.$$

<sup>5</sup>This can be made more formal by taking all length  $k$  sequences of  $[n]$  consisting of distinct elements and considering the equivalence classes formed by the equivalence relation defined on the previous line.

Now,

$$\begin{aligned}
 \mathbb{E}[N(G_{n,p})] &= \mathbb{E} \left[ \sum_{(v_1, \dots, v_k) \in \mathcal{C}} \mathbb{1}_{v_1 v_2, v_2 v_3, \dots, v_{k-1} v_k, v_k v_1 \text{ are edges}} \right] \\
 &= \sum_{(v_1, \dots, v_k) \in \mathcal{C}} \Pr \left[ \mathbb{1}_{v_1 v_2, v_2 v_3, \dots, v_{k-1} v_k, v_k v_1 \text{ are edges}} \right] && \text{(linearity of expectation)} \\
 &= \sum_{(v_1, \dots, v_k) \in \mathcal{C}} p^k \\
 &= \frac{n!}{2k \cdot k!} p^k.
 \end{aligned}$$

This quantity is obviously less than  $(np)^k/2k$ . If  $p$ , and thus the expectation is small, we expect to not see many  $C_k$ s. On the other hand,

$$\begin{aligned}
 \mathbb{E}[e(G_{n,p})] &= \mathbb{E} \left[ \sum_{\{i,j\} \in \binom{[n]}{2}} \mathbb{1}_{ij \text{ is an edge}} \right] \\
 &= \binom{n}{2} p.
 \end{aligned}$$

Given a graph, if we delete an (arbitrary) edge from each copy of  $C_k$  in it, we will be left with no cycles. That is, given any graph  $G$ , there is a graph on the same vertex set with  $e(G) - N(G)$  edges that is  $C_k$ -free. Inspired by this, by the linearity of expectation,

$$\mathbb{E}[e(G_{n,p}) - N(G_{n,p})] \geq \binom{n}{2} p - \frac{(np)^k}{2k}.$$

If we set  $p = \left(\frac{k}{2}\right)^{1/(k-1)} n^{-1+1/(k-1)}$ , then the above quantity is at least  $n(n-1)p/4 \geq c \cdot n^{1+1/(k-1)}$  for an appropriate constant  $c$  and  $n \gg 0$ , completing the proof. ■

### 2.2.2. Digression: A coloring result of Erdős

The question we consider in this section is:

Are there  $C_3$ -free graphs with large chromatic number?

More generally,

Are there graphs with large chromatic number and large girth?

For example, if the girth of a graph is 7, then there cannot be adjacent vertices  $v, w$  such that  $\Gamma(v) \cap \Gamma(u)$  and  $\Gamma(w) \cap \Gamma(u)$  are non-empty for some  $u$  distinct from  $v, w$ . As a result, we can draw a “2-step tree” rooted at any  $u$ , which has  $\Gamma(u)$  at the first level and the neighbours (other than  $u$ ) of vertices of  $\Gamma(u)$  at the second.

This seems to suggest some level of sparseness in the graph, due to which there are not too many edges and as a result, the chromatic number is low. However, it turns out that this intuition is not true, as proved by Erdős in [Erd59].

**Theorem 2.7.** There exist graphs with arbitrarily large girth and chromatic number. That is, given a  $g, k \geq 3$ , there exists a graph  $G_n$  such that  $\text{girth}(G) > g$  and  $\chi(G) > k$ .

*Proof.* Consider  $G_{n,p}$  for some  $p$  we fix later. Further assume that  $np \geq 1$ .

Given a graph  $G$ , let  $N_i(G)$  (for  $3 \leq i \leq g$ ) be the number of cycles of size  $i$  in  $G$ . As we saw in the proof of Theorem 2.6,

$$\mathbf{E}[N_i(G_{n,p})] = \frac{n!}{2i \cdot i!} p^i < \frac{(np)^i}{6}.$$

Let  $N = \sum_{i=3}^g N_i$ . Then

$$\begin{aligned} \mathbf{E}[N(G_{n,p})] &= \mathbf{E}\left[\sum_{i=3}^g N_i(G_{n,p})\right] \\ &< \frac{(np)^3}{6} \left(\frac{(np)^{g-2} - 1}{np - 1}\right) \\ &< \frac{(np)^g}{3}. \end{aligned} \quad (\text{since } np \geq 1)$$

Using Markov's inequality,

$$\Pr\left[N(G_{n,p}) > \frac{2}{3}(np)^g\right] < \frac{1}{2} \quad (2.4)$$

This takes care of the girth (we want the above probability to be small). On the other hand, we need to make the chromatic number large. Towards this, observe that  $\chi(G) \geq n/\alpha(G)$  (Why?). We have

$$\begin{aligned} \Pr[\alpha(G_{n,p}) \geq r] &= \Pr\left[\bigcup_{X \subseteq [n]: |X|=r} \{X \text{ is independent}\}\right] \\ &\leq \sum_{X \subseteq [n]: |X|=r} \Pr[X \text{ is independent}] \\ &= \sum_{X \subseteq [n]: |X|=r} (1-p)^{\binom{r}{2}} \\ &\leq \binom{n}{r} \cdot e^{-pr(r-1)/2} \\ &\leq \left(\frac{en}{r}\right)^r \cdot e^{-pr^2/3} \\ &= \left(\frac{e^{1-pr/3}n}{r}\right)^r. \end{aligned}$$

We shall choose  $r$  and  $p$  such that with positive probability,  $N(G_{n,p}) \leq 2(np)^g/3$  and  $\alpha(G_{n,p}) < r$ . This implies the existence of a graph  $G_n$  such that both of the above hold.

We cannot use the tactic of removing edges we did in the earlier proof since that might increase  $\alpha$ . Deleting vertices on the other hand works, since this can increase neither  $\alpha$  nor  $N$ .

If we delete a single vertex from each cycle involved in  $N$ , the resulting graph will have girth greater than  $g$ . That is, given a graph  $G_n$ , there exists a graph with at least  $n - N(G_n)$  vertices that has girth greater than  $g$ . Denote this corresponding graph as  $G'_n$ .

Set  $p = n^{1/(g+1)-1}$ . In this case,  $2(np)^g/3 < n/2$  for  $n \gg 0$  and using (2.4),

$$\Pr\left[n - N(G_{n,p}) > \frac{n}{2}\right] \geq \frac{1}{2}. \quad (2.5)$$

Set  $r = 4 \log n/p = 4n^{1-1/(g+1)} \log n$ . For these values of  $p$  and  $r$ ,

$$\Pr[\alpha(G_{n,p}) \geq r] \xrightarrow{n \rightarrow \infty} 0. \quad (2.6)$$

Using (2.5), our construction of  $G'$ , and (2.7), it is true with positive probability that

$$|G'_{n,p}| \geq \frac{n}{2}, \text{girth}(G'_{n,p}) > g,$$

and

$$\chi(G'_{n,p}) \geq \frac{n}{\alpha(G'_{n,p})} \geq \frac{n}{r} = \frac{n^{1/(g+1)}}{4 \log n} \xrightarrow{n \rightarrow \infty} \infty.$$

Therefore, for  $g, k \geq 3$ , there exists  $n \gg 0$  and graph  $G$  on  $n$  vertices such that  $\text{girth}(G) > g$  and  $\chi(G) > k$ . ■

### 2.3. Szemerédi's Regularity Lemma

The second of our powerful results in extremal graph theory (after the Erdős-Stone-Simonovits Theorem) is Szemerédi's Regularity Lemma, which says that any sufficiently large graph behaves in some way like a random graph.

#### 2.3.1. Motivation

First, let us give a bound from probability theory that will be useful.

**Lemma 2.8** (Chernoff Bound). Suppose  $X \sim B(n, p)$ , the **binomial distribution** with parameters  $n, p$ . Then for any  $t \geq 0$ ,

$$\Pr [X - \mathbf{E}[X] \geq t] \leq \exp \left( -\frac{t^2}{2(\mathbf{E}[X] + t/3)} \right)$$

and

$$\Pr [X - \mathbf{E}[X] \leq -t] \leq \exp \left( -\frac{t^2}{2\mathbf{E}[X]} \right).$$

Fix disjoint  $A, B \subseteq [n]$  and let  $|A| = a, |B| = b$ . Then given a graph  $G$  on  $[n]$ ,

$$e(A, B) = \sum_{x \in A, y \in B} \mathbb{1}_{xy \in E(G)}.$$

Fix  $0 < p < 1$ . Then if  $G \sim G_{n,p}$ ,

$$e(A, B) \sim B(ab, p).$$

By the **Chernoff Bound**, for some fixed constant  $c$ ,

$$\Pr \left[ |e(A, B) - pab| > c \left( b \sqrt{pa \log \left( \frac{2n}{b} \right)} \right) \right] \xrightarrow{n \rightarrow \infty} 0.$$

In particular, if  $a = b = \alpha n$  for some  $0 < \alpha < 1/3$ , then with high probability,

$$|e(A, B) - pab| = \mathcal{O} \left( b \sqrt{ap \log \left( \frac{2n}{b} \right)} \right)$$

for any sets  $A, B$  of sizes  $a$  and  $b$  respectively.

This seems to say that the actual number of edges between two sets of the given size does not deviate very much from the expected number of edges between the two sets. The expression on the right is of the order of  $\mathcal{O}(n\sqrt{n})$ , which is asymptotically less than the expectation  $pab = \mathcal{O}(n^2)$ .

## 2.3.2. The result

The regularity lemma gives a qualitative version of the above observation. Before we move to the actual result, let us provide some notation.

**Definition 2.4** (Density). Given a graph  $G = (V, E)$  and  $U, W \subseteq V$ , the **density**  $d(U, W)$  is equal to  $e(U, W)/|U||W|$ .

*Remark.* Here,  $e(U, W)$  is  $\{(u, w) \in U \times W : \{u, w\} \in E\}$ . If  $U$  and  $W$  are disjoint, this is the same as our earlier definition of  $e(\cdot, \cdot)$ . If they are not disjoint however, edges within the intersection are counted *twice* in our current definition.

This does not matter all that much since we usually apply the regularity lemma on disjoint sets.

**Definition 2.5** ( $\varepsilon$ -regular pair). Suppose  $0 < \varepsilon < 1$ . A pair of subsets  $(U, W)$  is said to be  **$\varepsilon$ -regular** if for any  $A \subseteq U$ ,  $B \subseteq W$  with  $|A| \geq \varepsilon|U|$  and  $|B| \geq \varepsilon|W|$ , we have

$$|d(A, B) - d(U, W)| \leq \varepsilon$$

This corresponds to some sort of uniform behaviour throughout the sets, where subsets behave similarly to their parent sets in terms of density. If  $(U, W)$  is  $\varepsilon$ -regular, all sufficiently large subsets of  $U, W$  have roughly the same edge density as  $(U, W)$ .

**Theorem 2.9** (Szemerédi's Regularity Lemma). Given  $0 < \varepsilon < 1$ , there exists  $M$  such that for  $n \gg 0$ , any graph  $G_n$  admits a vertex partition  $\mathcal{P} = (V_0, V_1, \dots, V_k)$ , where

- $k \leq M$ ,
- $|V_0| \leq \varepsilon n$  ( $V_0$  is known as an “exceptional set”),
- all the  $V_i$  for  $1 \leq i \leq k$  are of equal size, and
- the number of  $\varepsilon$ -irregular pairs  $(V_i, V_j)$  ( $1 \leq i, j \leq k$ ) is at most  $\varepsilon k^2$ .

Such a partition where the number of  $\varepsilon$ -regular pairs is at most  $\varepsilon k^2$  is often referred to as an  **$\varepsilon$ -regular partition**.

We present the proof of the above, that uses an “energy increment” argument, over a series of lemmas.

**Definition 2.6** (Energy). Given a graph  $G_n$  with vertex set  $V$ , for disjoint  $U, W \subseteq V$ , define the **energy** of the pair  $(U, W)$  by

$$q(U, W) = \frac{|U||W|}{n^2} d^2(U, W).$$

If  $\mathcal{U} = \{U_1, \dots, U_m\}$  and  $\mathcal{W} = \{W_1, \dots, W_\ell\}$  are partitions of  $U$  and  $W$  respectively, then the energy of the pair  $(\mathcal{U}, \mathcal{W})$  is

$$q(\mathcal{U}, \mathcal{W}) = \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq \ell}} q(U_i, W_j).$$

For a partition  $\mathcal{P}$  of  $V$ , let

$$q(\mathcal{P}) = q(\mathcal{P}, \mathcal{P}) = \sum_{U, W \in \mathcal{P}} q(U, W).$$

**Lemma 2.10.** If  $U, W$  are disjoint subsets of  $V$  and  $\mathcal{U}, \mathcal{W}$  are partitions of  $U, W$  respectively, then

$$q(\mathcal{U}, \mathcal{W}) \geq q(U, W).$$

*Proof.* Independently pick  $u, w$  uniformly randomly from  $V$ . Define the random variable

$$Z(u, w) = \begin{cases} d(U', W'), & u \in U' \in \mathcal{U}, w \in W' \in \mathcal{W}, \\ 0, & \text{otherwise.} \end{cases}$$

Then,

$$\begin{aligned} \mathbf{E}[Z]^2 &= \left( \sum_{\substack{U' \in \mathcal{U} \\ W' \in \mathcal{W}}} d(U', W') \left( \frac{|U'| |W'|}{|U| |W|} \right) \right)^2 \\ &= \left( \frac{1}{|U| |W|} \sum_{\substack{U' \in \mathcal{U} \\ W' \in \mathcal{W}}} e(U', W') \right)^2 \\ &= d(U, W)^2 \\ &= \frac{n^2}{|U| |W|} q(U, W) \text{ and} \\ \mathbf{E}[Z^2] &= \sum_{\substack{U' \in \mathcal{U} \\ W' \in \mathcal{W}}} d^2(U', W') \left( \frac{|U'| |W'|}{|U| |W|} \right) \\ &= \frac{n^2}{|U| |W|} q(\mathcal{U}, \mathcal{W}). \end{aligned}$$

Since variance is always non-negative,  $\mathbf{E}[Z^2] \geq \mathbf{E}[Z]^2$  and thus,

$$q(\mathcal{U}, \mathcal{W}) \geq q(U, W). \quad \blacksquare$$

Partitioning increases energy! Also observe that since the random variable  $Z$  is at most 1 (1/2, in fact), so is the energy.

This also implies that if the partition  $\mathcal{Q}$  is a refinement of  $\mathcal{P}$ , then

$$q(\mathcal{Q}) \geq q(\mathcal{P}).$$

**Lemma 2.11.** If  $(U, W)$  is not  $\varepsilon$ -regular, there is a partition  $\mathcal{U} = \{U_1, U_2\}$  and  $\mathcal{W} = \{W_1, W_2\}$  of  $U, W$  respectively such that

$$q(\mathcal{U}, \mathcal{W}) > q(U, W) + \varepsilon^4 \frac{|U| |W|}{n^2}.$$

*Proof.* By the  $\varepsilon$ -irregularity, there exist  $U_1 \subseteq U$  and  $W_1 \subseteq W$  with  $|U_1| \geq \varepsilon |U|$ ,  $|W_1| \geq \varepsilon |W|$ , and

$$|d(U_1, W_1) - d(U, W)| > \varepsilon.$$



Consider the partitions  $\mathcal{U} = \{U_1, U \setminus U_1\}$  and  $\mathcal{W} = \{W_1, W \setminus W_1\}$ . With the same  $Z$  as in the proof of the previous lemma,

$$\begin{aligned} \mathbf{E}[Z^2] - \mathbf{E}[Z]^2 &= \mathbf{E}[(Z - \mathbf{E}[Z])^2] \\ &\geq \mathbf{E}[(Z - \mathbf{E}[Z])^2 \mathbb{1}_{u \in U', w \in W'}] \\ &= \varepsilon^4. \end{aligned}$$

The claim follows. ■

**Lemma 2.12.** Suppose  $0 < \varepsilon < 1/4$  and let  $\mathcal{P} = (V_0, V_1, \dots, V_k)$  be a partition of  $V$  such that

- $\mathcal{P} \setminus \{V_0\}$  is not  $\varepsilon$ -regular (there are at least  $\varepsilon k^2$  irregular pairs),
- $\mathcal{P} \setminus \{V_0\}$  is equitable<sup>6</sup>, and
- $|V_0| \leq \varepsilon n$ .

Then, there is a refinement  $\mathcal{Q} = (V'_0, V'_1, \dots, V'_\ell)$  of  $\mathcal{P}$  such that

- $\ell \leq k4^k$ ,
- $\mathcal{Q} \setminus \{V'_0\}$  is equitable,
- $|V'_0| \leq |V_0| + n/2^k$ , and
- $q(\mathcal{Q}) \geq q(\mathcal{P}) + \varepsilon^5/2$ .

*Proof.* Let  $|V_i| = t$  for all  $1 \leq i \leq k$ .

Suppose  $(V_i, V_j)$  is an  $\varepsilon$ -irregular pair in  $\mathcal{P}$  and let  $(V_{i,1}, V_{i,2})$  and  $(V_{j,1}, V_{j,2})$  be the partitions of  $V_i, V_j$  respectively described in Lemma 2.11. If  $\mathcal{Q}_1$  is this particular refinement

$$\mathcal{P} \cup \{V_{i,1}, V_{i,2}, V_{j,1}, V_{j,2}\} \setminus \{V_i, V_j\}$$

of  $\mathcal{P}$ , then

$$q(\mathcal{Q}_1) \geq q(\mathcal{P}) + \varepsilon^4 \frac{|V_i||V_j|}{n^2}$$

Let us similarly produce refinements corresponding to all irregular pairs of  $\mathcal{P}$ . Let  $\mathcal{Q}'$  be the “common” refinement of all these partitions  $(\mathcal{Q}_i)$ . That is, for each  $v \in V$ ,  $v$  is placed in the subset  $\bigcap_{W \in \mathcal{Q}_i: v \in W} W$ .

We then have

$$\begin{aligned} q(\mathcal{Q}') &\geq q(\mathcal{P}) + \varepsilon^4 \cdot \frac{t^2}{n^2} (\varepsilon k^2) \\ &= q(\mathcal{P}) + \varepsilon^5 \frac{(tk)^2}{n^2} \\ &\geq q(\mathcal{P}) + \frac{\varepsilon^5}{2}. \end{aligned} \quad (tk = n - |V_0| \geq (1 - \varepsilon)n \geq 3n/4 \text{ and } 9/16 \geq 1/2) \quad (2.7)$$

To make the partition  $\mathcal{Q}'$  equitable, create the partition  $\mathcal{Q}$  using it as follows.

Suppose we partition  $V_i$  into  $(V_{ij})$ , where  $V_{ij}$  is formed due to the irregularity of  $(V_i, V_j)$ . Partition each of these parts of  $\mathcal{Q}'$  into sets of size  $b := \lfloor t/4^k \rfloor$ . Whatever residual part cannot be cut out in this manner, we merge with  $V_0$ .

Since partitioning can only increase energy, this operation will only strengthen (2.7), if anything.

---

<sup>6</sup>All the blocks of the partition are of equal size.

Since the size of any block of  $\mathcal{Q}$  is  $b$ , the number of blocks is at most  $n/b = k4^k$ .

Finally, what is the size of  $|V'_0|$ ?

$\mathcal{Q}'$  has at most  $k \cdot 2^{k-1}$  parts (there are at most  $(k-1)$  2-part partitions of each block of  $\mathcal{P}$ , and together they give a partition of size at most  $2^{k-1}$ ). So,

$$\begin{aligned} |V'_0| &\leq |V_0| + |\mathcal{Q}'|b \\ &\leq |V_0| + k2^{k-1} \frac{t}{4^k} \\ &\leq |V_0| + \frac{n}{2^k}, \end{aligned}$$

as desired. ■

Observe that with the above lemma, **Szemerédi's Regularity Lemma** follows without too much difficulty.

Begin with a partition  $\mathcal{P}_0$  with  $k_0$  parts, where  $2^{k_0} \geq 2/\varepsilon$ .

Given  $\mathcal{P}_k$ , let  $\mathcal{P}_{k+1}$  be the partition defined by Lemma 2.12. Since the energy of any partition is bounded above by 1, this process must terminate after at most  $2/\varepsilon^5$  steps.

Further, since the size of the refined partition is bounded between quantities dependent solely on the old partition, the size of the final partition after termination of the above process is at most some quantity dependent only on  $\varepsilon$  (this quantity might be massive<sup>7</sup>, but that is besides the point).

## 2.4. Some Corollaries of Szemerédi's Regularity Lemma

In this section, we cover several interesting corollaries of **Szemerédi's Regularity Lemma**.

Before we begin however, how do we process the regularity lemma? We typically begin with a “cleaning” of the graph as follows.

- Given  $\varepsilon > 0$ , get a partition as described in the lemma.
- First, delete all edges between irregular pairs. This loses at most  $(\varepsilon k^2)t^2 < \varepsilon n^2$  edges.
- Delete all edges between “sparse” pairs, where we say that a pair  $(V_i, V_j)$  is sparse if  $d(V_i, V_j) < \varepsilon$  (say). This loses at most  $k^2(\varepsilon t^2) < (\varepsilon/2)n^2$  edges.
- Delete all edges inside the exceptional part  $V_0$ . This loses at most  $(\varepsilon^2/2)n^2$  edges.

All of these deletions cost at most  $2\varepsilon n^2$  edges. In the remaining graph, *all* pairs  $(V_i, V_j)$  for  $1 \leq i, j \leq k$  are  $\varepsilon$ -regular. Further, if  $e(V_i, V_j) > 0$ , then  $d(V_i, V_j) \geq \varepsilon$ .

### 2.4.1. The graph counting and removal lemmas

**Theorem 2.13** (Triangle Counting Lemma). Suppose  $V_1, V_2, V_3$  forms a pairwise  $\varepsilon$ -regular partition of the vertex set of graph  $G_n$ , and that  $d(V_i, V_j) \geq d$  (for some  $d \geq 2\varepsilon$ ). Then, there are at least

$$\varepsilon^2(1 - 2\varepsilon)(d - \varepsilon)|V_1||V_2||V_3|$$

triangles  $xyz$  with  $x \in V_1, y \in V_2, z \in V_3$ .

<sup>7</sup>Unbelievably so. For example,  $\varepsilon = 1/8$  gives a bound of the order of  $4 \uparrow\uparrow 2^{15}$  (using **Knuth's up-arrow notation**). It further turns out that such a massive tetration-type bound is necessary, as proved in [Gow97].

*Proof.* Let

$$X_1 = \{x \in V_1 : d(x, V_2) \leq \varepsilon|V_2|\}.$$

We claim that  $|X_1| < \varepsilon|V_1|$ . Suppose otherwise. Then  $e(X_1, V_2)/|X_1||V_2| < \varepsilon$  and further,  $\varepsilon$ -regularity implies that  $d(X_1, V_2) \geq (d - \varepsilon)$ , which leads to a contradiction.

Similarly,  $X_2 := \{x \in V_1 : d(x, V_3) \leq \varepsilon|V_3|\}$  has size less than  $\varepsilon|V_1|$  too.

For all  $x \in V_1 \setminus (X_1 \cup X_2)$ ,  $d(x, V_2) \geq \varepsilon|V_2|$  and  $d(x, V_3) \geq \varepsilon|V_3|$ . Fix such an  $x$  and let  $X'_i = \Gamma(x) \cap V_i$  (for  $i = 2, 3$ ).

By the  $\varepsilon$ -regularity of  $(V_2, V_3)$ ,  $d(V_2, V_3) - d(X'_2, X'_3) \leq \varepsilon$  so  $d(X'_2, X'_3) \geq d - \varepsilon$ . As a result,

$$e(X'_2, X'_3) \geq (d - \varepsilon)|X'_2||X'_3| \geq (d - \varepsilon)\varepsilon^2|V_2||V_3|.$$

Therefore, the number of triangles of the desired form is at least

$$\underbrace{(1 - 2\varepsilon)|V_1|}_{\text{the number of such } x} \cdot \underbrace{(d - \varepsilon)\varepsilon^2|V_2||V_3|}_{\text{the number of triangles corresponding to each } x},$$

completing the proof. ■

Along similar lines is the following result.

**Theorem 2.14** (Triangle Removal Lemma). Given  $\varepsilon > 0$ , there exists  $\delta$  (depending only on  $\varepsilon$ ) such that for  $n \gg 0$ , any graph  $G_n$  with at most  $\delta n^3$  triangles can be made triangle-free by deleting at most  $\varepsilon n^2$  edges.

*Proof.* Start with an  $(\varepsilon/4)$  regular partition of  $G_n$  using Szemerédi's Regularity Lemma and the cleaning process from earlier. Delete all edges within each of the  $V_i$ . This costs at most  $k(t^2/2) < (kt)^2/2k \leq \varepsilon n^2/4$  edges.

So, we have lost  $\leq \varepsilon n^2$  edges in all. If there is a triangle remaining in the graph, it must come from a triple  $(V_i, V_j, V_k)$  with all three pairs being  $(\varepsilon/4)$ -regular and density at least  $\varepsilon/2$ . We can then use the triangle counting lemma to conclude that there are at least  $t^3(\varepsilon/4)^3(1 - \varepsilon/2) > n^3(\varepsilon/8)^3/M(\varepsilon)^3$ . Letting  $\delta$  to be the  $1/6$  this quantity, we are done. ■

Let  $(V_1, V_2)$ ,  $(V_2, V_3)$ , and  $(V_3, V_1)$  all be  $\varepsilon$ -regular pairs with densities  $d_{12}, d_{23}, d_{13} > 2\varepsilon$ . A random graph on  $V_1 \sqcup V_2 \sqcup V_3$  with edge probabilities  $d_{ij}$  between  $V_i$  and  $V_j$  has an expected number of triangles of  $|V_1||V_2||V_3|d_{12}d_{23}d_{31}$ . Ideally, we would have a result that the number of triangles is indeed to this quantity (so the graph behaves almost randomly, in a sense similar to that in Szemerédi's Regularity Lemma).

**Theorem 2.15** (Graph Counting Lemma). Let  $G$  be a graph on  $n$  vertices and  $H$  a graph on  $[k]$ . Let  $V_1, \dots, V_k \subseteq V(G)$  such that  $(V_i, V_j)$  is  $\varepsilon$ -regular whenever  $ij \in E(H)$ . Then,

$$|\{(v_1, \dots, v_k) : v_i \in V_i \text{ and } \{v_1, \dots, v_k\} \text{ form a copy of } H \text{ in } G\}|$$

is within  $\varepsilon e(H)|V_1||V_2| \cdots |V_k|$  of

$$\prod_{i=1}^k |V_i| \prod_{ij \in E(H)} d(V_i, V_j),$$

assuming that  $\prod_{ij \in E(H)} d(V_i, V_j) > \varepsilon e(H)$ .

*Proof.* We shall prove this by inducting on  $e(H)$ . If  $e(H) = 0$ , the result is trivial.

Let us rephrase the problem probabilistically. Pick  $v_i \in V_i$  independently and uniformly. Then, we wish to prove that

$$\left| \Pr[v_i v_j \in E(G) \text{ for all } ij \in E(H)] - \prod_{ij \in E(H)} d(V_i, V_j) \right| \leq \varepsilon e(H). \quad (2.8)$$

Suppose  $\{1, 2\} \in E(H)$ . It suffices to show that

$$\left| \Pr[v_i v_j \in E(G) \text{ for all } ij \in E(H)] - d(V_1, V_2) \Pr[v_i v_j \in E(G) \text{ for all } E(H) \setminus \{1, 2\}] \right| \leq \varepsilon. \quad (2.9)$$

Indeed, by induction,

$$\left| \Pr[v_i v_j \in E(G) \text{ for all } E(H) \setminus \{1, 2\}] - \prod_{\substack{ij \in H \\ ij \neq \{1, 2\}}} d(V_i, V_j) \right| < \varepsilon(e(H) - 1), \quad (2.10)$$

and (2.8) follows from (2.9) and (2.10) on using the **union bound**. We shall prove that (2.9) holds when we condition on the choices  $v_i$  for  $i > 2$ , and thus generally. Let

$$A_j = \{v_j \in V_j : \{v_j, v_i\} \in E(G) \text{ when } \{j, i\} \in E(H)\}.$$

for  $j = 1, 2$ . Equation (2.9) is then equivalent to

$$\left| \frac{e(A_1, A_2)}{|V_1||V_2|} - d(V_1, V_2) \frac{|A_1||A_2|}{|V_1||V_2|} \right| \leq \varepsilon. \quad (2.11)$$

We claim that (2.11) holds for all  $A_1, A_2$ . If  $|A_1| \geq \varepsilon|V_1|$  and  $|A_2| \geq \varepsilon|V_2|$ , then  $\varepsilon$ -regularity implies that  $|d(A_1, A_2) - d(V_1, V_2)| \leq \varepsilon$ , that is,

$$\left| \frac{e(A_1, A_2)}{|A_1||A_2|} - d(V_1, V_2) \right| \leq \varepsilon.$$

So,

$$\frac{|V_1||V_2|}{|A_1||A_2|} \left| \frac{e(A_1, A_2)}{|V_1||V_2|} - d(V_1, V_2) \frac{|A_1||A_2|}{|V_1||V_2|} d(V_1, V_2) \right| \leq \varepsilon,$$

and (2.11) easily follows. If  $|A_1| < \varepsilon|V_1|$ , then the above follows immediately anyway, since

$$\begin{aligned} \left| \frac{e(A_1, A_2)}{|V_1||V_2|} - d(V_1, V_2) \frac{|A_1||A_2|}{|V_1||V_2|} \right| &= \frac{|A_1||A_2|}{|V_1||V_2|} \left| \frac{e(A_1, A_2)}{|A_1||A_2|} - d(V_1, V_2) \right| \\ &= \frac{|A_1||A_2|}{|V_1||V_2|} |d(A_1, A_2) - d(V_1, V_2)| \\ &\leq \frac{|A_1|}{|V_1|} < \varepsilon, \end{aligned}$$

so we are done. ■

The generalized version of the triangle removal lemma is the following.

**Theorem 2.16** (Graph Removal Lemma). Given  $\varepsilon > 0$  and any graph  $H$ , there exists  $\delta$  (depending only on  $\varepsilon$ ) such that for  $n \gg 0$ , any graph  $G_n$  with at most  $\delta n^{V(H)}$  subgraphs isomorphic to  $H$  can be made  $H$ -free by deleting at most  $\varepsilon n^2$  edges.

#### 2.4.2. Roth's Theorem and Corners

Next, we describe Roth's Theorem. The result deals with a conjecture of Erdős and Turán:

Given  $\varepsilon > 0$  and  $r \in \mathbb{N}$ , there exists  $N_0$  such that for all  $N \geq N_0$ , the following holds. If  $A \subseteq [N]$  with  $|A| \geq \varepsilon N$ , then  $A$  contains an arithmetic progression of length  $r$ .

Roth proved in the early 50s that the conjecture holds for  $r = 3$ . Szemerédi proved in the early 70s that it holds for all  $r$ . In fact, Szemerédi had originally come up with the regularity lemma to prove this result (this is why it is a *lemma*).

**Lemma 2.17.** Suppose that every edge of  $G_n$  is in exactly one triangle. Then,  $e(G_n) = o(n^2)$ .

How is this related to Roth's Theorem? Given  $N \gg 0$  and  $A \subseteq [N]$ , suppose  $A$  is 3-AP free. Let  $M = 2N + 1$  and construct a 3-partite graph  $G$  whose three parts  $X, Y$ , and  $Z$  are copies of  $\mathbb{Z}/M$ .

For  $x \in X, y \in Y$ , and  $z \in Z$ , keep an edge from  $x$  to  $y$  iff  $y - x \in A$ ,  $y$  to  $z$  iff  $z - y \in A$ , and  $z$  to  $x$  iff  $(z - x)/2 \in A$ . The final part is well-defined since 2 is invertible in  $\mathbb{Z}/M$  (Why?).

Observe that if  $xyz$  is a triangle with  $a = y - x, b = z - y$ , and  $2c = z - x$ , then  $a + b = 2c$ , so  $a, c, b$  are in AP.

Since  $a, b$ , and  $c$  are all in  $A$ , then by our assumption we must have  $a = b = c, y - x = z - y = (z - x)/2$ , and so  $x, y, z$  must be in AP in  $\mathbb{Z}/M$ .

This implies that any edge must be in a unique triangle.

By Lemma 2.17,  $e(G) = o((3M)^2) = o(N^2)$ . On the other hand,  $e(G) = 3M|A|$ . So,  $3M|A| = o(N^2)$  and  $|A| = o(N)$ .

*Proof of Lemma 2.17.* The number of triangles in  $G_n$  is exactly  $e(G_n)/3 = o(n^3)$ . By the **Triangle Removal Lemma**,  $G_n$  can be made triangle-free by removing  $o(n^2)$  edges. However, we must remove at least  $e(G_n)/3$  edges to make the graph triangle-free, so  $e(G_n) = o(n^2)$ . ■

Inspired by this problem, a natural question to ask is: what is the maximum sized  $A \subseteq [N]$  that is 3-AP free?

Picking  $A$  greedily gives  $O(\sqrt{n})$  elements.

It is possible to do better, as shown by *Behrend's construction*.

The basic idea is that on a sphere, the midpoint of any two points does not lie on the sphere.

Consider the set  $S = [k]^d \subseteq \mathbb{R}^d$  for some  $k, d$ . Observe that  $\|x\|^2 \leq k^2 d$  for any point in  $S$ . By the pigeonhole principle, there is a (centered) sphere with at least  $k^{d-2}/d$  of these points.

To get a subset  $A$  from these points, project  $x = (a_1, \dots, a_d)$  to  $\sum_{j=1}^d a_j(2k+1)^{j-1}$ , thus converting it to an integer in base  $(2k+1)$ . Observe that if  $x_1 \mapsto m_1$  and  $x_2 \mapsto m_2$ , then  $(x_1 + x_2)/2 \mapsto (m_1 + m_2)/2$ . As a result, if  $X$  is a subset of  $S$  on a single sphere, its projection by this map gives a 3-AP free subset.

The maximum element by this projection is  $k \sum_{i=0}^{d-1} (2k+1)^i = ((2k+1)^d - 1)/2$ . Set  $(2k+1)^d = 2N+1$ . By the argument from before, there is a 3-AP free set of size  $k^{d-2}/d$ .

This achieves a bound of

$$\frac{N}{e^{c\sqrt{\log n}}}$$

for some constant  $c$ . This is  $\Omega(N^{1-\delta})$  for any (fixed)  $\delta > 0$ !

**Definition 2.7 (Corner).** An  $A \subseteq [N]^2$  is said to have a **corner** if the points  $(x, y)$ ,  $(x + d, y)$ , and  $(x, y + d)$  are in  $A$  for some  $d > 0$  and  $x, y \in [N]$ .

**Theorem 2.18 (No Corners Theorem).** Suppose  $A \subseteq [N]^2$  has no corners. Then  $|A| = o(N^2)$ .

*Proof.* First, let us get rid of the  $d > 0$  clause.

Denote by  $A + A$  the set  $\{a + b : a, b \in A\} \subseteq [2N]^2$  (the **Minkowski sum** of the two sets) and by  $x + A$   $\{x + a : a \in A\}$ . By the pigeonhole principle, there exists a  $z \in [2N]^2$  such that  $z = a + b$  for at least  $|A|^2/4N^2$  pairs  $(a, b) \in A^2$ .

Consider  $A' = A \cap (z - A)$  for such a  $z$ . Then

$$|A'| \geq \frac{|A|^2}{4N^2}$$

by the definition of  $z$ . It is not too difficult to see that  $A' = (z - A')$ . As a result, there is a correspondence between corners with positive  $d$  and negative  $d$  in  $A'$ . Further, if  $A$  is corner-free, so is  $A'$ . By the cardinality bound on  $A'$ , it suffices to show that  $|A'| = o(N^2)$ .

So, let us drop the  $d > 0$  condition on corners and work with a corner-free  $A$ .

We construct a 3-partite graph  $G$ . Let the three parts of the vertex set  $V$  be the set  $H$  of horizontal lines in  $[N]^2$ , the set  $V$  of vertical lines in  $[N]^2$ , and the set  $D$  of lines with slope  $-1$  in  $[2N]^2$ .

Given  $\ell, \ell'$  in distinct parts of  $V$ , let  $\ell$  and  $\ell'$  be adjacent in  $G$  iff  $\ell \cap \ell' \in A$ . Triangles in this graph correspond to either corners or three lines all passing through a single point (that is in  $A$ ). However, there are no corners in  $A$ , so each edge is in a unique triangle. The result then follows from Lemma 2.17. ■

#### 2.4.3. A weaker version of the Erdős-Stone-Simonovits Theorem

Next, we shall prove a slightly weaker version of the **Erdős-Stone-Simonovits Theorem, Version 2** using **Szemerédi's Regularity Lemma**.

**Theorem 2.19** (Erdős-Stone-Simonovits Theorem, Version 3). Fix  $t \in \mathbb{N}$  and  $\varepsilon > 0$ . For  $n \gg 0$ , if  $G_n$  is  $K_{r+1}(t)$ -free,<sup>8</sup> then

$$e(G_n) \leq \left(1 - \frac{1}{r}\right) \frac{n^2}{2} + \varepsilon n^2.$$

This is weaker than the original result we saw since here,  $t$  is fixed and so cannot grow to the  $\log n$  estimate we gave earlier.

*Proof.* Suppose  $e(G_n) > (1 - 1/r)n^2/2 + \varepsilon n^2$  edges.

Let  $\delta > 0$  which we shall fix later. By Szemerédi's regularity lemma, we get a partition  $(V_0, V_1, \dots, V_k)$  such that  $|V_0| \leq \delta n$ ,  $|V_i| = \ell$  (for some  $\ell$ ) for  $1 \leq i \leq k$ , and at most  $\delta k^2$  of the pairs are  $\delta$ -irregular.

'Clean' the graph by deleting all edges within the  $V_i$  (losing  $\leq \ell^2 k/2$  edges), edges adjacent to  $V_0$  (losing  $\leq \delta n^2$  edges), edges in irregular pairs (losing  $\leq \delta k^2 \ell^2$  edges), and edges between pairs of density less than  $\delta$  (losing  $\leq \ell \delta (k^2/2)$  edges).

The number of deleted edges is thus at most  $4\delta n^2$ . As a result, the graph after purification still has  $> (1 - 1/r)n^2/2 + (\varepsilon - 4\delta)n^2$  edges.

Let  $H$  be a graph with vertex set  $\{V_i : 1 \leq i \leq k\}$ , and  $V_i V_j \in E(H)$  iff  $(V_i, V_j)$  is  $(\varepsilon/8)$ -regular with density at least  $\varepsilon/4$ .

By the pigeonhole principle, since  $e(V_i, V_j) \leq \ell^2$

$$e(H) \geq \left(1 - \frac{1}{r} + \varepsilon\right) \frac{n^2}{2\ell^2} \geq \left(1 - \frac{1}{r} + \varepsilon\right) \frac{k^2}{2}.$$

By Turán's Theorem,  $H$  contains  $K_{r+1}$ . Say  $(V_1, \dots, V_{r+1})$  form this  $K_{r+1}$ . All  $(V_i, V_j)$  are  $(\varepsilon/8)$ -regular with density at least  $\varepsilon/4$ .

We now use the **Graph Counting Lemma**. Take  $F = K_{r+1}(t)$ . The number of copies of  $F$  in  $(V_1, \dots, V_{r+1})$  is at least

$$\prod_{i=1}^{r+1} |V_i|^t \left( (\varepsilon/4)^{t^2 \binom{r+1}{2}} - \delta t^2 \binom{r+1}{2} \right).$$

So, if

$$\delta \leq \frac{(\varepsilon/4)^{t^2 \binom{r+1}{2}}}{2t^2 \binom{r+1}{2}},$$

then there are several copies of  $K_{r+1}$  in  $G$ , completing the proof. ■

<sup>8</sup>This is the complete  $(r+1)$ -partite graph with  $t$  vertices in each part, that is,  $T_{r+1}(t(r+1))$ .

## 2.4.4. An application in computer science

Now, we give a delightful application from computer science.

Suppose we are given a graph  $G_n$  and we wish to check (algorithmically) whether  $G$  has any triangles. An easy way to do this is to iterate through all triples and check if any of them form a triangle, which takes  $\mathcal{O}(n^3)$  time.

In this context, we give a paradigm due to Shafi Goldwasser and Madhu Sudan. We want to come up with an algorithm that takes a *constant* number of verifications, and still returns a reasonable output. It is clearly impossible to return the answer correctly all the time. The output we desire is such that with high probability,

- if  $G_n$  has no triangles, our output is correct.
- if  $G_n$  has triangles and we output otherwise, then  $G_n$  must be “close” to triangle-free.

Such a randomized constant time algorithm is known as a **property tester**. The question we wish to answer is: which graph properties admit such testers?

For starters, does triangle-freeness admit a property tester? It turns out that it *does*! The algorithm is in fact very simple.

- Pick a random triple  $(x, y, z)$  and check if it forms a triangle.
- If yes, return “not triangle-free”.
- If no, return “ $\varepsilon$ -close to triangle-free”, where at most  $\varepsilon n^2$  edges need to be deleted to remove all triangles.

Why is this a tester?

By the **Triangle Removal Lemma**, if  $G_n$  is  $\varepsilon$ -close to being triangle-free, then  $G_n$  has at most  $\delta n^3$  triangles for some  $\delta$ . That is,

$$\Pr[xyz \text{ is not a triangle}] \geq 1 - \frac{\delta n^3}{\binom{n}{3}} \geq 1 - 6\delta,$$

so our algorithm is indeed a property tester.

We shall return to property testing in more detail later.

## 2.4.5. The Frankl-Pach Theorem

**Theorem 2.20** (Frankl-Pach Theorem). Suppose  $s, r \in \mathbb{N}$  and  $0 < c < 1$ . Then, for  $n \gg 0$ ,

$$\text{ex}(n; K_{r+1}, K_{s, \lceil cn \rceil}) \leq \left( c^{1/s} \left( 1 - \frac{1}{r} \right)^{1/s} + o(1) \right) n^2.$$

Observe that one of the graphs we are forbidding depends on  $n$  itself.

*Proof.* Let  $\varepsilon > 0$  and  $G$  be a graph on  $n$  vertices that is  $K_{r+1}$ - and  $K_{s, \lceil cn \rceil}$ -free. Using Szemerédi’s regularity lemma and the cleaning process, obtain a graph  $G'$  such that

- $V(G') = V_1 \sqcup \dots \sqcup V_k$  where  $k = O_\delta(1)$ ,
- $|V_i| = \ell$  for all  $i$  and  $V(G') \geq (1 - \delta)n$ ,
- All pairs  $(V_i, V_j)$  are  $\delta$ -regular,
- if  $e(V_i, V_j) > 0$ ,  $d(V_i, V_j) \geq \eta$ , and
- $e(G) - e(G') \leq \varepsilon n^2$

for some  $\delta, \eta > 0$  depending only on  $\varepsilon$ . Henceforth, we operate only in  $G'$ .

Construct a graph  $H$  on vertex set  $[k]$  with  $ij \in E(H)$  iff  $d(V_i, V_j) \geq \eta$ . By the **Graph Counting Lemma**, if  $H$  contains  $K_{r+1}$ , so does  $G'$  and thus  $G$ . Therefore,  $H$  is  $K_{r+1}$ -free and by Turán's Theorem,  $e(H) \leq (1 - 1/r)k^2/2$ .

Since  $n \gg 0$  and  $k = O_\delta(1)$ , if  $K_{s, \lceil cn \rceil}$  appears in  $G$ , one of the  $V_i$ s must have at least  $s$  vertices for  $n \gg 0$  (by a simple pigeonhole argument).

So, let us bound  $e(G)$  given that  $K_{s, \lceil cn \rceil}$  is forbidden and the size  $s$  part of the bipartite graph does not come from a single  $V_i$ .

Similar to what we did in the **Kővári-Sós-Turán Theorem**, let us count pairs  $(x, S)$  where  $|S| = s$ ,  $S \subseteq V_i$  for some  $i$ , and  $x$  is adjacent to every  $y \in S$ . We call such a structure an  $(i, s)$ -claw. We have that

$$\text{number of } (i, s)\text{-claws} = \sum_{x \in V'} \sum_{i=1}^k \binom{|N(x) \cap V_i|}{s}.$$

On the other hand, by our assumption regarding  $K_{s, \lceil cn \rceil}$ -freeness,

$$\text{number of } (i, s)\text{-claws} \leq (\lceil cn \rceil - 1) \sum_{i=1}^k \binom{|V_i|}{s} \leq cnk \binom{\ell}{s}.$$

Therefore,

$$\sum_{x \in V'} \sum_{i=1}^k \binom{|N(x) \cap V_i|}{s} \leq cnk \binom{\ell}{s}.$$

Define  $\mathcal{P} = \{(x, i) : |N(x) \cap V_i| > 0\}$ . Applying Jensen's inequality in a manner similar to that in proof of the Kővári-Sós-Turán Theorem,

$$\begin{aligned} \sum_{(x,i) \in \mathcal{P}} \binom{|N(x) \cap V_i|}{s} &\geq |\mathcal{P}| \binom{(1/|\mathcal{P}|) \sum_{(x,i) \in \mathcal{P}} |N(x) \cap V_i|}{s} \\ &= |\mathcal{P}| \binom{2e(G')/|\mathcal{P}|}{s} \\ &= |\mathcal{P}| \binom{u}{s}, \end{aligned}$$

where  $u = 2e(G')/|\mathcal{P}|$ . Then,

$$\begin{aligned} cnk \frac{\ell^s}{s!} &\geq cnk \binom{\ell}{s} \\ &\geq \sum_{x \in V'} \sum_{i=1}^k \binom{|N(x) \cap V_i|}{s} \\ &= \sum_{(x,i) \in \mathcal{P}} \binom{|N(x) \cap V_i|}{s} \\ &\geq |\mathcal{P}| \binom{u}{s} \\ &\geq |\mathcal{P}| \frac{(u - s + 1)^s}{s!}. \end{aligned}$$

Simplifying,

$$e(G') \leq \frac{u|\mathcal{P}|}{2} \leq \frac{1}{2} \left( (cnk)^{1/s} \ell |\mathcal{P}|^{1-1/s} + |\mathcal{P}|(s-1) \right). \quad (2.12)$$

To bound  $|\mathcal{P}|$  from above, observe that

$$|\mathcal{P}| \leq e(H) \cdot 2\ell$$



since each edge in  $H$  gives at most  $2\ell$  choices for  $x$  (if the edge is  $ij$ ,  $x$  is in  $V_i$  or  $V_j$ ). By  $K_{r+1}$ -freeness,  $e(H) \leq (1 - 1/r)k^2/2$  so

$$|\mathcal{P}| \leq \left(1 - \frac{1}{r}\right) k^2 \ell.$$

Substituting the above back in (2.12),

$$\begin{aligned} e(G') &\leq \frac{1}{2} \left( (cnk)^{1/s} \ell |\mathcal{P}|^{1-1/s} + |\mathcal{P}|(s-1) \right) \\ &\leq \frac{1}{2} \left( (cn)^{1/s} \left(1 - \frac{1}{r}\right)^{1/s} (k\ell)^{2-1/s} + \left(1 - \frac{1}{r}\right) k^2 \ell (s-1) \right) \\ &\leq \frac{1}{2} \left( c^{1/s} \left(1 - \frac{1}{r}\right)^{1/s} n^2 + \left(1 - \frac{1}{r}\right) k(s-1)n \right) \\ &= \left( c^{1/s} \left(1 - \frac{1}{r}\right)^{1/s} + o(1) \right) \frac{n^2}{2}. \end{aligned}$$

The result follows since  $e(G) - e(G') \leq \varepsilon n^2$ . ■

It turns out that the bound given by the Frankl-Pach Theorem is asymptotically tight!

#### 2.4.6. A look at $\text{ex}(n; C_{2k+1})$

As we saw back in Corollary 2.3,

$$\frac{n^2}{4} \leq \text{ex}(n; C_{2k+1}) \leq \left( \frac{1}{4} + o(1) \right) n^2.$$

Recall that if  $e(G_n) \geq n^2/4 - t$  and  $G_n$  is  $K_3$ -free, then  $G_n$  can be made bipartite by deleting at most  $t$  edges.

**Lemma 2.21.** Given  $\varepsilon > 0$ , there exists  $\delta > 0$  such that the following holds for  $n \gg 0$ . If  $G_n$  is  $C_{2k+1}$ -free and has  $\geq (1/4 - \delta)n^2$  edges, it can be made bipartite by deleting at most  $\varepsilon n^2$  edges.

*Proof.* First, we shall try to make  $\delta(G_n)$  ‘large’. If  $G_n$  has a vertex of degree  $< (1/2 - 2\sqrt{\delta})n$ , delete this vertex to get a graph  $G_{n-1}$ . Perform this process repeatedly to finally arrive at some  $G_\ell$  such that

$$\begin{aligned} e(G_\ell) &> \left( \frac{1}{4} - \delta \right) n^2 - \left( \frac{1}{2} - 2\sqrt{\delta} \right) (n + (n-1) + \cdots + (\ell+1)) \\ &= \left( \frac{1}{4} - \delta \right) n^2 - \left( \frac{1}{2} - 2\sqrt{\delta} \right) \left( \binom{n+1}{2} - \binom{\ell+1}{2} \right) \\ &= \left( \frac{1}{4} - \delta \right) n^2 - \left( \frac{1}{4} - \sqrt{\delta} \right) (n^2 + n - \ell^2 - \ell) \\ &\geq \left( \frac{1}{4} - \delta \right) \ell^2 + (\sqrt{\delta} - \delta)n^2 - \left( \frac{1}{4} - \delta \right) n. \end{aligned}$$

If the quantity above is at least  $(1/4 + \eta)\ell^2$  for some constant  $\eta > 0$ , we have

$$\ell < \sqrt{\frac{\sqrt{\delta} - \delta}{\eta + \delta}} n.$$

This implies that for sufficiently large  $\ell$  (by virtue of sufficiently large  $n$ ),  $G_\ell$  has  $C_{2k+1}$  which yields a contradiction. Therefore, setting  $\eta$  appropriately, this process must terminate with  $\ell \geq (1 - 4\sqrt{\delta})n$ .

Therefore, suppose  $G' = (V', E')$  is a subgraph of  $G_n$  such that  $|V'| = n' \geq (1 - 4\sqrt{\delta})n$ ,  $\delta(G') \geq (1/2 - 2\sqrt{\delta})n'$ , and  $G'$  is  $C_{2k+1}$ -free. We have  $e(G') \geq (1/4 - \sqrt{\delta})n'^2$ .

Recall that  $\text{ex}(n; C_{2k})$  is  $o(n^2)$ . As a result, it follows that if  $n \gg 0$ ,  $G'$  contains a copy of  $C_{2k}$ , say  $v_1 v_2 \cdots v_{2k} v_1$ . Letting  $W = V' \setminus \{v_1, \dots, v_k\}$ , we must have that  $A_1 = N(v_1) \cap W$  and  $A_2 = N(v_2) \cap W$  are disjoint. Since  $G'$  is  $C_{2k+1}$ -free, the subgraph induced on  $A_1$  (or  $A_2$ ) is  $P_{2k}$ -free. By Theorem 1.7,  $e(A_1)$  (or  $e(A_2)$ ) is at most  $kn'$ . To summarize,

- $|A_1|, |A_2| \geq (1/2 - 2\sqrt{\delta})n' - 2k \geq (1/2 - \sqrt{\delta})n'$ .
- $e(A_1), e(A_2) \leq kn'$ .
- $A_1 \cap A_2 = \emptyset$ .

Delete all edges that are not between  $A_1$  and  $A_2$  (namely those within the  $A_i$ s and those that touch at least one vertex not in  $A_1 \cup A_2$ ). This deletes at most

$$\underbrace{2kn'}_{e(A_i)} + \underbrace{2\sqrt{\delta}n'^2}_{|V' \setminus (A_1 \cup A_2)| \geq 2\sqrt{\delta}n'}.$$

The resulting graph is clearly bipartite. Thus, from  $G_n$ , we have deleted at most, say,  $5\sqrt{\delta}n^2$  edges, so setting  $\delta$  less than say  $\varepsilon^2/100$  works.  $\blacksquare$

**Theorem 2.22.** For  $n \gg 0$ ,

$$\text{ex}(n; C_{2k+1}) = \left\lfloor \frac{n^2}{4} \right\rfloor.$$

*Proof.* Let  $G_n$  be  $C_{2k+1}$ -free and have the maximum possible number of edges. Since the complete bipartite graph is  $C_{2k+1}$ -free,  $e(G_n) \geq \lfloor n^2/4 \rfloor$ .

Let  $\varepsilon > 0$ . For starters, let us assume that  $\delta(G_n) \geq (1/2 - 2\sqrt{\varepsilon})n$ . By Lemma 2.21,  $G_n$  can be made bipartite by deleting at most  $\varepsilon n^2$  edges (due to the assumption on  $\delta(G_n)$  and the previous proof). Let  $(A, B)$  be the parts of such a bipartite graph. Further assume that  $e(A, B)$  is the maximum possible (we choose that bipartite subgraph with the most edges). We claim that  $|A|$  and  $|B|$  are at least  $(1/2 - \varepsilon)n$ . Indeed, otherwise, since  $|A| + |B| = n$ ,

$$e(A, B) \leq |A||B| < \left(\frac{1}{2} - \sqrt{\varepsilon}\right) \left(\frac{1}{2} + \varepsilon\right) n^2 = \left(\frac{1}{4} - \varepsilon\right) n^2.$$

In this case,  $e(G_n) \leq e(A, B) + \varepsilon n^2 < \lfloor n^2/4 \rfloor$ , proving the claim.

Since  $e(A, B)$  is the maximum possible, it follows that for any  $a \in A$ ,  $d(a, B) \leq d(a, A)$  (otherwise we can switch  $a$  from  $A$  to  $B$ ).

Next, we claim that no vertex  $a$  of  $A$  is such that  $d(a, A) \geq 2\sqrt{\varepsilon}n$ . Suppose otherwise. We then have  $d(a, B) \geq 2\sqrt{\varepsilon}n$  as well. Let  $A_1 = A \cap \Gamma(a)$  and  $B_1 = B \cap \Gamma(a)$ . Consider the bipartite subgraph on  $(A_1, B_1)$ . This subgraph is  $P_{2k}$ -free. By Theorem 1.7,

$$e(A_1, B_1) \leq 2kn.$$

In particular, the number of missing edges between  $A_1$  and  $B_1$  is at least  $4\varepsilon n^2 - 2kn > \varepsilon n^2 \geq e(A) + e(B)$  for  $n \gg 0$ . As a result,

$$\begin{aligned} e(G_n) &= e(A, B) + e(A) + e(B) \\ &\leq e(A, B) + (\text{number of missing pairs between } A_1 \text{ and } B_1) \\ &< |A||B| \leq n^2/4, \end{aligned}$$

which gives a contradiction.

Let  $a, a' \in A$  be adjacent. We have

$$\begin{aligned}
 |(\Gamma(a) \cap B) \cap (\Gamma(a') \cap B)| &\geq d(a, B) + d(a', B) - |B| \\
 &> d(a) - 2\sqrt{\varepsilon}n + d(a') - 2\sqrt{\varepsilon}n - \left(\frac{1}{2} + \sqrt{\varepsilon}\right)n \\
 &\geq 2\delta(G_n) - 4\sqrt{\varepsilon}n - \left(\frac{1}{2} + \sqrt{\varepsilon}\right)n \\
 &\geq \left(\frac{1}{2} - 9\sqrt{\varepsilon}\right)n.
 \end{aligned}$$

Let  $B' = \Gamma(a) \cap \Gamma(a') \cap B$  and  $A' = A \setminus \{a, a'\}$  and consider the bipartite subgraph between these two parts. Since the previous subgraph was  $C_{2k+1}$ -free, there cannot be a  $P_{2k-1}$  between two vertices in  $B'$  (or  $A'$ ). It follows that there does not exist a path of length  $2k$  between any two vertices of  $A' \cup B'$ . So,

$$e(A', B') \leq 2kn.$$

Estimating the number of edges again,

$$\begin{aligned}
 e(G_n) &\leq e(A', B') + e(A \setminus A', V) + e(B \setminus B', V) \\
 &= 2kn + \underbrace{2n}_{|A \setminus A'|=2} + \underbrace{10\sqrt{\varepsilon}n}_{\substack{|B| \leq (1/2 + \sqrt{\varepsilon})n \\ |B'| \geq (1/2 - 9\sqrt{\varepsilon})n}} \\
 &< \frac{n^2}{4} \text{ for } n \gg 0,
 \end{aligned}$$

giving a contradiction again.

Therefore, it now suffices to show why we can make our initial assumption that  $\delta(G_n) \geq (1/2 - 2\sqrt{\varepsilon})n$ . As in the proof of the previous lemma, repeatedly delete any vertex that has degree less than  $(1/2 - 2\sqrt{\varepsilon})n$ . We must stop at some  $G_\ell$  with  $\ell > (1 - 4\sqrt{\varepsilon})n$ . Further,  $G_\ell$  must have greater than  $\ell^2/4$  edges and is  $C_{2k+1}$ -free, so we arrive at a contradiction anyway due to our proof taking the assumption at the beginning. ■

## 2.5. Pseudorandomness

Szemerédi's regularity lemma, in layman terms, says that any dense graph consists 'mostly' of a 'small' number of 'random-like' bipartite graphs. The meaning of the first two words within the quotes should be clear; they just say that  $|V_0|$  is small and that  $k$  is bounded. The question then is: what does it mean to say that a graph is random-like? We have a single graph, not a distribution.

### 2.5.1. Notions of randomness

Let us first list a couple of features that feel like something a random-like graph should have. Suppose that  $G_n$  has  $(p + o(1))\binom{n}{2}$  edges, where  $p$  is a fixed constant.

1. *Disc (Discrepancy)*. For all  $X, Y \subseteq V(G)$ ,

$$|e(X, Y) - p|X||Y|| = o(n^2).$$

2. *Disc'*. For all  $X \subseteq V(G)$ ,

$$\left| e(X) - p \binom{|X|}{2} \right| = o(n^2).$$

3. *Count*. For a fixed  $H$ , the number of labelled copies of  $H$  in  $G$  is  $(1 + o(1))p^{e(H)}n^{|V(H)|}$  (Why choose this number?).

4. Codegree. We have

$$\sum_{x,y \in V} |d(x,y) - p^2 n| = o(n^3).$$

5. Eigen. If  $\lambda_1 \geq \dots \geq \lambda_n$  are the eigenvalues of the adjacency matrix of  $G_n$ , then

$$\lambda_1 = pn(1 + o(1))$$

and

$$\max_{i>1} |\lambda_i| = \max\{|\lambda_2|, |\lambda_n|\} = o(n).$$

Observe that there are already asymptotics at play here since  $o(\cdot)$  is not very meaningful when we are dealing with a single graph.

While the reader is likely familiar with what an adjacency matrix is, we define it here for the sake of completeness.

**Definition 2.8** (Adjacency Matrix). Given a graph  $G = (V, E)$ , the *adjacency matrix*  $A$  of  $G$  is a matrix whose rows and columns are indexed by  $V$ , and for  $v, w \in V$

$$A_{vw} = \begin{cases} 1, & vw \in E, \\ 0, & vw \notin E. \end{cases}$$

The eigenvalues of  $A$  are sometimes referred to as the eigenvalues of  $G$ .

While the first four properties above might make sense based on our discussion thus far, the last seems to come out of nowhere.

To motivate Eigen, we give the following result.

**Theorem 2.23** (Expander Mixing Lemma). Suppose  $G = (V, E)$  is a  $d$ -regular graph on  $n$  vertices and  $\lambda = \max\{|\lambda_2|, |\lambda_n|\}$ , where  $\lambda_1 \geq \dots \geq \lambda_n$  are the eigenvalues of  $G$ . Then for any  $X, Y \subseteq V$ ,

$$\left| e(X, Y) - \frac{d}{n} |X| |Y| \right| \leq \lambda \sqrt{|X| |Y|}.$$

*Proof.* Observe that

$$e(X, Y) = \mathbb{1}_X^\top A \mathbb{1}_Y,$$

where  $\mathbb{1}_X$  is the ‘indicator vector’ of  $X$ , a vector indexed by the vertices of  $G$ , with 1 at the positions in  $X$  and 0 elsewhere.

Let  $A$  be the adjacency matrix of  $G$ . Since  $A$  is real symmetric, we can apply the spectral theorem to write  $A = \sum_i \lambda_i v_i v_i^\top$ , where  $v_i$  is an eigenvector for  $\lambda_i$ .

Observe that  $d$  is an eigenvalue of  $A$  with corresponding eigenvector  $\mathbf{1}$ , the all 1s vector. In fact,  $d$  is the largest eigenvalue – if  $\lambda_1$  is the largest eigenvalue and  $x$  is an eigenvector with  $x_v = \max_u |x_u|$ ,

$$|\lambda_1 x_v| = |(Ax)_v| = \left| \sum_{u \in \Gamma(v)} x_u \right| \leq d(v) |x_v| \leq \Delta(G) |x_v|.$$

Letting  $J$  be the all 1s matrix of appropriate size,

$$\begin{aligned} \left| e(X, Y) - \frac{d}{n} |X| |Y| \right| &= \left| \mathbb{1}_X^\top A \mathbb{1}_Y - \frac{d}{n} \mathbb{1}_X^\top J \mathbb{1}_Y \right| \\ &= \left| \mathbb{1}_X^\top \left( A - \frac{d}{n} J \right) \mathbb{1}_Y \right| \\ &\leq \left\| \mathbb{1}_X^\top \right\| \left\| A - \frac{d}{n} J \right\| \left\| \mathbb{1}_Y \right\|. \end{aligned}$$

We claim that  $\|A - (d/n)J\| \leq \lambda$ . Since  $A$  is symmetric with all row sums as  $d$ , it commutes with  $J$ , and so they can be simultaneously diagonalized. Further,  $n$  is an eigenvalue of  $J$  and  $d$  is an eigenvalue of  $A$  with eigenvector  $\mathbf{1}$ . ■

Next, the following result says that all five notions of randomness we came up with are equivalent!

**Theorem 2.24** (Chung-Graham-Wilson Theorem). *Disc, Disc', Count, Codegree, and Eigen are equivalent.*

*Proof.* To prove this we shall show that all of our 5 are equivalent to the following.

$C_4$ -Count. The number of labelled  $C_4$ s is  $(p^4 + o(1))n^4$  (this is just Count for  $H = C_4$  specifically).

- Disc to Disc'. Setting  $X = Y$  then dividing by 2 gives the result directly.
- Disc' to Disc. Let  $X, Y \subseteq V$ . We have

$$e(X, Y) = e(X \cup Y) + e(X \cap Y) - e(X \setminus Y) - e(Y \setminus X)$$

Applying Disc' to each of these 4 sets gives the results near directly:

$$\begin{aligned} e(X, Y) &= p \binom{|X \cup Y|}{2} + p \binom{|X \cap Y|}{2} - p \binom{|X \setminus Y|}{2} - p \binom{|Y \setminus X|}{2} + o(n^2) \\ &= p|X||Y| + o(n^2). \end{aligned}$$

- Disc to Count. This is just the **Graph Counting Lemma** (we only use the Disc part of regularity in the proof).
- Count to  $C_4$ -Count. This is clear from the definition.
- $C_4$ -Count to Codegree. Now,

$$\begin{aligned} \sum_{x, y \in V} d(x, y) &= \sum_{x \in V} d(x)^2 \\ &\geq \frac{1}{n} \left( \sum_{x \in V} d(x) \right)^2 \\ &= \frac{4e(G)^2}{n} = (1 + o(1))p^2n^3. \end{aligned}$$

Also note that

$$\sum_{x, y \in V} d(x, y)^2 = \text{number of labelled } C_4\text{s} + o(n^4) = (1 + o(1))p^4n^4.$$

So,

$$\begin{aligned} \sum_{x, y \in V} |d(x, y) - p^2n| &\leq n \left( \sum_{x, y \in V} (d(x, y) - p^2n)^2 \right)^{1/2} \\ &= n \left( \sum_{x, y \in V} d(x, y)^2 - 2p^2n \sum_{x, y \in V} d(x, y) + p^4n^2 \right)^{1/2} \\ &\leq n \left( (1 + o(1))p^4n^4 - 2p^2n(1 + o(1))p^2n^3 + p^4n^2 \right)^{1/2} \\ &= n(o(1)n^4)^{1/2} \\ &= o(n^3). \end{aligned}$$

- Codegree to Disc. We have

$$\begin{aligned}
\sum_{x \in V} |d(x) - pn| &\leq \sqrt{n} \left( \sum_{x \in V} (d(x) - pn)^2 \right)^{1/2} \\
&= \sqrt{n} \left( \sum_{x \in V} d^2(x) - 2pn \sum_{x \in V} d(x) + p^2 n^3 \right) \\
&= \sqrt{n} \left( \sum_{x, y \in V} d(x, y) - 2p^2 n^3 + p^2 n^3 o(n) \right) \\
&= \sqrt{n} \left( \sum_{x, y \in V} (d(x, y) - p^2 n) \right) \\
&\leq \sqrt{n} \left( \sum_{x, y \in V} |d(x, y) - p^2 n| \right) = o(n^2).
\end{aligned}$$

For  $X, Y \subseteq V$ ,

$$\begin{aligned}
|e(X, Y) - p|X||Y|| &\leq \sum_{x \in X} |d(x, Y) - p|Y|| \\
&\leq \sqrt{|X|} \left( \sum_{x \in X} (d(x, Y) - p|Y|)^2 \right)^{1/2} \\
&\leq \sqrt{|X|} \left( \sum_{v \in V} (d(v, Y) - p|Y|)^2 \right)^{1/2} \\
&= \sqrt{|X|} \left( \sum_{v \in V} d(v, Y)^2 - 2p|Y| \sum_{v \in V} d(v, Y) + p^2 |Y|^2 n \right)^{1/2} \\
&= \sqrt{|X|} \left( \sum_{y, y' \in Y} d(y, y')^2 - 2p|Y| \sum_{y \in Y} d(y) + p^2 |Y|^2 n \right)^{1/2} \\
&= \sqrt{|X|}
\end{aligned}$$

- Eigen to  $C_4$ -Count. Note that the number of labelled  $C_4$ s in  $G$  equals the number of closed walks of length 4 minus  $O(n^3)$ . Now, observe that given two vertices  $v, w$ , the number of walks from  $v$  to  $w$  of length 4 is just  $(A^4)_{vw}$  (Why?). In particular, the number of closed walks from  $v$  to  $v$  is  $(A^4)_{vv}$ . Recalling that the trace of a matrix is the sum of its eigenvalues,

$$\begin{aligned}
\text{number of } C_4\text{s} &= \text{Tr}(A^4) - O(n^3) \\
&\leq \lambda_1^4 + \sum_{i>1} \lambda_i^4 \\
&\leq (1 + o(1))(pn)^4 + \left( \max_{i>1} \lambda_i^2 \right) \sum_{i=1}^n \lambda_i^2 \\
&= (1 + o(1))(pn)^4 + o(n^2) \cdot 2e(G) \\
&= (1 + o(1))(pn)^4 + o(n^4) = (1 + o(1))p^4 n^4.
\end{aligned}$$

- $C_4$ -Count to Eigen. The maximum eigenvalue of  $A$  is just

$$\lambda_1 = \sup_{\|x\|_2=1} x^\top A x.$$

Taking  $x = (1/\sqrt{n})\mathbf{1}$ , we have

$$\lambda_1 \geq \frac{2e(G)}{n} = (p - o(1))n.$$

For the remaining eigenvalues,

$$\begin{aligned} \max_{i \neq 1} |\lambda_i|^4 &\leq \text{Tr}(A^4) - \lambda_1^4 \\ &\leq \underbrace{(p^4 + o(1))n^4}_{\text{from } C_4\text{-Count}} - (p - o(1))^4 n^4 \\ &= o(n^4), \end{aligned}$$

completing the proof. ■

*Remark.* • The proof requires  $p = \Omega(1)$ , in particular to show that Disc implies Count.

- Count does not hold in sparse regimes. In fact, Noga Alon in the 90s gave a graph with  $p = \Omega(n^{-1/3})$  which is Eigen-pseudorandom, but triangle-free.

### 2.5.2. Strongly regular graphs

Do we have any ‘known’ families of pseudorandom graphs?

**Definition 2.9.** Suppose  $q$  is a prime power such that  $q \equiv 1 \pmod{4}$ . Consider the **Paley graph**  $P_q$  with vertex set  $V(P_q) = \mathbb{F}_q$  and  $xy$  is an edge iff  $x - y$  is a quadratic residue<sup>9</sup>.

Let us see a couple of properties of the Paley graph.

**Definition 2.10.** A graph  $G$  is said to be an  $(n, d, \lambda)$ -**graph** if it is a regular graph of degree  $d$  over  $n$  vertices with  $\max_{i \neq 1} |\lambda_i| \leq \lambda$ , where  $\lambda_1 \geq \dots \geq \lambda_n$  are the eigenvalues of the graph.

**Definition 2.11.** A graph  $G = (V, E)$  is a  $(n, d, \lambda, \mu)$ -**strongly regular graph** if

- $|V| = n$ ,
- $G$  is regular with degree  $d$ ,
- Any two adjacent vertices have exactly  $\lambda$  common neighbours, and
- Any two non-adjacent vertices have exactly  $\mu$  common neighbours.

If a graph is  $(n, d, \lambda, \mu)$ -strongly regular for some  $n, d, \lambda, \mu$ , it is just said to be strongly regular.

---

<sup>9</sup>  $x - y = a^2$  for some  $a \in \mathbb{F}_q^\times$ .

For example, a complete graph on  $n$  vertices is  $(n, n-1, n-1, 0)$ -strongly regular and  $C_5$  is  $(5, 2, 0, 1)$ -strongly regular.

Slightly more complicatedly, the Petersen graph is  $(10, 3, 0, 1)$ -strongly regular.

Observe that a graph is  $(n, d, \lambda, \mu)$ -strongly regular iff its adjacency matrix  $A$  is such that

$$(A^2)_{vw} = \begin{cases} d, & i = j, \\ \lambda, & v, w \text{ are adjacent,} \\ \mu, & v, w \text{ are not adjacent.} \end{cases}$$

More succinctly,

$$A^2 = dI + \lambda A + \mu(J - I - A).$$

We see that

$$\begin{aligned} A^3 &= dA + \lambda A^2 + \mu dJ - \mu(A + A^2) \\ 0 &= A^3 + (\mu - \lambda)A^2 - dA - \mu dJ. \end{aligned}$$

The graph is strongly regular iff its adjacency matrix satisfies a cubic with integer coefficients and leading coefficient 1!

This further implies that a strongly regular graph has only 3 eigenvalues, with one of them being  $d$ .

Returning to the Paley graph, it is not too difficult to check that it is  $(q, (q-1)/2, (q-5)/4, (q-1)/4)$ -strongly regular. Consequently, it is a  $(q, (q-1)/2, (\sqrt{q}+1)/2)$ -graph.

Observe that any strongly regular graph has eigenvalues (other than  $\lambda_1$ ) bounded in a similar fashion.

### 2.5.3. The sparse regime

Let us restrict ourselves to  $(n, d, \lambda)$ -graphs. A popular family of  $(n, d, \lambda)$ -graphs comes from the following.

**Definition 2.12.** Suppose  $\Gamma$  is a finite group and  $S \subseteq \Gamma \setminus \{1\}$  is such that  $S = S^{-1}$ . We define the **Cayley graph**  $\text{Cay}(\Gamma, S)$  as that on vertex set  $\Gamma$  with  $gh \in E(G)$  iff  $g^{-1}h \in S$ .

It is easy to see that  $\text{Cay}(\Gamma, S)$  is  $|S|$ -regular. In fact,  $P_q$  is just a Cayley graph with  $\Gamma = \mathbb{F}_q$  (under addition) and  $S$  is the set of quadratic residues in  $\mathbb{F}_q$  other than 1.

**Theorem 2.25** (Conlon-Zhao, 2017). For Cayley graphs, Disc and Eigen are equivalent.

*Proof.* First, note that Eigen implies Disc with the same constant for both. That is, if

$$\max_{i \neq 1} |\lambda_i| \leq \varepsilon n.$$

then

$$|e(X, Y) - p|X||Y|| \leq \varepsilon n^2$$

by the **Expander Mixing Lemma**.

To show the other side, we require a result of Grothendieck's:

There exists an absolute constant  $K > 0$  such that for all real matrices  $A = (a_{ij})$  and a Hilbert space  $\mathcal{H}$ ,

$$\sup_{\substack{x_i, y_j \in \mathcal{H} \\ \|x\|, \|y\|=1}} \sum_{i,j} a_{ij} \langle x_i, y_j \rangle \leq K \sup_{\{u,v\} \in \{-1,1\}^n} \sum_{i,j} a_{ij} u_i v_j.$$



The best possible  $K$  is unknown, but we do know that  $K < 1.79$  is possible. We skip the proof of this. Suppose that Disc holds with a constant of  $\varepsilon$  and let  $A$  be the adjacency matrix of  $G = \text{Cay}(\Gamma, S)$ . Let  $u, v \in \{-1, 1\}^n$ . Denote by  $u^+$  and  $u^-$  the positive and negative parts of  $u$  respectively. Letting  $U^+ \subseteq V$  be the support of  $u^+$  (the subset of vertices with non-zero components) then using Grothendieck's lemma,

$$(u^\pm)^\top \left( A - \frac{d}{n} J \right) v^\pm = e(U^\pm, V^\pm) - \frac{d}{n} |U^\pm| |V^\pm|,$$

where the sign over the  $U$ s (resp.  $V$ s) are the same as that over  $u$  (resp.  $v$ ). Using Disc,

$$u^\top \left( A - \frac{d}{n} J \right) v \leq 4(\varepsilon d)n.$$

Now, since

$$\max_{i \neq 1} |\lambda_i| = \sup_{\|x\|=\|y\|=1} x^\top \left( A - \frac{d}{n} J \right) y,$$

it suffices to show that the quantity on the right is 'small'.

Let  $B = A - (d/n)J$ . We have

$$\begin{aligned} x^\top B y &= \sum_{g,h \in \Gamma} x_g B_{g,h} y_h \\ &= \frac{1}{n} \sum_{s,g,h \in \Gamma} x_{sg} B_{sg,sh} y_{sh} \\ &= \frac{1}{n} \sum_{s,g,h \in \Gamma} x_{sg} B_{g,h} y_{sh} && (\Gamma \text{ is an automorphism group of } G) \\ &= \frac{1}{n} \sum_{g,h \in \Gamma} \left( B_{g,h} \sum_{s \in \Gamma} x_{sg} y_{sh} \right) \\ &= \frac{1}{n} \sum_{g,h \in \Gamma} B_{g,h} \langle x^{(g)}, y^{(h)} \rangle && \text{where } x_k^{(g)} = x_{kg} \\ &\leq \frac{1}{n} \sup_{\|x\|=\|y\|=1} \sum_{g,h \in \Gamma} B_{g,h} \langle x^{(g)}, y^{(h)} \rangle && (\|x^{(g)}\| = \|x\| = 1 \text{ and } \|y^{(h)}\| = \|y\| = 1) \\ &\leq \frac{1}{n} \cdot K \cdot (4\varepsilon d n) && (\text{using Grothendieck's result and the above result}) \\ &< 8\varepsilon d, \end{aligned}$$

using the fact that  $\|x^{(g)}\|^2 = \|x\|^2$ , completing the proof. ■

**Theorem 2.26.** If  $G$  is  $d$ -regular, then  $\lambda = \max\{|\lambda_2(G)|, |\lambda_n(G)|\}$  is at least  $\sqrt{d}(1 - o_n(1))$ .

*Proof.* Let  $A$  be the adjacency matrix of  $G$ . Observe that

$$\begin{aligned} nd &= \text{Tr}(A^2) \\ &= \sum_i \lambda_i^2 \\ &\leq d^2 + (n-1)\lambda^2, \end{aligned}$$

so

$$\lambda \geq \sqrt{d}(1 - o_n(1)). \quad \blacksquare$$

However, we can get a better bound.

**Theorem 2.27** (Alon-Boppana bound). If  $G$  is a connected  $d$ -regular graph, then  $\lambda(G) = \max\{|\lambda_2(G)|, |\lambda_n(G)|\}$  is at least  $2\sqrt{d-1}(1 - o_n(1))$ .

*Proof.* To prove the result, we shall get a vector  $x \perp \mathbf{1}$  such that

$$\frac{x^\top Ax}{x^\top x} \geq 2\sqrt{d-1}(1 - o_n(1)).$$

Fix a vertex  $v$  and let  $r \in \mathbb{N}$  (which we shall fix later). For  $u \in V(G)$ , consider the vector  $x$  defined by

$$x_u = \begin{cases} (d-1)^{-i/2}, & d(u, v) = i \leq r-1, \\ 0, & \text{otherwise,} \end{cases}$$

where  $d(u, v)$  is the distance between the vertices  $u$  and  $v$ . Let  $V_i = \{u : d(u, v) = i\}$ . We have

$$x^\top x = \sum_{i=0}^{r-1} \frac{|V_i|}{(d-1)^i}$$

and

$$\begin{aligned} x^\top Ax &= \sum_{u \in V} x_u \sum_{w \in \Gamma(u)} x_w \\ &\geq \sum_{i=0}^{r-2} |V_i| \cdot \frac{1}{(d-1)^{i/2}} \left( \frac{1}{(d-1)^{(i-1)/2}} + \frac{(d-1)}{(d-1)^{(i+1)/2}} \right) + \frac{|V_{r-1}|}{(d-1)^{(r-1)/2}} \cdot \frac{1}{(d-1)^{r/2}} \\ &\quad \text{(if } u \in V_i, \text{ then } \Gamma(u) \subseteq V_{i-1} \cup V_{i+1}, \text{ and } u \text{ must have a neighbour in } V_{i-1}) \\ &\geq \sum_{i=0}^{r-2} |V_i| \cdot \frac{2\sqrt{d-1}}{(d-1)^i} + \frac{|V_{r-1}|(d-1)}{(d-1)^{(2r-1)/2}} \\ &\geq 2\sqrt{d-1} \left( \sum_{i=0}^{r-2} \frac{|V_i|}{(d-1)^i} + \frac{1}{2} \frac{|V_{r-1}|}{(d-1)^r} \right) \\ &\geq 2\sqrt{d-1} \left( 1 - \frac{1}{2r} \right) \sum_{i=0}^{r-2} \frac{|V_i|}{(d-1)^i}. \end{aligned} \quad (|V_i|/(d-1)^i \leq |V_{i-1}|/(d-1)^{i-1})$$

So,

$$\frac{x^\top Ax}{x^\top x} \geq 2\sqrt{d-1} \left( 1 - \frac{1}{2r} \right).$$

However,  $x$  is not orthogonal to  $\mathbf{1}$ , so this does not directly yield a bound on  $\lambda(G)$ .

Now note that if  $n > 1 + (d-1) + (d-1)^2 + \dots + (d-1)^{k-1}$ , then there must be two vertices that are at a distance of at least  $k$ . Equivalently,

$$n > \frac{(d-1)^k - 1}{d-2},$$

which gives a lower bound on  $k$  that is  $\Omega(\log n)$ .

Let  $v_1, v_2$  be two farthest vertices in  $G$ , and let  $r$  such that  $d(v_1, v_2) \geq 2r$ . Let the vector  $x_1$  be that as defined above with  $v = v_1$  with

$$\frac{x_1^\top Ax_1}{x_1^\top x_1} \geq 2\sqrt{d-1} \left( 1 - \frac{1}{2r} \right).$$

Similarly get  $x_2$  using  $v_2$ . Note that  $x_1$  and  $x_2$  must be orthogonal. Let  $c$  be a constant such that  $y = (x_1 + cx_2)$  is orthogonal to  $\mathbf{1}$  (Why does such a  $c$  exist?). We then have

$$y^\top y = \|x_1\|^2 + c^2 \|x_2\|^2$$

and

$$y^\top Ay = x_1^\top Ax_1 + c^2 x_2^\top Ax_2 \geq 2\sqrt{d-1} \left(1 - \frac{1}{2r}\right) (\|x_1\|^2 + c^2 \|x_2\|^2),$$

completing the proof. ■

This leads to the following definition, which we shall not get into.

**Definition 2.13.** An  $(n, d, \lambda)$ -graph is called a **Ramanujan graph** if  $\lambda \leq 2\sqrt{d-1}$ .

This notion is due to Lubotzky, Phillips, and Sarnak, who constructed what are called “LPS graphs”, that are Cayley graphs on the group  $\text{PSL}(2, \mathbb{F}_q)$ , that are  $(q+1)$ -regular and Ramanujan.

Alon conjectured (mayhaps surprisingly) that random regular graphs are Ramanujan with high probability. This was resolved by Joel Friedman in the early 2000s.

Questions on the existence of Ramanujan graphs (and explicit constructions) are largely open, and an active area of research in combinatorics and theoretical computer science.

For  $\alpha > 0$ , one can also define  $(n, d, \alpha)$ -expanders, which are  $(n, d, \lambda)$ -graphs with  $\lambda \leq \alpha d$ .

### §3. Matching Theory

**Definition 3.1.** Given a graph  $G$ , a **matching**  $\mathcal{M}$  in  $G$  is a collection of pairwise disjoint edges in  $G$  (no two of the edges have a common vertex).

For example,  $\{\{1, 6\}, \{2, 5\}, \{3, 4\}\}$  is a matching in  $K_6$ .

**Definition 3.2.** Given a graph  $G$ , the **matching number**  $\nu(G)$  of  $G$  is the size of a maximum matching in  $G$ .

Given a graph, how do we compute a maximum matching, or simpler yet,  $\nu(G)$ ?

#### 3.1. The Bipartite Setting

For this section, suppose that  $G = (X, Y, E)$  is bipartite, where all edges in  $E$  are between  $X$  and  $Y$ . Assume  $|X| \leq |Y|$ .

It is clear that  $\nu(G) \leq |X|$ .

The question is: under what conditions on  $G$  is  $\nu(G) = |X|$ ?

This (the bipartite scenario) is sometimes referred to as a *system of distinct representatives* (SDR). That is,  $G$  has an SDR if  $\nu(G) = |X|$ .

##### 3.1.1. Hall's Marriage Theorem

An obvious necessary condition is that given any  $S \subseteq X$ ,

$$|\Gamma(S)| \geq |S|.$$

Otherwise, there cannot exist a matching since two vertices in  $S$  would be forced to map to the same element in  $Y$ . This is known as *Hall's condition*.

It in fact turns out that Hall's condition is sufficient for the existence of a matching too!

**Theorem 3.1** (Hall's Marriage Theorem). Suppose  $G = (X, Y, E)$  is bipartite with  $|X| \leq |Y|$ . Then  $\nu(G) = |X|$  if and only if it satisfies Hall's condition.

Much like Turán's theorem, this has a large number of proofs.

*Proof 1 of Hall's Marriage Theorem.* We begin with a proof by induction over  $|X|$ . If  $|X| = 1$ , the theorem obviously holds.

Make the stronger assumption that for all  $S \subseteq X$  of size  $|X| - 1$ ,  $|\Gamma(S)| > |S|$ , and  $|\Gamma(X)| \geq |X|$ . Pick  $x \in X$  and pair it with an arbitrary neighbour  $y \in Y$ . Using the inductive hypothesis on the subgraph induced on  $(X \setminus \{x\}) \cup (Y \setminus \{y\})$ , we get a matching on  $G$ .

Now, let  $S \subseteq X$  with  $|\Gamma(S)| = |S| = |X| - 1$ . Using the inductive hypothesis on the subgraph induced on  $S \cup \Gamma(S)$ , together with the edge from the (single) element in  $X \setminus S$  to any element in  $\Gamma(X) \setminus \Gamma(S)$ , we get a matching on  $G$ . ■

An alternate way to prove this is by doing the first part of the proof with all non-empty  $S \subseteq X$  (instead of only those of size  $|X| - 1$ ). In the case where we have some non-empty  $S \subseteq X$  with  $|\Gamma(S)| = |S|$ , we can use the inductive hypothesis twice to get two matchings  $\mathcal{M}_1$  and  $\mathcal{M}_2$  on the subgraphs induced by  $S \cup \Gamma(S)$  and  $(X \setminus S) \cup (Y \setminus \Gamma(S))$ , then take the union of the two matchings to get a matching  $\mathcal{M}$  on the original graph  $G$ .

A nice consequence of the above proof is that it allows us to extend Hall's Theorem to an infinite setting as well.

**Definition 3.3.** Given a matching  $\mathcal{M}$  on a graph  $G = (V, E)$ , call a  $v \in V$  **unsaturated** if no edge of  $\mathcal{M}$  is incident on  $v$ .

For a matching  $\mathcal{M}$ ,  $x \in X$ ,  $y \in Y$  such that  $x, y$  are unsaturated (with respect to  $\mathcal{M}$ ), a path  $P$  from  $x$  to  $y$  is called  **$\mathcal{M}$ -augmenting** if every alternate edge in it is a  $\mathcal{M}$ -edge.

Observe that if  $\mathcal{M}$  admits an augmenting path, then  $\mathcal{M}$  is not maximal. Indeed, we can 'flip' the edges in the path to show this. That is, given an augmenting path  $\mathcal{P}$ , where every  $y_i x_i$  is in  $\mathcal{M}$ , it is not too difficult to show that  $\mathcal{M} \triangle \mathcal{P}$  is a matching that has strictly more edges.

*Proof 2 of Hall's Marriage Theorem.* Let  $\mathcal{M}$  be a maximum matching of a bipartite graph  $G = (X, Y, E)$  that satisfies Hall's condition. Suppose there exists  $x_0 \in X$  such that  $x_0$  is unsaturated by  $\mathcal{M}$ . Let  $y_1$  be a neighbour of  $x_0$  (such a  $y_1$  exists by Hall's condition).

If  $y_1$  is unsaturated by  $\mathcal{M}$ , then  $x_0 y_1$  is a  $\mathcal{M}$ -augmenting path, contradicting its maximality. So, suppose  $x_1 y_1 \in \mathcal{M}$ .

In general, if we have  $y_1, \dots, y_r, x_1, \dots, x_r$ , we can pick  $y_{r+1} \in \Gamma(\{x_0, \dots, x_r\})$  (distinct from all the  $y_i$  for  $1 \leq i \leq r$ ).

If  $y_{r+1}$  is unsaturated, we have a  $\mathcal{M}$ -augmenting path from  $x_0$  to  $y_{r+1}$ , contradicting the maximality of  $\mathcal{M}$ .

If it is not, we can jump back to the  $x_{r+1} \in X$  (distinct from the  $x_i$  for  $1 \leq i \leq r$ ) such that  $y_{r+1} x_{r+1} \in E$ , then continue the process. This must terminate at some point since the number of vertices in  $Y$  incident on some element of  $\mathcal{M}$  is at most  $|X \setminus \{x_0\}| < |Y|$ , completing the proof. ■

### 3.1.2. Some applications

Next, let us look at a few applications of Hall's Theorem.

**Definition 3.4 (Perfect Matching).** A matching  $\mathcal{M}$  in a graph  $G = (V, E)$  is said to be a **perfect matching** if  $|\mathcal{M}| = |V|/2$ , that is, every vertex in  $V$  is saturated by  $\mathcal{M}$ .

**Theorem 3.2.** If a bipartite graph  $G = (X, Y, E)$  is  $d(> 0)$ -regular, it has a perfect matching.

*Proof.* It suffices to check that Hall's condition holds. Let  $S \subseteq X$ . We clearly have  $e(S, \Gamma(S)) = d|S|$ . However, the total number of edges coming into  $\Gamma(S)$  must include all the edges coming from  $S$ , that is,  $S \subseteq \Gamma(\Gamma(S))$ . So,

$$\begin{aligned} d|\Gamma(S)| &= e(\Gamma(S), \Gamma(\Gamma(S))) \\ &\geq e(\Gamma(S), S) \\ &= d|S|, \end{aligned}$$

completing the proof. ■

For the next result, consider the following definition.

**Definition 3.5.** A matrix  $A \in \mathbb{R}^{n \times n}$  is said to be **doubly stochastic**<sup>10</sup> if for all  $i, j$ ,

$$0 \leq a_{ij} \leq 1 \text{ and } \sum_k a_{ik} = \sum_k a_{kj} = 1.$$

It is not too difficult to see that any permutation matrix is doubly stochastic.

Suppose  $A_1, A_2$  are doubly stochastic and  $0 \leq \lambda \leq 1$ . Then observe that  $\lambda A_1 + (1 - \lambda)A_2$  is doubly stochastic too (Why?).

As a result, a convex combination of doubly stochastic matrices is doubly stochastic.

**Theorem 3.3** (Birkhoff-von Neumann Theorem). A matrix is doubly stochastic iff it is a convex combination of permutation matrices.

## References

- [Erd59] P. Erdős. Graph theory and probability. *Canadian Journal of Mathematics*, 11:34–38, 1959.
- [Gow97] W. T. Gowers. Lower bounds of tower type for Szemerédi’s uniformity lemma. *Geometric & Functional Analysis GAFA*, 7(2):322–337, May 1997.
- [KRS96] János Kollár, Lajos Rónyai, and Tibor Szabó. Norm-graphs and bipartite Turán numbers. *Combinatorica*, 16(3):399–406, 1996.

---

<sup>10</sup>These are interesting to study since they arise as the transition matrices of a certain class of discrete Markov chains. The stationary distribution of a discrete Markov chain with a doubly stochastic transition matrix is the uniform distribution.