

# Linear Algebra Applications - Discrete Linear Dynamical Systems

Alexander Xie<sup>1\*</sup>

[alex.xie1@student.unsw.edu.au](mailto:alex.xie1@student.unsw.edu.au)

February 2025

## Abstract

Dynamical systems are processes that evolve in time. Through forms of mathematical modelling, the study of dynamical systems is beneficial to comprehend a variety of natural phenomena. In particular, discrete linear dynamical systems are useful to model a scenario as a series of state snapshots. In this paper, we focus on discrete linear dynamical systems in the form of first-order homogenous matrix recurrence relation. We then illustrate the application of these systems through a real-world case study by modelling the spread of a virus.

## Contents

<b>1</b>	<b>Discrete Linear Dynamical Systems</b>	<b>2</b>
1.1	A matrix recurrence relation . . . . .	2
1.2	Diagonalisation . . . . .	2
<b>2</b>	<b>More Examples</b>	<b>4</b>
<b>A</b>	<b>Omitted Proof in Section 2</b>	<b>4</b>

---

\*With support from Pau Battle

# 1 Discrete Linear Dynamical Systems

We will begin by exploring the theory central to our case study in Section 2.

## 1.1 A matrix recurrence relation

For a square matrix  $A$  and an initial state vector  $x_0$ , define the first-order homogenous recurrence:

$$x_t = Ax_{t-1}, \quad \text{for } t \geq 1.$$

Iterating yields a sequence of vectors  $x_0, Ax_0, A^2x_0, \dots, A^kx_0, \dots$  which represent the evolution of the system starting from  $x_0$ . This sequence is called the **orbit** of the system starting from  $x_0$ .

An important question to raise about this dynamical system is: what is its long-term behaviour? For example, if our system models a particular population of rabbits, it would be worthwhile to observe the population a few months ahead. What happens as  $t \rightarrow \infty$ ? Does the system have limiting behaviours?

To compute the elements in our orbit, we require an efficient method to compute  $A^n$  for large  $n$ , as repeated matrix multiplication is cumbersome.

## 1.2 Diagonalisation

Recall that it is easy to multiply diagonal matrices (square matrices in which the entries outside the main diagonal are all zero):

$$\begin{pmatrix} a_{11} & & & \\ & a_{22} & & \\ & & \ddots & \\ & & & a_{nn} \end{pmatrix} \begin{pmatrix} b_{11} & & & \\ & b_{22} & & \\ & & \ddots & \\ & & & b_{nn} \end{pmatrix} = \begin{pmatrix} a_{11}b_{11} & & & \\ & a_{22}b_{22} & & \\ & & \ddots & \\ & & & a_{nn}b_{nn} \end{pmatrix}$$

This simplicity of multiplying diagonal matrices prompts us to investigate if  $A$  is similar to a diagonal matrix; do there exist an invertible matrix  $P$  and a diagonal matrix  $D$  such that  $A$  can be written as  $PDP^{-1}$ ? If so, then we say that  $A$  is **diagonalisable**. This condition is equivalent to  $AP = PD$ . Observe that it is easier to compute powers of a diagonalisable matrix:

$$\begin{aligned} A^n &= (PDP^{-1})(PDP^{-1}) \dots (PDP^{-1}) \\ &= PD \underbrace{(P^{-1}P)}_I D \underbrace{(P^{-1}P)}_I D \dots \underbrace{(P^{-1}P)}_I DP^{-1} \\ &= PD^n P^{-1} \end{aligned}$$

**Theorem 1.1.** *A square matrix is diagonalisable if and only if it has  $n$  linearly independent eigenvectors.*

*Proof.* ( $\implies$ ) Suppose that  $A \in \mathbb{R}^{n \times n}$  is diagonalisable. Then there exists an invertible matrix  $P \in \mathbb{R}^{n \times n}$  and a diagonal matrix  $D \in \mathbb{R}^{n \times n}$  such that  $AP = PD$ .

Let  $P = (v_1 \ v_2 \ \dots \ v_n)$  where  $v_i$  are linearly independent columns, and let  $D = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$ .

Writing  $AP = PD$  as columns, we have  $(Av_1 \ Av_2 \ \dots \ Av_n) = (\lambda_1 v_1 \ \lambda_2 v_2 \ \dots \ \lambda_n v_n)$ .

Thus,  $Av_i = \lambda_i v_i$  for each  $i = 1, \dots, n$ . Hence,  $v_i$  is an eigenvector associated to  $\lambda_i$ .

( $\impliedby$ ) Conversely, suppose that  $A \in \mathbb{R}^{n \times n}$  has  $n$  linearly independent eigenvectors  $v_1, v_2, \dots, v_n$  associated to the eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$  respectively. Then for each  $i = 1, \dots, n$  we have  $Av_i = \lambda_i v_i$ .

Let  $P = (v_1 \ v_2 \ \dots \ v_n)$  where  $v_i$  are the columns. Note that  $P$  is invertible as the columns are linearly independent. Also, let  $D = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$ .

Now the equations  $Av_i = \lambda_i v_i$  for each  $i = 1, \dots, n$  can be compactly expressed as  $AP = PD$ , which gives  $A = PDP^{-1}$ . Hence,  $A$  is diagonalisable.  $\square$

**Corollary 1.2.** *An eigen-formula for diagonalisable matrices.*

In 1.1, observe that  $P$  and  $D$  are constructed by the eigenvectors and eigenvalues of  $A$ . So if  $A \in \mathbb{R}^{n \times n}$  has  $n$  linearly independent eigenvectors  $v_1, v_2, \dots, v_n$  associated to the eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ , then

$$A = (v_1 \ v_2 \ \cdots \ v_n) \begin{pmatrix} \lambda_1 & & \\ & \lambda_2 & \\ & & \ddots \\ & & & \lambda_n \end{pmatrix} (v_1 \ v_2 \ \cdots \ v_n)^{-1}$$

**Example 1.3.** Diagonalise  $A = \begin{pmatrix} 2 & 0 & 0 \\ -2 & 1 & 1 \\ 1 & 0 & -1 \end{pmatrix}$ , if possible.

We will first find the eigenvalues of  $A$  by solving  $\det(\lambda I - A) = 0$ .

$$\begin{aligned} 0 &= \det \begin{pmatrix} \lambda - 2 & 0 & 0 \\ 2 & \lambda - 1 & -1 \\ -1 & 0 & \lambda + 1 \end{pmatrix} \\ &= (\lambda - 2) \det \begin{pmatrix} \lambda - 1 & -1 \\ 0 & \lambda + 1 \end{pmatrix} \\ &= (\lambda - 2)(\lambda - 1)(\lambda + 1) \end{aligned}$$

Thus the eigenvalues of  $A$  are  $\lambda_1 = 1, \lambda_2 = 2, \lambda_3 = -1$ .

To find the eigenvectors of  $A$ , we solve  $(\lambda_i I - A)v_i = 0$  for each  $i = 1, 2, 3$ .

By row reducing each equation, it can be shown that we have  $v_1 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ ,  $v_2 = \begin{pmatrix} 3 \\ -5 \\ 1 \end{pmatrix}$  and  $v_3 = \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix}$ .

The eigenvectors of  $A$  are clearly linearly independent, thus by 1.1,  $A$  is diagonalisable, and by 1.2, we can write  $A$  as  $PDP^{-1}$  where  $P = \begin{pmatrix} 0 & 3 & 0 \\ 1 & -5 & -1 \\ 0 & 1 & 2 \end{pmatrix}$  and  $D = \begin{pmatrix} 1 & & \\ & 2 & \\ & & -1 \end{pmatrix}$ .

Unordered List (taken from Overleaf)

- The individual entries are indicated with a black dot, a so-called bullet.
- The text in the entries may be of any length.

Ordered List (taken from Overleaf)

1. The labels consists of sequential numbers.
2. The numbers starts at 1 with every call to the enumerate environment.

Odd	Even
One	Two
Three	Four

Table 1: This is a table

Table 1 is an example of a table.

## 2 More Examples

Now we include a figure. (See Figure 1.)




Figure 1: An example of a figure

**Acknowledgements** Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

## A Omitted Proof in Section 2

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.