

## Supplemental Material

Alexander Lalejini, Austin J. Ferguson, Nkrumah Grant, and Charles Ofria

2021-02-18



# Contents

<b>1</b>	<b>Introduction</b>	<b>5</b>
<b>2</b>	<b>Validation experiment</b>	<b>9</b>
2.1	Overview . . . . .	9
2.2	Analysis dependencies . . . . .	10
2.3	Setup . . . . .	10
2.4	Evolution of phenotypic plasticity . . . . .	12
<b>3</b>	<b>Evolutionary change</b>	<b>15</b>
3.1	Overview . . . . .	15
3.2	Analysis dependencies . . . . .	15
3.3	Setup . . . . .	16
3.4	Evolution of phenotypic plasticity . . . . .	17
3.5	Average generation . . . . .	19
3.6	Selective sweeps . . . . .	21
3.7	Phenotypic volatility along dominant lineage . . . . .	26
3.8	Mutation accumulation along the dominant lineage . . . . .	34
3.9	Depth of MRCA . . . . .	51
3.10	Manuscript figures . . . . .	52
<b>4</b>	<b>Evolution and maintenance of novel traits</b>	<b>59</b>
4.1	Overview . . . . .	59
4.2	Analysis dependencies . . . . .	60
4.3	Setup . . . . .	61
4.4	Evolution of phenotypic plasticity . . . . .	63
4.5	Final dominant novel task performance . . . . .	64
4.6	Final population novel task performance . . . . .	67
4.7	Population-level novel tasks discovered . . . . .	69
4.8	Novel tasks along lineage of final dominant genotype . . . . .	72
4.9	Extra task performance over time . . . . .	91
4.10	Manuscript figures . . . . .	93
<b>5</b>	<b>Genetic hitchhiking</b>	<b>99</b>
5.1	Overview . . . . .	99

5.2	Analysis dependencies . . . . .	99
5.3	Setup . . . . .	100
5.4	Evolution of phenotypic plasticity . . . . .	102
5.5	Hitchhiking instruction execution . . . . .	104
5.6	Characterizing mutations that increase hitchhiker instruction execution . . . . .	115
5.7	Manuscript figures . . . . .	128

# Chapter 1

## Introduction




Figure 1.1: Experimental design overview




Figure 1.2: Fluctuating environment




Figure 1.3: Static environment





## Chapter 2

# Validation experiment

In this experiment, we validate that (1) we observe the evolution of phenotypic plasticity in a changing environment when digital organisms have access to sensory instructions (capable of differentiating environmental states) and (2) that adaptive phenotypic plasticity does not evolve when populations lack access to sensory instructions.

### 2.1 Overview

```
total_updates <- 200000
replicates <- 100

all_traits <- c("not","nand","and","ornot","or","andnot")
traits_set_a <- c("not", "and", "or")
traits_set_b <- c("nand", "ornot", "andnot")

# Relative location of data.
working_directory <- "experiments/2021-01-07-validation/analysis/" # << For bookdown
# working_directory <- "./" # << For local analysis
```

We evolved populations of digital organisms under four conditions:

1. A fluctuating environment with access to sensory instructions
2. A fluctuating environment without access to sensory instructions (i.e., sensory instructions are no-operations)
3. A constant environment with access to sensory instructions
4. A constant environment without access to sensory instructions

In fluctuating environments, we alternate between rewarding and punishing different sets of computational tasks. In one environment, we reward tasks not,

and, or and punish tasks nand, ornot, andnot. In the alternative environment, we reward tasks nand, ornot, andnot and punish tasks not, and, or. In constant environments, we reward all tasks (not, nand, and, ornot, or, andnot).

For each replicate of each condition, we extract the dominant (i.e., most numerous) genotype at the end of the run to analyze further. We expect to observe the evolution of adaptive phenotypic plasticity in only the first experimental condition. In conditions without sensors, plasticity in any form should be unable to evolve.

## 2.2 Analysis dependencies

Load all required R libraries.

```
library(ggplot2)
library(tidyverse)
library(cowplot)
source("https://gist.githubusercontent.com/benmarwick/2a1bb0133ff568cbe28d/raw/fb53bd9")
```

These analyses were conducted/knitted with the following computing environment:

```
print(version)

##
## platform      x86_64-pc-linux-gnu
## arch          x86_64
## os            linux-gnu
## system        x86_64, linux-gnu
## status
## major         4
## minor         0.4
## year          2021
## month         02
## day           15
## svn rev       80002
## language      R
## version.string R version 4.0.4 (2021-02-15)
## nickname      Lost Library Book
```

## 2.3 Setup

```
data_loc <- paste0(working_directory, "data/aggregate.csv")
data <- read.csv(data_loc, na.strings="NONE")

data$DISABLE_REACTION_SENSORS <- as.factor(data$DISABLE_REACTION_SENSORS)
```

```

data$chg_env <- as.factor(data$chg_env)
data$dom_plastic_odd_even <- as.factor(data$dom_plastic_odd_even)
data$sensors <- data$DISABLE_REACTION_SENSORS == "0"
data$is_plastic <- data$dom_plastic_odd_even == "True"

env_label_fun <- function(chg_env) {
  if (chg_env) {
    return("Fluctuating")
  } else {
    return("Constant")
  }
}

sensors_label_fun <- function(has_sensors) {
  if (has_sensors) {
    return("Sensors")
  } else {
    return("No sensors")
  }
}

# Count observed plasticity for each condition (I'm sure there's a 'tidier' way to do this..)
observed_plasticity <- data.frame(
  environment=character(),
  sensors=character(),
  plastic=integer(),
  nonplastic=integer(),
  plastic_adaptive=integer(),
  plastic_optimal=integer(),
  plastic_nonadaptive=integer()
)

for (env_chg in levels(data$chg_env)) {
  for (disabled_sensors in levels(data$DISABLE_REACTION_SENSORS)) {
    cond_data <- filter(data, chg_env == env_chg & data$DISABLE_REACTION_SENSORS == disabled_sensors)
    environment_label <- env_label_fun(env_chg)
    sensors_label <- sensors_label_fun(disabled_sensors == "0")

    observed_plasticity <- observed_plasticity %>% add_row(
      environment=environment_label,
      sensors=sensors_label,
      plastic=nrow(filter(cond_data, is_plastic==TRUE)),
      nonplastic=nrow(filter(cond_data, is_plastic==FALSE)),
      plastic_adaptive=nrow(filter(cond_data, dom_adaptive_plasticity=="True")),
      plastic_optimal=nrow(filter(cond_data, dom_optimal_plastic=="True")),
      plastic_nonadaptive=nrow(filter(cond_data, is_plastic==TRUE & dom_adaptive_plasticity=="False"))
    )
  }
}

```

```

    )
  }
}

observed_plasticity <- pivot_longer(
  observed_plasticity,
  cols=c("plastic", "plastic_adaptive", "plastic_optimal", "plastic_nonadaptive", "nonplastic"),
  names_to="phenotype",
  values_to="phenotype_cnt"
)

##### misc #####
# Configure our default graphing theme
theme_set(theme_cowplot())

```

## 2.4 Evolution of phenotypic plasticity

For each experimental condition, do we observe the evolution of phenotypic plasticity? To test for phenotypic plasticity, we culture digital organisms in both environments from the fluctuating condition (including organisms evolved in a constant environment). Any plasticity that we observe from digital organisms evolved under constant conditions is cryptic variation (as these organisms were never exposed to these culturing environments).

```

ggplot(filter(observed_plasticity, phenotype %in% c("plastic", "nonplastic")), aes(x=phenotype, y=phenotype_cnt)) +
  geom_bar(
    stat="identity",
    position=position_dodge(0.9)
  ) +
  geom_text(
    stat="identity",
    mapping=aes(label=phenotype_cnt),
    vjust=0.05
  ) +
  scale_fill_brewer(palette="Accent") +
  scale_x_discrete(
    name="Phenotype",
    limits=c("plastic", "nonplastic"),
    labels=c("Plastic", "Non-plastic")
  ) +
  facet_grid(sensors~environment) +
  theme(
    legend.position="none"
  )


```



Indeed, we do not observe the evolution of phenotypic plasticity in any replicates in which digital organisms do not have access to sensory instructions. We do observe the evolution of plasticity (not necessarily adaptive plasticity) in both constant and fluctuating environments where sensors are enabled.

To what extent is the observed phenotypic plasticity adaptive?

```
ggplot(filter(observed_plasticity, environment=="Fluctuating" & sensors == "Sensors" & phenotype
  geom_bar(
    stat="identity",
    position=position_dodge(0.9)
  ) +
  geom_text(
    stat="identity",
    mapping=aes(label=phenotype_cnt),
    vjust=0.05
  ) +
  scale_fill_brewer(palette="Accent") +
  scale_x_discrete(
    name="Phenotype",
    limits=c("plastic", "plastic_adaptive", "plastic_optimal", "plastic_nonadaptive"),
    labels=c("Total plastic", "Adaptive plasticity", "Optimal plasticity", "Non-adaptive plasticity")
  ) +
  facet_grid(sensors~environment) +
  theme(
    legend.position="none"
```



## Chapter 3

# Evolutionary change

The effect of adaptive phenotypic plasticity on evolutionary change.

### 3.1 Overview

```
total_updates <- 200000
replicates <- 100

all_traits <- c("not","nand","and","ornot","or","andnot")
traits_set_a <- c("not", "and", "or")
traits_set_b <- c("nand", "ornot", "andnot")

# Relative location of data.
working_directory <- "experiments/2021-02-08-evo-dynamics/analysis/" # << For bookdown
# working_directory <- "./" # << For local analysis
```

### 3.2 Analysis dependencies

Load all required R libraries.

```
library(ggplot2)
library(tidyverse)
library(cowplot)
library(RColorBrewer)
library(Hmisc)
library(boot)
source("https://gist.githubusercontent.com/benmarwick/2a1bb0133ff568cbe28d/raw/fb53bd97121f7f9ce9")
```

These analyses were conducted/knitted with the following computing environ-

ment:

```
print(version)

##
## platform      x86_64-pc-linux-gnu
## arch          x86_64
## os            linux-gnu
## system        x86_64, linux-gnu
## status
## major         4
## minor         0.4
## year          2021
## month         02
## day           15
## svn rev       80002
## language      R
## version.string R version 4.0.4 (2021-02-15)
## nickname      Lost Library Book
```

### 3.3 Setup

```
summary_data_loc <- paste0(working_directory, "data/aggregate.csv")
summary_data <- read.csv(summary_data_loc, na.strings="NONE")

summary_data$DISABLE_REACTION_SENSORS <- as.factor(summary_data$DISABLE_REACTION_SENSORS)
summary_data$chg_env <- summary_data$chg_env == "True"
summary_data$dominant_plastic_odd_even <- as.factor(summary_data$dominant_plastic_odd_even)
summary_data$sensors <- summary_data$DISABLE_REACTION_SENSORS == "0"
summary_data$is_plastic <- summary_data$dominant_plastic_odd_even == "True"

env_label_fun <- function(chg_env) {
  if (chg_env) {
    return("Fluctuating")
  } else {
    return("Constant")
  }
}

sensors_label_fun <- function(has_sensors) {
  if (has_sensors) {
    return("Sensors")
  } else {
    return("No sensors")
  }
}
```



```

}

# note that this labeler makes assumptions about how we set up our experiment
condition_label_fun <- function(has_sensors, env_chg) {
  if (has_sensors && env_chg) {
    return("PLASTIC")
  } else if (env_chg) {
    return("NON-PLASTIC")
  } else {
    return("STATIC")
  }
}

summary_data$env_label <- mapply(
  env_label_fun,
  summary_data$chg_env
)
summary_data$sensors_label <- mapply(
  sensors_label_fun,
  summary_data$sensors
)
summary_data$condition <- mapply(
  condition_label_fun,
  summary_data$sensors,
  summary_data$chg_env
)

condition_order = c(
  "STATIC",
  "NON-PLASTIC",
  "PLASTIC"
)

##### misc #####
# Configure our default graphing theme
theme_set(theme_cowplot())
dir.create(paste0(working_directory, "plots"), showWarnings=FALSE)

```

### 3.4 Evolution of phenotypic plasticity


For sensor-enabled populations in fluctuating environments, we only transferred populations containing an optimally plastic genotype to phase-two.

```

summary_data_grouped = dplyr::group_by(summary_data, condition)
summary_data_group_counts = dplyr::summarize(summary_data_grouped, n=dplyr::n())

```


```
ggplot(summary_data_group_counts, aes(x=condition, y=n, fill=condition)) +
  geom_col(position=position_dodge(0.9)) +
  geom_text(aes(label=n, y=n+2)) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylab("Number of replicates in phase two") +
  theme(
    legend.position="none"
  )
)
```



We can confirm our expectation that the dominant genotypes in non-plastic conditions are not phenotypically plastic.

```
summary_data_grouped = dplyr::group_by(summary_data, condition, is_plastic)
summary_data_group_counts = dplyr::summarize(summary_data_grouped, n=dplyr::n())
ggplot(filter(summary_data_group_counts, is_plastic), aes(x=condition, y=n, fill=condition)) +
  geom_col(
    position=position_dodge(0.9)
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  )
```

```
geom_text(aes(label=n, y=n+1)) +
ylab("Number of plastic replicates") +
ylim(0, 100) +
theme(
  legend.position="none"
)
```




### 3.5 Average generation

```
ggplot(summary_data, aes(x=condition, y=time_average_generation, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
```

```

alpha = 0.5
) +
scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
ylab("average generation") +
theme(
  legend.position="none"
)

```



```

paste0(
  "PLASTIC median: ",
  median(filter(summary_data, condition=="PLASTIC"))$time_average_generation)
)

```

```
## [1] "PLASTIC median: 31697.65"
```

```

paste0(
  "STATIC median: ",
  median(filter(summary_data, condition=="STATIC"))$time_average_generation)
)

```

```
## [1] "STATIC median: 30839.75"
```

```

paste0(
  "NON-PLASTIC median: ",

```

```

    median(filter(summary_data, condition=="NON-PLASTIC")$time_average_generation)
)

## [1] "NON-PLASTIC median: 41768.65"
kruskal.test(
  formula=time_average_generation~condition,
  data=summary_data
)

##
## Kruskal-Wallis rank sum test
##
## data:  time_average_generation by condition
## Kruskal-Wallis chi-squared = 177.33, df = 2, p-value < 2.2e-16
pairwise.wilcox.test(
  x=summary_data$time_average_generation,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)

##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data:  summary_data$time_average_generation and summary_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.004
##
## P value adjustment method: bonferroni

```

## 3.6 Selective sweeps

The number of times the most recent common ancestor changes gives us the number of selective sweeps that occur during the experiment.

```


ggplot(summary_data, aes(x=condition, y=phylo_mrca_changes, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,

```

```

alpha = 0.8
) +
geom_boxplot(
  width = .1,
  outlier.shape = NA,
  alpha = 0.5
) +
scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
ylab("Number of selective sweeps") +
theme(
  legend.position="none"
)

```



```

paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC")$phylo_mrca_changes)
)

```

```
## [1] "PLASTIC: 45.5"
```

```

paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC")$phylo_mrca_changes)
)

```

```

)

## [1] "STATIC: 45"
paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$phylo_mrca_changes)
)

## [1] "NON-PLASTIC: 663.5"
kruskal.test(
  formula=phylo_mrca_changes~condition,
  data=summary_data
)

##
## Kruskal-Wallis rank sum test
##
## data: phylo_mrca_changes by condition
## Kruskal-Wallis chi-squared = 175.46, df = 2, p-value < 2.2e-16
pairwise.wilcox.test(
  x=summary_data$phylo_mrca_changes,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)

##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: summary_data$phylo_mrca_changes and summary_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      1
##
## P value adjustment method: bonferroni

```

### 3.6.1 Average number of generations between selective sweeps

```

summary_data$generations_per_mrca_change <- summary_data$time_average_generation / summary_data$phylo_mrca_change


ggplot(summary_data, aes(x=condition, y=generations_per_mrca_change, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),

```

```

alpha = .8
) +
geom_point(
  mapping=aes(color=condition),
  position = position_jitter(width = .15),
  size = .5,
  alpha = 0.8
) +
geom_boxplot(
  width = .1,
  outlier.shape = NA,
  alpha = 0.5
) +
scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
theme(
  legend.position="none"
)

```



```

paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC")$generations_per_mrca_change)
)

```



```
## [1] "PLASTIC: 685.001780758557"
```

```
paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC")$generations_per_mrca_change)
)
```

```
## [1] "STATIC: 693.676265008576"
```

```
paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$generations_per_mrca_change)
)
```

```
## [1] "NON-PLASTIC: 62.0184902295191"
```

```
kruskal.test(
  formula=generations_per_mrca_change~condition,
  data=summary_data
)
```


```
##
## Kruskal-Wallis rank sum test
##
## data: generations_per_mrca_change by condition
## Kruskal-Wallis chi-squared = 175.33, df = 2, p-value < 2.2e-16
```

```
pairwise.wilcox.test(
  x=summary_data$generations_per_mrca_change,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)
```

```
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: summary_data$generations_per_mrca_change and summary_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      1
##
## P value adjustment method: bonferroni
```

### 3.7 Phenotypic volatility along dominant lineage

```
ggplot(summary_data, aes(x=condition, y=dominant_lineage_trait_volatility, fill=condition)) +  
  geom_flat_violin(  
    position = position_nudge(x = .2, y = 0),  
    alpha = .8  
  ) +  
  geom_point(  
    mapping=aes(color=condition),  
    position = position_jitter(width = .15),  
    size = .5,  
    alpha = 0.8  
  ) +  
  geom_boxplot(  
    width = .1,  
    outlier.shape = NA,  
    alpha = 0.5  
  ) +  
  scale_x_discrete(  
    name="Condition",  
    limits=condition_order  
  ) +  
  scale_y_continuous(  
    name="Phenotypic volatility (log scale)",  
    trans="pseudo_log",  
    breaks=c(0, 10, 100, 1000, 10000),  
    limits=c(-1, 10000)  
  ) +  
  theme(  
    legend.position="none"  
  )
```



```
paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC"))$dominant_lineage_trait_volatility)
)

## [1] "PLASTIC: 2"

paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC"))$dominant_lineage_trait_volatility)
)

## [1] "STATIC: 0"

paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC"))$dominant_lineage_trait_volatility)
)

## [1] "NON-PLASTIC: 1868"

kruskal.test(
  formula=dominant_lineage_trait_volatility~condition,
  data=summary_data
)

##
## Kruskal-Wallis rank sum test
```

```
##
## data: dominant_lineage_trait_volatility by condition
## Kruskal-Wallis chi-squared = 190.78, df = 2, p-value < 2.2e-16
pairwise.wilcox.test(
  x=summary_data$dominant_lineage_trait_volatility,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)


##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: summary_data$dominant_lineage_trait_volatility and summary_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC < 2e-16      -
## STATIC  < 2e-16      8.7e-07
##
## P value adjustment method: bonferroni
```

### 3.7.1 Phenotypic volatility normalized by generations elapsed

```
summary_data$dominant_lineage_trait_volatility_per_generation <- summary_data$dominant_lineage_trait_volatility / summary_data$generations_elapsed

ggplot(summary_data, aes(x=condition, y=dominant_lineage_trait_volatility_per_generation)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
```

```
theme(
  legend.position="none"
)
```



```
paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC"))$dominant_lineage_trait_volatility_per_generation
)
```

```
## [1] "PLASTIC: 6.33339279717772e-05"
```

```
paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC"))$dominant_lineage_trait_volatility_per_generation
)
```

```
## [1] "STATIC: 0"
```

```
paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC"))$dominant_lineage_trait_volatility_per_generation
)
```

```
## [1] "NON-PLASTIC: 0.0447440145638177"
```

```
kruskal.test(
  formula=dominant_lineage_trait_volatility_per_generation~condition,
```

```

data=summary_data
)

##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_trait_volatility_per_generation by condition
## Kruskal-Wallis chi-squared = 189.62, df = 2, p-value < 2.2e-16

pairwise.wilcox.test(
  x=summary_data$dominant_lineage_trait_volatility_per_generation,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)

##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: summary_data$dominant_lineage_trait_volatility_per_generation and summary_data$dominant_lineage_trait_volatility_per_lineage_step
##
##      NON-PLASTIC PLASTIC
## PLASTIC < 2e-16      -
## STATIC  < 2e-16    4.2e-06
##
## P value adjustment method: bonferroni

```

### 3.7.2 Phenotypic volatility normalized by lineage length

Lineage length = number of genotypes along the lineage.

```

summary_data$dominant_lineage_trait_volatility_per_lineage_step <- summary_data$dominant_lineage_trait_volatility_per_generation / summary_data$dominant_lineage_length


ggplot(summary_data, aes(x=condition, y=dominant_lineage_trait_volatility_per_lineage_step)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  )

```

```

) +
scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
theme(
  legend.position="none"
)

```



```

paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC"))$dominant_lineage_trait_volatility_per_lineage_std
)

```

```
## [1] "PLASTIC: 0.00224688783339238"
```

```

paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC"))$dominant_lineage_trait_volatility_per_lineage_std
)

```

```
## [1] "STATIC: 0"
```

```

paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC"))$dominant_lineage_trait_volatility_per_lineage_std
)

```

```

)

## [1] "NON-PLASTIC: 0.437482522172625"

kruskal.test(
  formula=dominant_lineage_trait_volatility_per_lineage_step~condition,
  data=summary_data
)

##
##  Kruskal-Wallis rank sum test
##
## data:  dominant_lineage_trait_volatility_per_lineage_step by condition
## Kruskal-Wallis chi-squared = 191.23, df = 2, p-value < 2.2e-16

pairwise.wilcox.test(
  x=summary_data$dominant_lineage_trait_volatility_per_lineage_step,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)

##
##  Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data:  summary_data$dominant_lineage_trait_volatility_per_lineage_step and summary_data$condition
##
##           NON-PLASTIC PLASTIC
## PLASTIC < 2e-16      -
## STATIC  < 2e-16      2.3e-07
##
## P value adjustment method: bonferroni

```

### 3.7.3 Phenotypic fidelity

Frequency that an offspring's genotype is identical to a parent genotype (along the dominant lineage).

```

summary_data$dominant_lineage_trait_fidelity <- (summary_data$dominant_generation_born == summary_data$offspring_generation_born)

ggplot(summary_data, aes(x=condition, y=dominant_lineage_trait_fidelity, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
  )


```



```

    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  theme(
    legend.position="none"
  )

```



```

paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC"))$dominant_lineage_trait_fidelity)
)

```

```
## [1] "PLASTIC: 0.999936666072028"
```

```

paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC"))$dominant_lineage_trait_fidelity)
)

```

```
## [1] "STATIC: 1"
paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$dominant_lineage_trait_fidelity,
)

## [1] "NON-PLASTIC: 0.955255985436182"
kruskal.test(
  formula=dominant_lineage_trait_fidelity~condition,
  data=summary_data
)

##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_trait_fidelity by condition
## Kruskal-Wallis chi-squared = 189.62, df = 2, p-value < 2.2e-16
pairwise.wilcox.test(
  x=summary_data$dominant_lineage_trait_fidelity,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)


##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: summary_data$dominant_lineage_trait_fidelity and summary_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC < 2e-16      -
## STATIC  < 2e-16    4.2e-06
##
## P value adjustment method: bonferroni
```

### 3.8 Mutation accumulation along the dominant lineage

```
ggplot(summary_data, aes(x=condition, y=dominant_lineage_total_mut_cnt, fill=condition))
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
```

### 3.8. MUTATION ACCUMULATION ALONG THE DOMINANT LINEAGE<sup>35</sup>

```
mapping=aes(color=condition),
position = position_jitter(width = .15),
size = .5,
alpha = 0.8
) +
geom_boxplot(
width = .1,
outlier.shape = NA,
alpha = 0.5
) +
scale_x_discrete(
name="Condition",
limits=condition_order
) +
ylab("Mutation accumulation") +
theme(
legend.position="none"
)
)
```



```
paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC"))$dominant_lineage_total_mut_cnt)
)
```

```
## [1] "PLASTIC: 998.5"
```

```

paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC"))$dominant_lineage_total_mut_cnt)
)

## [1] "STATIC: 1100"

paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC"))$dominant_lineage_total_mut_cnt)
)

## [1] "NON-PLASTIC: 4657.5"

kruskal.test(
  formula=dominant_lineage_total_mut_cnt~condition,
  data=summary_data
)

##
## Kruskal-Wallis rank sum test
##
## data:  dominant_lineage_total_mut_cnt by condition
## Kruskal-Wallis chi-squared = 179.33, df = 2, p-value < 2.2e-16

pairwise.wilcox.test(
  x=summary_data$dominant_lineage_total_mut_cnt,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)

##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data:  summary_data$dominant_lineage_total_mut_cnt and summary_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.0019
##
## P value adjustment method: bonferroni


```

### 3.8.1 Mutation accumulation normalized by generations elapsed

```
summary_data$mutations_per_generation <- summary_data$dominant_lineage_total_mut_cnt /
```

### 3.8. MUTATION ACCUMULATION ALONG THE DOMINANT LINEAGE<sup>37</sup>

```
ggplot(summary_data, aes(x=condition, y=mutations_per_generation, fill=condition)) +  
  geom_flat_violin(  
    position = position_nudge(x = .2, y = 0),  
    alpha = .8  
  ) +  
  geom_point(  
    mapping=aes(color=condition),  
    position = position_jitter(width = .15),  
    size = .5,  
    alpha = 0.8  
  ) +  
  geom_boxplot(  
    width = .1,  
    outlier.shape = NA,  
    alpha = 0.5  
  ) +  
  scale_x_discrete(  
    name="Condition",  
    limits=condition_order  
  ) +  
  ylab("Mutation accumulation / generation") +  
  theme(  
    legend.position="none"  
  )  
)
```



```

paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC")$mutations_per_generation)
)

## [1] "PLASTIC: 0.0319267181456982"

paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC")$mutations_per_generation)
)

## [1] "STATIC: 0.0368157192941933"

paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$mutations_per_generation)
)

## [1] "NON-PLASTIC: 0.112804526786948"

kruskal.test(
  formula=mutations_per_generation~condition,
  data=summary_data
)

##
##  Kruskal-Wallis rank sum test
##
## data:  mutations_per_generation by condition
## Kruskal-Wallis chi-squared = 180.11, df = 2, p-value < 2.2e-16

pairwise.wilcox.test(
  x=summary_data$mutations_per_generation,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)


##
##  Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data:  summary_data$mutations_per_generation and summary_data$condition
##
##           NON-PLASTIC PLASTIC
## PLASTIC <2e-16          -
## STATIC  <2e-16        2e-04
##
## P value adjustment method: bonferroni

```

### 3.8. MUTATION ACCUMULATION ALONG THE DOMINANT LINEAGE<sup>39</sup>

#### 3.8.2 Mutation accumulation normalized by lineage length

```
summary_data$mutations_per_lineage_step <- summary_data$dominant_lineage_total_mut_cnt / summary_data$lineage_length
ggplot(summary_data, aes(x=condition, y=mutations_per_lineage_step, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylab("Mutation accumulation / lineage length") +
  theme(
    legend.position="none"
  )
```



```
paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC")$mutations_per_lineage_step)
)

## [1] "PLASTIC: 1.0328599144651"

paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC")$mutations_per_lineage_step)
)

## [1] "STATIC: 1.03794597464116"

paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$mutations_per_lineage_step)
)

## [1] "NON-PLASTIC: 1.10048311715591"

kruskal.test(
  formula=mutations_per_lineage_step~condition,
  data=summary_data
)

##
## Kruskal-Wallis rank sum test
```



### 3.8. MUTATION ACCUMULATION ALONG THE DOMINANT LINEAGE<sup>41</sup>

```
##
## data: mutations_per_lineage_step by condition
## Kruskal-Wallis chi-squared = 178.92, df = 2, p-value < 2.2e-16
pairwise.wilcox.test(
  x=summary_data$mutations_per_lineage_step,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)

##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: summary_data$mutations_per_lineage_step and summary_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.0034
##
## P value adjustment method: bonferroni
```


#### 3.8.3 Genotypic fidelity

The frequency that an offspring's genotype is the same as a parent's genotype.

```
summary_data$dominant_lineage_genotypic_fidelity <- (summary_data$dominant_generation_born - summary_data$dominant_generation) / summary_data$dominant_generation_born

ggplot(summary_data, aes(x=condition, y=dominant_lineage_genotypic_fidelity, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
```

```
theme(
  legend.position="none"
)
```



```
paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC")$dominant_lineage_genotypic_fidelity)
)

## [1] "PLASTIC: 0.969286906891951"

paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC")$dominant_lineage_genotypic_fidelity)
)

## [1] "STATIC: 0.964620594632577"

paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$dominant_lineage_genotypic_fidelity)
)

## [1] "NON-PLASTIC: 0.89754902563783"

kruskal.test(
  formula=dominant_lineage_genotypic_fidelity~condition,
```

### 3.8. MUTATION ACCUMULATION ALONG THE DOMINANT LINEAGE<sup>43</sup>

```
data=summary_data
)

##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_genotypic_fidelity by condition
## Kruskal-Wallis chi-squared = 179.86, df = 2, p-value < 2.2e-16
pairwise.wilcox.test(
  x=summary_data$dominant_lineage_genotypic_fidelity,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)


##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: summary_data$dominant_lineage_genotypic_fidelity and summary_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      2e-04
##
## P value adjustment method: bonferroni
```

#### 3.8.4 Characterizing variation along lineages

##### 3.8.4.1 How many mutation-steps along the lineage result in phenotypic changes?


```
summary_data$frac_phenotype_changing_mut_steps <- summary_data$dominant_lineage_num_mut_steps_tha

ggplot(filter(summary_data, dominant_lineage_num_mut_steps > 0), aes(x=frac_phenotype_changing_mu
  geom_density() +
  facet_wrap(
    ~condition,
    nrow=3,
    labeller=label_both
  ) +
  xlim(0, 1.0) +
  theme(
    legend.position="none"
  )
)
```



```
ggplot(summary_data, aes(x=condition, y=frac_phenotype_changing_mut_steps, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  theme(
    legend.position="none"
  )
```

### 3.8. MUTATION ACCUMULATION ALONG THE DOMINANT LINEAGE<sup>45</sup>



```
paste0(
  "PLASTIC: ",
  median(filter(summary_data, condition=="PLASTIC"))$frac_phenotype_changing_mut_steps)
)

## [1] "PLASTIC: 0.00224941742616098"

paste0(
  "STATIC: ",
  median(filter(summary_data, condition=="STATIC"))$frac_phenotype_changing_mut_steps)
)

## [1] "STATIC: 0"

paste0(
  "NON-PLASTIC: ",
  median(filter(summary_data, condition=="NON-PLASTIC"))$frac_phenotype_changing_mut_steps)
)

## [1] "NON-PLASTIC: 0.437583018324547"

kruskal.test(
  formula=frac_phenotype_changing_mut_steps~condition,
  data=summary_data
)

##
## Kruskal-Wallis rank sum test
```


```
##
## data: frac_phenotype_changing_mut_steps by condition
## Kruskal-Wallis chi-squared = 191.23, df = 2, p-value < 2.2e-16
pairwise.wilcox.test(
  x=summary_data$frac_phenotype_changing_mut_steps,
  g=summary_data$condition,
  p.adjust.method="bonferroni",
)
```

```
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: summary_data$frac_phenotype_changing_mut_steps and summary_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC < 2e-16      -
## STATIC  < 2e-16      2.3e-07
##
## P value adjustment method: bonferroni
```

```
summary_data$frac_unexpressed_mut_steps <- summary_data$dominant_lineage_num_mut_steps
summary_data$frac_expressed_mut_steps <- summary_data$dominant_lineage_num_mut_steps

ggplot(filter(summary_data, condition=="PLASTIC" & dominant_lineage_num_mut_steps_that,
  geom_density() +
  theme(
    legend.position="none"
  )
)
```

### 3.8. MUTATION ACCUMULATION ALONG THE DOMINANT LINEAGE<sup>47</sup>



```
print(paste0("PLASTIC - Mean with bootstrapped 95% CI"))
```

```
## [1] "PLASTIC - Mean with bootstrapped 95% CI"
```

```
bo <- boot(filter(summary_data, condition=="PLASTIC" & dominant_lineage_num_mut_steps_that_change
print(bo)
```

```
##
```

```
## ORDINARY NONPARAMETRIC BOOTSTRAP
```

```
##
```

```
##
```

```
## Call:
```

```
## boot(data = filter(summary_data, condition == "PLASTIC" & dominant_lineage_num_mut_steps_that_change
##       0)$frac_unexpressed_mut_steps, statistic = samplemean, R = 10000)
```

```
##
```

```
##
```

```
## Bootstrap Statistics :
```

```
##      original      bias    std. error
```

```
## t1* 0.8247126 -0.0002882184 0.04009371
```

```
print(boot.ci(bo, conf=0.95, type="perc"))
```

```
## BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
```

```
## Based on 10000 bootstrap replicates
```

```
##
```

```
## CALL :
```

```
## boot.ci(boot.out = bo, conf = 0.95, type = "perc")
```

```
##
## Intervals :
## Level      Percentile
## 95%      ( 0.7443,  0.9023 )
## Calculations and Intervals on Original Scale
plastic_summary_data <- filter(summary_data, condition=="PLASTIC")
aggregate_frac_mut_steps_that_change_unexpressed_phenotype <- sum(plastic_summary_data$dominant_lineage_num_mut_steps_that_change_unexpressed_phenotype)

## [1] 83
sum(plastic_summary_data$dominant_lineage_num_mut_steps_that_change_aggregate_phenotype)

## [1] 102
aggregate_frac_mut_steps_that_change_unexpressed_phenotype

## [1] 0.8137255

83 / 102 (0.8137255)
```

#### 3.8.4.3 For PLASTIC populations, what fraction of mutations that affect the unexpressed phenotype are deleterious versus beneficial?

```
aggregate_frac_unexpressed_deleterious_mut_steps <- sum(plastic_summary_data$dominant_lineage_num_mut_steps_that_change_unexpressed_phenotype > 0)
aggregate_frac_unexpressed_beneficial_mut_steps <- sum(plastic_summary_data$dominant_lineage_num_mut_steps_that_change_unexpressed_phenotype < 0)
```


Deleterious

```
summary_data$frac_unexpressed_deleterious_mut_steps <- summary_data$dominant_lineage_num_mut_steps_that_change_unexpressed_phenotype > 0

ggplot(
  filter(summary_data, condition=="PLASTIC" & dominant_lineage_num_mut_steps_that_change_unexpressed_phenotype > 0)
) +
  geom_density() +
  theme(
    legend.position="none"
  )
```



### 3.8. MUTATION ACCUMULATION ALONG THE DOMINANT LINEAGE<sup>49</sup>



```
bo <- boot(filter(summary_data, condition=="PLASTIC" & dominant_lineage_num_mut_steps_that_change
print(bo)
```

```
##
## ORDINARY NONPARAMETRIC BOOTSTRAP
##
## Call:
## boot(data = filter(summary_data, condition == "PLASTIC" & dominant_lineage_num_mut_steps_that_change
##       0)$frac_unexpressed_deleterious_mut_steps, statistic = samplemean,
##       R = 10000)
##
## Bootstrap Statistics :
##      original      bias    std. error
## t1* 0.5172414 0.000384023 0.03925893
print(boot.ci(bo, conf=0.95, type="perc"))


## BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
## Based on 10000 bootstrap replicates
##
## CALL :
## boot.ci(boot.out = bo, conf = 0.95, type = "perc")
##
## Intervals :
```

```
## Level      Percentile
## 95%      ( 0.4425,  0.5977 )
## Calculations and Intervals on Original Scale
```

Beneficial

```
summary_data$frac_unexpressed_beneficial_mut_steps <- summary_data$dominant_lineage_num_mut_steps_that_changed /
summary_data$dominant_lineage_num_mut_steps

ggplot(
  filter(summary_data, condition=="PLASTIC" & dominant_lineage_num_mut_steps_that_changed > 0)
  aes(x=frac_unexpressed_beneficial_mut_steps)
) +
  geom_density() +
  theme(
    legend.position="none"
  )
```



```
bo <- boot(filter(summary_data, condition=="PLASTIC" & dominant_lineage_num_mut_steps_that_changed > 0),
  statistic = samplemean,
  print=bo)
```


```
##
## ORDINARY NONPARAMETRIC BOOTSTRAP
##
##
## Call:
## boot(data = filter(summary_data, condition == "PLASTIC" & dominant_lineage_num_mut_steps_that_changed > 0),
##       statistic = samplemean,
```

```
##      R = 10000)
##
##
## Bootstrap Statistics :
##      original      bias      std. error
## t1* 0.4827586 -9.83908e-05  0.03960692
print(boot.ci(bo, conf=0.95, type="perc"))

## BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
## Based on 10000 bootstrap replicates
##
## CALL :
## boot.ci(boot.out = bo, conf = 0.95, type = "perc")
##
## Intervals :
## Level      Percentile
## 95%      ( 0.4046,  0.5609 )
## Calculations and Intervals on Original Scale
```

### 3.9 Depth of MRCA

```
ggplot(summary_data, aes(x=condition, y=phylo_mrca_depth, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylab("MRCA Depth") +
  theme(
    legend.position="none"
```



### 3.10 Manuscript figures

Figures styled for the paper.


#### 3.10.1 Evolutionary change panel

Selective sweeps, mutation accumulation, phenotypic volatility.

Mutation accumulation:

```
# dominant_lineage_total_mut_cnt or mutations_per_lineage_step?
mutation_count_fig <- ggplot(
  summary_data,
  aes(x=condition, y=dominant_lineage_total_mut_cnt, fill=condition)
) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
```

```
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order,
    labels=condition_order
  ) +
  scale_y_continuous(
    name="Mutations accumulated (log scale)",
    trans="log10",
    breaks=c(100, 1000, 10000),
    limits=c(100, 10000)
  ) +
  scale_fill_brewer(
    palette="Paired"
  ) +
  scale_color_brewer(
    palette="Paired"
  ) +
  coord_flip() +
  theme(
    legend.position="none"
  ) +
  ggsave(
    paste0(working_directory, "plots/", "mutation-accumulation.pdf"),
    width=5,
    height=4
  )
mutation_count_fig
```



Phenotypic volatility:


```
phenotypic_volatility_fig <- ggplot(
  summary_data,
  aes(x=condition, y=dominant_lineage_trait_volatility, fill=condition)
) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order,
    labels=condition_order
  )
```

```

) +
scale_y_continuous(
  name="Phenotypic volatility (log scale)",
  trans="pseudo_log",
  breaks=c(0, 10, 100, 1000, 10000),
  limits=c(-1,10000)
) +
scale_fill_brewer(
  palette="Paired"
) +
scale_color_brewer(
  palette="Paired"
) +
coord_flip() +
theme(
  legend.position="none"
) +
ggsave(
  paste0(working_directory, "plots/", "phenotypic-volatility.pdf"),
  width=4,
  height=4
)

```


phenotypic\_volatility\_fig



Selective sweeps:

```
selective_sweeps_fig <- ggplot(
  summary_data,
  aes(x=condition, y=phylo_mrca_changes, fill=condition)
) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order,
    labels=condition_order
  ) +
  scale_y_continuous(
    name="Coalescence Events (log scale)",
    trans="log10",
    breaks=c(10, 100, 1000),
    limits=c(10, 1000)
  ) +
  scale_fill_brewer(
    palette="Paired"
  ) +
  scale_color_brewer(
    palette="Paired"
  ) +
  coord_flip() +
  theme(
    legend.position="none"
  ) +
  ggsave(
    paste0(working_directory, "plots/", "selective-sweeps.pdf"),
    width=4,
    height=4
  )
```






All together:

```
grid <- plot_grid(
  selective_sweeps_fig + theme(
    legend.position="none"
  ),
  mutation_count_fig + theme(
    legend.position="none",
    axis.ticks.y=element_blank(),
    axis.text.y=element_blank(),
    axis.title.y=element_blank()
  ),
  phenotypic_volatility_fig + theme(
    legend.position="none",
    axis.ticks.y=element_blank(),
    axis.text.y=element_blank(),
    axis.title.y=element_blank()
  ),
  nrow=1,
  align="v",
  labels="auto"
```

```
)
grid
```



```
save_plot(
  paste0(working_directory, "plots/", "evolutionary-dynamics.pdf"),
  grid,
  base_height=6,
  base_asp=2.5
)
```

## Chapter 4

# Evolution and maintenance of novel traits

The effect of adaptive phenotypic plasticity on the evolution and maintenance of novel traits.

### 4.1 Overview

```
total_updates <- 200000
replicates <- 100

focal_traits <- c("not", "nand", "and", "ornot", "or", "andnot")
traits_set_a <- c("not", "and", "or")
traits_set_b <- c("nand", "ornot", "andnot")
extra_traits <- c(
  "nor", "xor", "equals",
  "logic_3aa", "logic_3ab", "logic_3ac",
  "logic_3ad", "logic_3ae", "logic_3af",
  "logic_3ag", "logic_3ah", "logic_3ai",
  "logic_3aj", "logic_3ak", "logic_3al",
  "logic_3am", "logic_3an", "logic_3ao",
  "logic_3ap", "logic_3aq", "logic_3ar",
  "logic_3as", "logic_3at", "logic_3au",
  "logic_3av", "logic_3aw", "logic_3ax",
  "logic_3ay", "logic_3az", "logic_3ba",
  "logic_3bb", "logic_3bc", "logic_3bd",
  "logic_3be", "logic_3bf", "logic_3bg",
  "logic_3bh", "logic_3bi", "logic_3bj",
  "logic_3bk", "logic_3bl", "logic_3bm",
```

```

    "logic_3bn", "logic_3bo", "logic_3bp",
    "logic_3bq", "logic_3br", "logic_3bs",
    "logic_3bt", "logic_3bu", "logic_3bv",
    "logic_3bw", "logic_3bx", "logic_3by",
    "logic_3bz", "logic_3ca", "logic_3cb",
    "logic_3cc", "logic_3cd", "logic_3ce",
    "logic_3cf", "logic_3cg", "logic_3ch",
    "logic_3ci", "logic_3cj", "logic_3ck",
    "logic_3cl", "logic_3cm", "logic_3cn",
    "logic_3co", "logic_3cp"
  )

# Relative location of data.
working_directory <- "experiments/2021-01-31-complex-features/analysis/" # << For book
# working_directory <- "/"

```

## 4.2 Analysis dependencies

Load all required R libraries.

```

library(ggplot2)
library(tidyverse)
library(cowplot)
library(RColorBrewer)
library(Hmisc)
library(boot)
source("https://gist.githubusercontent.com/benmarwick/2a1bb0133ff568cbe28d/raw/fb53bd9")

```

These analyses were conducted/knitted with the following computing environment:

```

print(version)

##
## platform      _
## arch          x86_64-pc-linux-gnu
## os            linux-gnu
## system        x86_64, linux-gnu
## status
## major         4
## minor         0.4
## year          2021
## month         02
## day           15
## svn rev       80002
## language      R

```

```
## version.string R version 4.0.4 (2021-02-15)
## nickname      Lost Library Book
```

## 4.3 Setup

```
##### summary data #####
summary_data_loc <- paste0(working_directory, "data/aggregate.csv")
summary_data <- read.csv(summary_data_loc, na.strings="NONE")

summary_data$DISABLE_REACTION_SENSORS <- as.factor(summary_data$DISABLE_REACTION_SENSORS)
summary_data$chg_env <- summary_data$chg_env == "True"
summary_data$dominant_plastic_odd_even <- as.factor(summary_data$dominant_plastic_odd_even)
summary_data$sensors <- summary_data$DISABLE_REACTION_SENSORS == "0"
summary_data$is_plastic <- summary_data$dominant_plastic_odd_even == "True"
summary_data$extra_task_value <- as.factor(summary_data$extra_task_value)
summary_data <- filter(summary_data, extra_task_value == 0.1)

env_label_fun <- function(chg_env) {
  if (chg_env) {
    return("Fluctuating")
  } else {
    return("Constant")
  }
}

sensors_label_fun <- function(has_sensors) {
  if (has_sensors) {
    return("Sensors")
  } else {
    return("No sensors")
  }
}

condition_label_fun <- function(has_sensors, env_chg) {
  if (has_sensors && env_chg) {
    return("PLASTIC")
  } else if (env_chg) {
    return("NON-PLASTIC")
  } else {
    return("STATIC")
  }
}

summary_data$env_label <- mapply(
```

```

    env_label_fun,
    summary_data$chg_env
  )
summary_data$sensors_label <- mapply(
  sensors_label_fun,
  summary_data$sensors
)
summary_data$condition <- mapply(
  condition_label_fun,
  summary_data$sensors,
  summary_data$chg_env
)

condition_order = c(
  "STATIC",
  "NON-PLASTIC",
  "PLASTIC"
)

##### time series #####
lineage_time_series_data_loc <- paste0(working_directory, "data/lineage_series.csv")
lineage_time_series_data <- read.csv(lineage_time_series_data_loc)

lineage_time_series_data$DISABLE_REACTION_SENSORS <- as.factor(lineage_time_series_data$DISABLE_REACTION_SENSORS == "True")
lineage_time_series_data$chg_env <- lineage_time_series_data$chg_env == "True"
lineage_time_series_data$sensors <- lineage_time_series_data$DISABLE_REACTION_SENSORS
lineage_time_series_data$extra_task_value <- as.factor(lineage_time_series_data$extra_task_value)

lineage_time_series_data$env_label <- mapply(
  env_label_fun,
  lineage_time_series_data$chg_env
)
lineage_time_series_data$sensors_label <- mapply(
  sensors_label_fun,
  lineage_time_series_data$sensors
)
lineage_time_series_data$condition <- mapply(
  condition_label_fun,
  lineage_time_series_data$sensors,
  lineage_time_series_data$chg_env
)

##### misc #####
# Configure our default graphing theme
theme_set(theme_cowplot())

```


```
dir.create(paste0(working_directory, "plots"), showWarnings=FALSE)
```

## 4.4 Evolution of phenotypic plasticity

For sensor-enabled populations in fluctuating environments, we only transferred populations containing an optimally plastic genotype to phase two.


```
summary_data_grouped = dplyr::group_by(summary_data, sensors, env_label, condition, extra_task_value)
summary_data_group_counts = dplyr::summarize(summary_data_grouped, n=dplyr::n())
```

```
ggplot(summary_data_group_counts, aes(x=condition, y=n, fill=condition)) +
  geom_col(position=position_dodge(0.9)) +
  geom_text(aes(label=n, y=n+2)) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylab("Number of replicates in phase two") +
  facet_wrap(~extra_task_value, labeller=label_both) +
  theme(
    legend.position="none"
  )
```



We can confirm our expectation that the dominant genotypes in non-plastic conditions are not phenotypically plastic.

```
summary_data_grouped = dplyr::group_by(summary_data, condition, is_plastic, extra_task_value)
summary_data_group_counts = dplyr::summarize(summary_data_grouped, n=dplyr::n())
ggplot(filter(summary_data_group_counts, is_plastic), aes(x=condition, y=n, fill=condition)) +
  geom_col(position=position_dodge(0.9)) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylim(0, 100) +
  geom_text(aes(label=n, y=n+1)) +
  ylab("Number of replicates with a plastic dominant genotype") +
  facet_wrap(~extra_task_value, labeller=label_both) +
  theme(
    legend.position="none"
  )
)
```




## 4.5 Final dominant novel task performance

How many novel tasks do final dominant genotypes perform?

```
ggplot(summary_data, aes(x=condition, y=dominant_extra_tasks, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  )
```



```
) +  
geom_point(  
  mapping=aes(color=condition),  
  position = position_jitter(width = .15),  
  size = .5,  
  alpha = 0.8  
) +  
geom_boxplot(  
  width = .1,  
  outlier.shape = NA,  
  alpha = 0.5  
) +  
scale_x_discrete(  
  name="Condition",  
  limits=condition_order  
) +  
ylab("Novel tasks performed by final dominant") +  
facet_wrap(  
  ~extra_task_value,  
  labeller=label_both  
) +  
theme(  
  legend.position="none"  
) +  
ggsave(  
  paste0(working_directory, "plots/dominant-extra-tasks.pdf"),  
  width=15,  
  height=10  
)
```



```
paste0(
  "PLASTIC median: ",
  median(filter(summary_data, condition=="PLASTIC")$dominant_extra_tasks)
)

## [1] "PLASTIC median: 3"

paste0(
  "STATIC median: ",
  median(filter(summary_data, condition=="STATIC")$dominant_extra_tasks)
)

## [1] "STATIC median: 3"

paste0(
  "NON-PLASTIC median: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$dominant_extra_tasks)
)

## [1] "NON-PLASTIC median: 0"

reward_level <- 0.1
dom_task_data <- filter(summary_data, extra_task_value==reward_level)
kruskal.test(
  formula=dominant_extra_tasks~condition,
  data=dom_task_data
)
```

```
##
## Kruskal-Wallis rank sum test
##
## data: dominant_extra_tasks by condition
## Kruskal-Wallis chi-squared = 177.17, df = 2, p-value < 2.2e-16

pairwise.wilcox.test(
  x=dom_task_data$dominant_extra_tasks,
  g=dom_task_data$condition,
  p.adjust.method="bonferroni",
  conf.int=TRUE,
  conf.level=0.95
)

##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: dom_task_data$dominant_extra_tasks and dom_task_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.9
##
## P value adjustment method: bonferroni
```

## 4.6 Final population novel task performance


How many novel tasks are performed across the final population (1% of organisms must perform to count)?

```
ggplot(summary_data, aes(x=condition, y=final_pop_extra_tasks_0.01, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
```

```

scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
facet_wrap(
  ~extra_task_value,
  labeller=label_both
) +
theme(
  legend.position="none"
)

```



```

paste0(
  "PLASTIC median: ",
  median(filter(summary_data, condition=="PLASTIC")$final_pop_extra_tasks_0.01)
)

```

```
## [1] "PLASTIC median: 3"
```

```

paste0(
  "STATIC median: ",
  median(filter(summary_data, condition=="STATIC")$final_pop_extra_tasks_0.01)
)

```

```
## [1] "STATIC median: 4"
```

```

paste0(
  "NON-PLASTIC median: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$final_pop_extra_tasks_0.01)
)

## [1] "NON-PLASTIC median: 0"

reward_level <- 0.1
dom_task_data <- filter(summary_data, extra_task_value==reward_level)
kruskal.test(
  formula=final_pop_extra_tasks_0.01~condition,
  data=dom_task_data
)

##
## Kruskal-Wallis rank sum test
##
## data: final_pop_extra_tasks_0.01 by condition
## Kruskal-Wallis chi-squared = 169.47, df = 2, p-value < 2.2e-16

pairwise.wilcox.test(
  x=dom_task_data$final_pop_extra_tasks_0.01,
  g=dom_task_data$condition,
  p.adjust.method="bonferroni",
  conf.int=TRUE,
  conf.level=0.95
)

##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: dom_task_data$final_pop_extra_tasks_0.01 and dom_task_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC < 2e-16      -
## STATIC  < 2e-16      0.00016
##
## P value adjustment method: bonferroni

```

## 4.7 Population-level novel tasks discovered

```


ggplot(summary_data, aes(x=condition, y=discovered_extra_tasks_0.01, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +

```

```

geom_point(
  mapping=aes(color=condition),
  position = position_jitter(width = .15),
  size = .5,
  alpha = 0.8
) +
geom_boxplot(
  width = .1,
  outlier.shape = NA,
  alpha = 0.5
) +
scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
facet_wrap(
  ~extra_task_value,
  labeller=label_both
) +
theme(
  legend.position="none"
)

```



```

paste0(
  "PLASTIC median: ",

```

```

    median(filter(summary_data, condition=="PLASTIC")$discovered_extra_tasks_0.01)
  )

## [1] "PLASTIC median: 8"
paste0(
  "STATIC median: ",
  median(filter(summary_data, condition=="STATIC")$discovered_extra_tasks_0.01)
)

## [1] "STATIC median: 9"
paste0(
  "NON-PLASTIC median: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$discovered_extra_tasks_0.01)
)

## [1] "NON-PLASTIC median: 13"
reward_level <- 0.1
dom_task_data <- filter(summary_data, extra_task_value==reward_level)
kruskal.test(
  formula=discovered_extra_tasks_0.01~condition,
  data=dom_task_data
)

##
## Kruskal-Wallis rank sum test
##
## data: discovered_extra_tasks_0.01 by condition
## Kruskal-Wallis chi-squared = 24.271, df = 2, p-value = 5.365e-06
pairwise.wilcox.test(
  x=dom_task_data$discovered_extra_tasks_0.01,
  g=dom_task_data$condition,
  p.adjust.method="bonferroni",
  conf.int=TRUE,
  conf.level=0.95
)

##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: dom_task_data$discovered_extra_tasks_0.01 and dom_task_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC 2.4e-05      -
## STATIC  0.00035     1.00000
##

```


```
## P value adjustment method: bonferroni
```

## 4.8 Novel tasks along lineage of final dominant genotype

```
ggplot(summary_data, aes(x=condition, y=dominant_lineage_extra_traits_discovered, fill=
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylab("Extra tasks discovered along dominant lineage") +
  facet_wrap(
    ~extra_task_value,
    labeller=label_both
  ) +
  theme(
    legend.position="none"
  ) +
  ggsave(
    paste0(working_directory, "plots/dominant-lineage-extra-tasks-discovered.pdf"),
    width=15,
    height=10
  )
)
```



#### 4.8. NOVEL TASKS ALONG LINEAGE OF FINAL DOMINANT GENOTYPE73



```
paste0(
  "PLASTIC median: ",
  median(filter(summary_data, condition=="PLASTIC")$dominant_lineage_extra_traits_discovered)
)
```

```
## [1] "PLASTIC median: 3.5"
```

```
paste0(
  "STATIC median: ",
  median(filter(summary_data, condition=="STATIC")$dominant_lineage_extra_traits_discovered)
)
```

```
## [1] "STATIC median: 4"
```

```
paste0(
  "NON-PLASTIC median: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$dominant_lineage_extra_traits_discovered)
)
```

```
## [1] "NON-PLASTIC median: 6"
```

```
reward_level <- 0.1
dom_task_data <- filter(summary_data, extra_task_value==reward_level)
kruskal.test(
  formula=dominant_lineage_extra_traits_discovered~condition,
  data=dom_task_data
)
```

```
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_extra_traits_discovered by condition
## Kruskal-Wallis chi-squared = 24.099, df = 2, p-value = 5.846e-06

pairwise.wilcox.test(
  x=dom_task_data$dominant_lineage_extra_traits_discovered,
  g=dom_task_data$condition,
  p.adjust.method="bonferroni",
  conf.int=TRUE,
  conf.level=0.95
)

##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: dom_task_data$dominant_lineage_extra_traits_discovered and dom_task_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC 1.7e-05      -
## STATIC  0.0035      0.0561
##
## P value adjustment method: bonferroni
```


#### 4.8.1.1 Novel traits discovered per step

This isn't totally fair to non-plastic lineages because they're continuously re-adapting, so they have more genotypes along lineage.

```
summary_data$dominant_lineage_extra_traits_discovered_per_step <- summary_data$dominant_lineage_extra_traits_discovered_per_step
ggplot(summary_data, aes(x=condition, y=dominant_lineage_extra_traits_discovered_per_step)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
```

#### 4.8. NOVEL TASKS ALONG LINEAGE OF FINAL DOMINANT GENOTYPE75

```
scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
ylab("Lineage task discovery (per step)" ) +
facet_wrap(
  ~extra_task_value,
  labeller=label_both
) +
theme(
  legend.position="none"
)
```



```
paste0(
  "PLASTIC median: ",
  median(filter(summary_data, condition=="PLASTIC"))$dominant_lineage_extra_traits_discovered_per_s
)
```

```
## [1] "PLASTIC median: 0.00484428434398198"
```

```
paste0(
  "STATIC median: ",
  median(filter(summary_data, condition=="STATIC"))$dominant_lineage_extra_traits_discovered_per_s
)
```

```
## [1] "STATIC median: 0.00480194844967106"
```

```

paste0(
  "NON-PLASTIC median: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$dominant_lineage_extra_traits_
)

## [1] "NON-PLASTIC median: 0.00139827576402932"

reward_level <- 0.1
dom_task_data <- filter(summary_data, extra_task_value==reward_level)
kruskal.test(
  formula=dominant_lineage_extra_traits_discovered_per_step~condition,
  data=dom_task_data
)

##
##  Kruskal-Wallis rank sum test
##
## data:  dominant_lineage_extra_traits_discovered_per_step by condition
## Kruskal-Wallis chi-squared = 106.72, df = 2, p-value < 2.2e-16

pairwise.wilcox.test(
  x=dom_task_data$dominant_lineage_extra_traits_discovered_per_step,
  g=dom_task_data$condition,
  p.adjust.method="bonferroni",
  conf.int=TRUE,
  conf.level=0.95
)

##
##  Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data:  dom_task_data$dominant_lineage_extra_traits_discovered_per_step and dom_task_
##
##           NON-PLASTIC PLASTIC
## PLASTIC 9.7e-11      -
## STATIC  < 2e-16      0.67
##
## P value adjustment method: bonferroni

```

#### 4.8.2 Novel tasks discovered per generation


```

summary_data$dominant_lineage_extra_traits_discovered_per_generation <- summary_data$d
ggplot(summary_data, aes(x=condition, y=dominant_lineage_extra_traits_discovered_per_g
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +

```

#### 4.8. NOVEL TASKS ALONG LINEAGE OF FINAL DOMINANT GENOTYPE77

```
geom_point(
  mapping=aes(color=condition),
  position = position_jitter(width = .15),
  size = .5,
  alpha = 0.8
) +
geom_boxplot(
  width = .1,
  outlier.shape = NA,
  alpha = 0.5
) +
scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
ylab("Lineage task discovery (per generation)") +
facet_wrap(
  ~extra_task_value,
  labeller=label_both
) +
theme(
  legend.position="none"
)
```



```

paste0(
  "PLASTIC median: ",
  median(filter(summary_data, condition=="PLASTIC"))$dominant_lineage_extra_traits_discovered_per_generation
)

## [1] "PLASTIC median: 0.000117695011124939"

paste0(
  "STATIC median: ",
  median(filter(summary_data, condition=="STATIC"))$dominant_lineage_extra_traits_discovered_per_generation
)

## [1] "STATIC median: 0.00015363220504867"

paste0(
  "NON-PLASTIC median: ",
  median(filter(summary_data, condition=="NON-PLASTIC"))$dominant_lineage_extra_traits_discovered_per_generation
)

## [1] "NON-PLASTIC median: 0.00014358046266055"

reward_level <- 0.1
dom_task_data <- filter(summary_data, extra_task_value==reward_level)
kruskal.test(
  formula=dominant_lineage_extra_traits_discovered_per_generation~condition,
  data=dom_task_data
)

##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_extra_traits_discovered_per_generation by condition
## Kruskal-Wallis chi-squared = 7.1465, df = 2, p-value = 0.02806

pairwise.wilcox.test(
  x=dom_task_data$dominant_lineage_extra_traits_discovered_per_generation,
  g=dom_task_data$condition,
  p.adjust.method="bonferroni",
  conf.int=TRUE,
  conf.level=0.95
)


##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: dom_task_data$dominant_lineage_extra_traits_discovered_per_generation and dom_task_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC 0.092          -

```

#### 4.8. NOVEL TASKS ALONG LINEAGE OF FINAL DOMINANT GENOTYPE79

```
## STATIC 1.000      0.025
##
## P value adjustment method: bonferroni
```

```
ggplot(summary_data, aes(x=condition, y=dominant_lineage_extra_traits_gained, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylab("Extra tasks gained along dominant lineage") +
  facet_wrap(
    ~extra_task_value,
    labeller=label_both
  ) +
  theme(
    legend.position="none"
  ) +
  ggsave(
    paste0(working_directory, "plots/dominant-lineage-extra-tasks-gained.pdf"),
    width=15,
    height=10
  )
```



#### 4.8.4 Novel tasks lost

```
ggplot(summary_data, aes(x=condition, y=dominant_lineage_extra_traits_lost, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylab("Extra tasks lost along dominant lineage") +
  facet_wrap(
```




#### 4.8. NOVEL TASKS ALONG LINEAGE OF FINAL DOMINANT GENOTYPE81

```

~extra_task_value,
labeller=label_both
) +
theme(
  legend.position="none"
) +
ggsave(
  paste0(working_directory, "plots/dominant-lineage-extra-tasks-lost.pdf"),
  width=15,
  height=10
)

```



```

paste0(
  "PLASTIC median: ",
  median(filter(summary_data, condition=="PLASTIC")$dominant_lineage_extra_traits_lost)
)

```

```
## [1] "PLASTIC median: 2"
```

```

paste0(
  "STATIC median: ",
  median(filter(summary_data, condition=="STATIC")$dominant_lineage_extra_traits_lost)
)

```

```
## [1] "STATIC median: 5"
```

```

paste0(
  "NON-PLASTIC median: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$dominant_lineage_extra_traits_lost)
)

## [1] "NON-PLASTIC median: 87.5"

reward_level <- 0.1
dom_task_data <- filter(summary_data, extra_task_value==reward_level)
kruskal.test(
  formula=dominant_lineage_extra_traits_lost~condition,
  data=dom_task_data
)

##
##  Kruskal-Wallis rank sum test
##
## data:  dominant_lineage_extra_traits_lost by condition
## Kruskal-Wallis chi-squared = 129.06, df = 2, p-value < 2.2e-16

pairwise.wilcox.test(
  x=dom_task_data$dominant_lineage_extra_traits_lost,
  g=dom_task_data$condition,
  p.adjust.method="bonferroni",
  conf.int=TRUE,
  conf.level=0.95
)

##
##  Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data:  dom_task_data$dominant_lineage_extra_traits_lost and dom_task_data$condition
##
##           NON-PLASTIC PLASTIC
## PLASTIC 2.7e-16      -
## STATIC  < 2e-16      0.0024
##
## P value adjustment method: bonferroni

```

#### 4.8.4.1 Novel traits lost per step

Again, not totally fair to non-plastic lineages.

```

summary_data$dominant_lineage_extra_traits_lost_per_step <- summary_data$dominant_lineage_extra_traits_lost /
  dominant_lineage_extra_traits_lost
ggplot(summary_data, aes(x=condition, y=dominant_lineage_extra_traits_lost_per_step, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  )


```

#### 4.8. NOVEL TASKS ALONG LINEAGE OF FINAL DOMINANT GENOTYPE83

```

) +
geom_point(
  mapping=aes(color=condition),
  position = position_jitter(width = .15),
  size = .5,
  alpha = 0.8
) +
geom_boxplot(
  width = .1,
  outlier.shape = NA,
  alpha = 0.5
) +
scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
ylab("Extra tasks lost along dominant lineage (per step)" ) +
facet_wrap(
  ~extra_task_value,
  labeller=label_both
) +
theme(
  legend.position="none"
)

```



```

paste0(
  "PLASTIC median: ",
  median(filter(summary_data, condition=="PLASTIC"))$dominant_lineage_extra_traits_lost_per_step
)

## [1] "PLASTIC median: 0.00238455242036334"

paste0(
  "STATIC median: ",
  median(filter(summary_data, condition=="STATIC"))$dominant_lineage_extra_traits_lost_per_step
)

## [1] "STATIC median: 0.00544747485837901"

paste0(
  "NON-PLASTIC median: ",
  median(filter(summary_data, condition=="NON-PLASTIC"))$dominant_lineage_extra_traits_lost_per_step
)

## [1] "NON-PLASTIC median: 0.0216427755153431"

reward_level <- 0.1
dom_task_data <- filter(summary_data, extra_task_value==reward_level)
kruskal.test(
  formula=dominant_lineage_extra_traits_lost_per_step~condition,
  data=dom_task_data
)

##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_extra_traits_lost_per_step by condition
## Kruskal-Wallis chi-squared = 65.779, df = 2, p-value = 5.204e-15

pairwise.wilcox.test(
  x=dom_task_data$dominant_lineage_extra_traits_lost_per_step,
  g=dom_task_data$condition,
  p.adjust.method="bonferroni",
  conf.int=TRUE,
  conf.level=0.95
)


##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: dom_task_data$dominant_lineage_extra_traits_lost_per_step and dom_task_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC 1.3e-10      -

```

#### 4.8. NOVEL TASKS ALONG LINEAGE OF FINAL DOMINANT GENOTYPE85

```
## STATIC 1.7e-10 0.0092
##
## P value adjustment method: bonferroni
```

```
summary_data$dominant_lineage_extra_traits_lost_per_generation <- summary_data$dominant_lineage_e
ggplot(summary_data, aes(x=condition, y=dominant_lineage_extra_traits_lost_per_generation, fill=c
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylab("Extra tasks lost along dominant lineage (per generation)") +
  facet_wrap(
    ~extra_task_value,
    labeller=label_both
  ) +
  theme(
    legend.position="none"
  )
)
```



```
paste0(
  "PLASTIC median: ",
  median(filter(summary_data, condition=="PLASTIC")$dominant_lineage_extra_traits_lost_per_generation)
)

## [1] "PLASTIC median: 6.25141973661864e-05"

paste0(
  "STATIC median: ",
  median(filter(summary_data, condition=="STATIC")$dominant_lineage_extra_traits_lost_per_generation)
)

## [1] "STATIC median: 0.000161396283669756"

paste0(
  "NON-PLASTIC median: ",
  median(filter(summary_data, condition=="NON-PLASTIC")$dominant_lineage_extra_traits_lost_per_generation)
)

## [1] "NON-PLASTIC median: 0.0022026054610079"

reward_level <- 0.1
dom_task_data <- filter(summary_data, extra_task_value==reward_level)
kruskal.test(
  formula=dominant_lineage_extra_traits_lost_per_generation~condition,
  data=dom_task_data
)
```

#### 4.8. NOVEL TASKS ALONG LINEAGE OF FINAL DOMINANT GENOTYPE<sup>87</sup>

```
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_extra_traits_lost_per_generation by condition
## Kruskal-Wallis chi-squared = 121.41, df = 2, p-value < 2.2e-16

pairwise.wilcox.test(
  x=dom_task_data$dominant_lineage_extra_traits_lost_per_generation,
  g=dom_task_data$condition,
  p.adjust.method="bonferroni",
  conf.int=TRUE,
  conf.level=0.95
)


##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: dom_task_data$dominant_lineage_extra_traits_lost_per_generation and dom_task_data$condi
##
##          NON-PLASTIC PLASTIC
## PLASTIC 1.1e-15      -
## STATIC  < 2e-16    0.0012
##
## P value adjustment method: bonferroni
```

##### 4.8.4.3 How many instances of novel trait loss co-occur with changes in base phenotype?

Task loss linked with primary trait changes.

```
lost_traits_summary_data <- filter(summary_data, extra_task_value==0.1 & dominant_lineage_extra_t
lost_traits_summary_data$frac_linked_extra_trait_loss <- lost_traits_summary_data$dominant_lineag

ggplot(lost_traits_summary_data, aes(x=frac_linked_extra_trait_loss)) +
  geom_density() +
  facet_grid(
    condition~extra_task_value,
    labeller=label_both
  ) +
  theme(
    legend.position="none"
  ) +
  ggsave(
    paste0(working_directory, "plots/dominant-lineage-extra-tasks-lost-linkage.pdf"),
    width=15,
    height=10
  )
```




```
ggplot(lost_traits_summary_data, aes(x=condition, y=frac_linked_extra_trait_loss, fill=
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  facet_wrap(
    ~extra_task_value,
    labeller=label_both
  ) +
  theme(
```



#### 4.8. NOVEL TASKS ALONG LINEAGE OF FINAL DOMINANT GENOTYPE89

```
legend.position="none"
)
```



```
paste0(
  "PLASTIC median: ",
  median(filter(loss_traits_summary_data, condition=="PLASTIC")$frac_linked_extra_trait_loss)
)
```

```
## [1] "PLASTIC median: 0.0192307692307692"
```

```
paste0(
  "STATIC median: ",
  median(filter(loss_traits_summary_data, condition=="STATIC")$frac_linked_extra_trait_loss)
)
```

```
## [1] "STATIC median: 0"
```

```
paste0(
  "NON-PLASTIC median: ",
  median(filter(loss_traits_summary_data, condition=="NON-PLASTIC")$frac_linked_extra_trait_loss)
)
```

```
## [1] "NON-PLASTIC median: 0.983803278688525"
```

```
kruskal.test(
  formula=frac_linked_extra_trait_loss~condition,
  data=loss_traits_summary_data
)
```

```

)

##
##  Kruskal-Wallis rank sum test
##
## data:  frac_linked_extra_trait_loss by condition
## Kruskal-Wallis chi-squared = 153.68, df = 2, p-value < 2.2e-16
pairwise.wilcox.test(
  x=lost_traits_summary_data$frac_linked_extra_trait_loss,
  g=lost_traits_summary_data$condition,
  p.adjust.method="bonferroni",
  conf.int=TRUE,
  conf.level=0.95
)

##
##  Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data:  lost_traits_summary_data$frac_linked_extra_trait_loss and lost_traits_summary_data$condition
##
##           NON-PLASTIC PLASTIC
## PLASTIC 1.9e-08      -
## STATIC  < 2e-16      1.8e-06
##
## P value adjustment method: bonferroni
sum(filter(lost_traits_summary_data, condition=="NON-PLASTIC")$dominant_lineage_extra_trait_loss)
## [1] 10998
sum(filter(lost_traits_summary_data, condition=="NON-PLASTIC")$dominant_lineage_extra_trait_loss)
## [1] 11229
aggregate_frac_linked_extra_trait_loss_nonplastic <- sum(filter(lost_traits_summary_data, condition=="NON-PLASTIC")$dominant_lineage_extra_trait_loss)
aggregate_frac_linked_extra_trait_loss_nonplastic
## [1] 0.9794283
sum(filter(lost_traits_summary_data, condition=="PLASTIC")$dominant_lineage_extra_trait_loss)
## [1] 29
sum(filter(lost_traits_summary_data, condition=="PLASTIC")$dominant_lineage_extra_trait_loss)
## [1] 142
aggregate_frac_linked_extra_trait_loss_plastic <- sum(filter(lost_traits_summary_data, condition=="PLASTIC")$dominant_lineage_extra_trait_loss)
aggregate_frac_linked_extra_trait_loss_plastic

```

```
## [1] 0.2042254
```

```
sum(filter(lost_traits_summary_data, condition=="STATIC")$dominant_lineage_extra_traits_lost_link)
```

```
## [1] 13
```

```
sum(filter(lost_traits_summary_data, condition=="STATIC")$dominant_lineage_extra_traits_lost)
```

```
## [1] 631
```

```
aggregate_frac_linked_extra_trait_loss_nonplastic <- sum(filter(lost_traits_summary_data, condition=="STATIC")$dominant_lineage_extra_traits_lost)
aggregate_frac_linked_extra_trait_loss_nonplastic
```


```
## [1] 0.02060222
```

## 4.9 Extra task performance over time

Match score over time

```
lineage_reward10 <- filter(lineage_time_series_data, extra_task_value=="0.1")
```

```
ggplot(filter(lineage_reward10, update>198000 & update<=200000), aes(x=update, y=match_score_even)) +
  stat_summary(fun="mean", geom="line") +
  stat_summary(
    fun.data="mean_cl_boot",
    fun.args=list(conf.int=0.95),
    geom="ribbon",
    alpha=0.2,
    linetype=0
  ) +
  ylab("Match score (even environment)") +
  facet_wrap(
    ~extra_task_value,
    labeller=label_both
  ) +
  ggsave(
    paste0(working_directory, "plots/dominant-lineage-match-score-even-val10.png"),
    width=15,
    height=10
  )
```



Extra tasks over time

```
ggplot(filter(lineage_reward10, update>198000 & update<=200000), aes(x=update, y=extra.
  stat_summary(fun="mean", geom="line") +
  stat_summary(
    fun.data="mean_cl_boot",
    fun.args=list(conf.int=0.95),
    geom="ribbon",
    alpha=0.2,
    linetype=0
  ) +
  ylab("Number of extra traits") +
  facet_wrap(
    ~extra_task_value,
    labeller=label_both
  ) +
  ggsave(
    paste0(working_directory, "plots/dominant-lineage-extra-traits-val10.png"),
    width=15,
    height=10
  )
)
```



## 4.10 Manuscript figures


Final dominant extra tasks.

```
extra_task_reward_value=0.1
dominant_extra_tasks_fig <- ggplot(
  filter(summary_data, extra_task_value==extra_task_reward_value),
  aes(x=condition, y=dominant_extra_tasks, fill=condition)
) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
```

```

scale_x_discrete(
  name="Condition",
  limits=condition_order,
  labels=condition_order
) +
scale_y_continuous(
  name="Final dominant novel traits"
) +
scale_fill_brewer(
  palette="Paired"
) +
scale_color_brewer(
  palette="Paired"
) +
theme(
  legend.position="none"
) +
coord_flip()
dominant_extra_tasks_fig

```




Final dominant lineage tasks discovered.

```

lineage_extra_tasks_discovered_fig <- ggplot(
  filter(summary_data, extra_task_value==extra_task_reward_value),
  aes(x=condition, y=dominant_lineage_extra_traits_discovered, fill=condition)
)

```

```
) +  
geom_flat_violin(  
  position = position_nudge(x = .2, y = 0),  
  alpha = .8  
) +  
geom_point(  
  mapping=aes(color=condition),  
  position = position_jitter(width = .15),  
  size = .5,  
  alpha = 0.8  
) +  
geom_boxplot(  
  width = .1,  
  outlier.shape = NA,  
  alpha = 0.5  
) +  
scale_x_discrete(  
  name="Condition",  
  limits=condition_order,  
  labels=condition_order  
) +  
scale_y_continuous(  
  name="Novel traits discovered on lineage"  
) +  
scale_fill_brewer(  
  palette="Paired"  
) +  
scale_color_brewer(  
  palette="Paired"  
) +  
theme(  
  legend.position="none"  
) +  
coord_flip()  
lineage_extra_tasks_discovered_fig
```



Final dominant lineage tasks lost.

```
lineage_extra_tasks_lost_fig <- ggplot(
  filter(summary_data, extra_task_value == extra_task_reward_value),
  aes(x=condition, y=dominant_lineage_extra_traits_lost, fill=condition)
) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order,
    labels=condition_order
  )
```




```

) +
scale_y_continuous(
  name="Novel traits lost on lineage (log scale)",
  trans="pseudo_log",
  breaks=c(0,10,100,1000),
  limits=c(-1,1000)
) +
scale_fill_brewer(
  palette="Paired"
) +
scale_color_brewer(
  palette="Paired"
) +
theme(
  legend.position="none"
) +
coord_flip()

```

lineage\_extra\_tasks\_lost\_fig



Pull it all together.

```


grid <- plot_grid(
  dominant_extra_tasks_fig,
  lineage_extra_tasks_discovered_fig + theme(axis.ticks.y=element_blank(),axis.text.y=element_bla

```

```

lineage_extra_tasks_lost_fig + theme(axis.ticks.y=element_blank(),axis.text.y=element_blank(),
nrow=1,
align="v",
labels="auto"
)
save_plot(
  paste0(working_directory, "plots/", "complex-traits-panel.pdf"),
  grid,
  base_height=6,
  base_asp=2.5
)
grid

```



## Chapter 5

# Genetic hitchhiking

The effect of adaptive phenotypic plasticity on (deleterious) genetic hitchhiking.

### 5.1 Overview

```
total_updates <- 200000
replicates <- 100

focal_traits <- c("not", "nand", "and", "ornot", "or", "andnot")
traits_set_a <- c("not", "and", "or")
traits_set_b <- c("nand", "ornot", "andnot")

# Relative location of data.
working_directory <- "experiments/2021-02-05-hitchhiking/analysis/" # << For bookdown
# working_directory <- "./"
```

### 5.2 Analysis dependencies

Load all required R libraries.

```
library(RColorBrewer)
library(ggplot2)
library(tidyverse)
library(cowplot)
library(Hmisc)
library(boot)
library(fmsb)
source("https://gist.githubusercontent.com/benmarwick/2a1bb0133ff568cbe28d/raw/fb53bd97121f7f9ce9")
```

These analyses were conducted/knitted with the following computing environment:

```
print(version)

##
## platform      x86_64-pc-linux-gnu
## arch          x86_64
## os            linux-gnu
## system        x86_64, linux-gnu
## status
## major         4
## minor         0.4
## year          2021
## month         02
## day           15
## svn rev       80002
## language      R
## version.string R version 4.0.4 (2021-02-15)
## nickname      Lost Library Book
```

### 5.3 Setup

```
##### summary data #####
summary_data_loc <- paste0(working_directory, "data/aggregate.csv")
summary_data <- read.csv(summary_data_loc, na.strings="NONE")

summary_data$DISABLE_REACTION_SENSORS <- as.factor(summary_data$DISABLE_REACTION_SENSORS)
summary_data$chg_env <- summary_data$chg_env == "True"
summary_data$dominant_plastic_odd_even <- as.factor(summary_data$dominant_plastic_odd_even)
summary_data$sensors <- summary_data$DISABLE_REACTION_SENSORS == "0"
summary_data$is_plastic <- summary_data$dominant_plastic_odd_even == "True"
summary_data$POISON_PENALTY <- as.factor(summary_data$POISON_PENALTY)

env_label_fun <- function(chg_env) {
  if (chg_env) {
    return("Fluctuating")
  } else {
    return("Constant")
  }
}

sensors_label_fun <- function(has_sensors) {
  if (has_sensors) {
    return("Sensors")
  }
}
```

```

    } else {
      return("No sensors")
    }
  }

condition_label_fun <- function(has_sensors, env_chg) {
  if (has_sensors && env_chg) {
    return("PLASTIC")
  } else if (env_chg) {
    return("NON-PLASTIC")
  } else {
    return("STATIC")
  }
}

summary_data$env_label <- mapapply(
  env_label_fun,
  summary_data$chg_env
)
summary_data$sensors_label <- mapapply(
  sensors_label_fun,
  summary_data$sensors
)
summary_data$condition <- mapapply(
  condition_label_fun,
  summary_data$sensors,
  summary_data$chg_env
)

condition_order = c(
  "STATIC",
  "NON-PLASTIC",
  "PLASTIC"
)

##### time series #####
lineage_time_series_data_loc <- paste0(working_directory, "data/lineage_series.csv")
lineage_time_series_data <- read.csv(lineage_time_series_data_loc)

lineage_time_series_data$DISABLE_REACTION_SENSORS <- as.factor(lineage_time_series_data$DISABLE_F
lineage_time_series_data$chg_env <- lineage_time_series_data$chg_env == "True"
lineage_time_series_data$sensors <- lineage_time_series_data$DISABLE_REACTION_SENSORS == "0"
lineage_time_series_data$POISON_PENALTY <- as.factor(lineage_time_series_data$POISON_VALUE)

lineage_time_series_data$env_label <- mapapply(

```

```

    env_label_fun,
    lineage_time_series_data$chg_env
  )
  lineage_time_series_data$sensors_label <- mapapply(
    sensors_label_fun,
    lineage_time_series_data$sensors
  )
  lineage_time_series_data$condition <- mapapply(
    condition_label_fun,
    lineage_time_series_data$sensors,
    lineage_time_series_data$chg_env
  )

##### misc #####
# Configure our default graphing theme
theme_set(theme_cowplot())
dir.create(paste0(working_directory, "plots"), showWarnings=FALSE)

```

## 5.4 Evolution of phenotypic plasticity

For sensor-enabled populations in fluctuating environments, we only transferred populations containing an optimally plastic genotype to phase-two.

```


summary_data_grouped = dplyr::group_by(summary_data, sensors, env_label, condition, POI)
summary_data_group_counts = dplyr::summarize(summary_data_grouped, n=dplyr::n())

```

```

ggplot(summary_data_group_counts, aes(x=condition, y=n, fill=condition)) +
  geom_col(position=position_dodge(0.9)) +
  geom_text(aes(label=n, y=n+2)) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylab("Number of replicates in phase two") +
  facet_wrap(~POISON_PENALTY, labeller=label_both) +
  theme(
    legend.position="none"
  )


```



We can confirm our expectation that the dominant genotypes in non-plastic conditions are not phenotypically plastic.

```
summary_data_grouped = dplyr::group_by(summary_data, condition, is_plastic, POISON_PENALTY)
summary_data_group_counts = dplyr::summarize(summary_data_grouped, n=dplyr::n())
```

```
## `summarise()` has grouped output by 'condition', 'is_plastic'. You can override using the `.gr
ggplot(filter(summary_data_group_counts, is_plastic), aes(x=condition, y=n, fill=condition)) +
  geom_col(position=position_dodge(0.9)) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  geom_text(aes(label=n, y=n+1)) +
  ylab("Number of replicates with a plastic dominant genotype") +
  ylim(0, 100) +
  facet_wrap(~POISON_PENALTY, labeller=label_both) +
  theme(
    legend.position="none"
  )
```



## 5.5 Hitchhiking instruction execution

### 5.5.1 Number of replicates where final dominant genotype executes hitchhiker instruction

```
hitchiker_penalty <- 0.1
```

```
occurrences <- c(
  length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="NON-PLASTIC")),
  length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="PLASTIC")),
  length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="STATIC"))
)
trials <- c(
  length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="NON-PLASTIC")),
  length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="PLASTIC")),
  length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="STATIC"))
)
names(trials) <- c(
  "NON-PLASTIC",
  "PLASTIC",
  "STATIC"
)
names(occurrences) <- c(
  "NON-PLASTIC",
```



```

    "PLASTIC",
    "STATIC"
)

pairwise.fisher.test(x=occurrences, n=trials, p.adjust.method="bonferroni")

##
## Pairwise comparisons using Pairwise comparison of proportions (Fisher)
##
## data:  occurrences out of trials
##
##          NON-PLASTIC PLASTIC
## PLASTIC 0.03212      -
## STATIC  0.00022      1.00000
##
## P value adjustment method: bonferroni

```

### 5.5.2 Final dominant genotype hitchhiker execution

```


ggplot(summary_data, aes(x=condition, y=dominant_times_poison_executed, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  ylab("poison instruction executions (final dominant)") +
  facet_wrap(
    ~POISON_PENALTY,
    labeller=label_both,
    scale="free_y"
  ) +

```

```

theme(
  legend.position="none"
) +
ggsave(
  paste0(working_directory, "plots/dominant-poison.pdf"),
  width=15,
  height=10
)

```



```

penalties <- levels(summary_data$POISON_PENALTY)
for (penalty in penalties) {
  stat_data <- filter(summary_data, POISON_PENALTY==penalty)
  print(
    paste0(
      "PENALTY: ", penalty
    )
  )
  kt <- kruskal.test(
    formula=dominant_times_poison_executed~condition,
    data=stat_data
  )
  print(
    kt
  )
  if (is.na(kt$p.value)) { next }
}

```

```

if (kt$p.value > 0.05) { next }
print(
  pairwise.wilcox.test(
    x=stat_data$dominant_times_poison_executed,
    g=stat_data$condition,
    p.adjust.method="bonferroni"
  )
)
}

## [1] "PENALTY: 0"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_times_poison_executed by condition
## Kruskal-Wallis chi-squared = 36.988, df = 2, p-value = 9.294e-09
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_times_poison_executed and stat_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC 2.8e-07      -
## STATIC  0.00015     0.00198
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.03"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_times_poison_executed by condition
## Kruskal-Wallis chi-squared = 72.995, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_times_poison_executed and stat_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC 2.0e-06      -
## STATIC  2.8e-13      1
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.1"

```

```
##
## Kruskal-Wallis rank sum test
##
## data: dominant_times_poison_executed by condition
## Kruskal-Wallis chi-squared = 21.157, df = 2, p-value = 2.546e-05
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_times_poison_executed and stat_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC 0.02034      -
## STATIC  0.00022      -
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.3"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_times_poison_executed by condition
## Kruskal-Wallis chi-squared = NaN, df = 2, p-value = NA
```


### 5.5.3 Hitchhiker instruction execution in final population

```
ggplot(summary_data, aes(x=condition, y=final_population_poison, fill=condition)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
```

```

scale_y_continuous(
  name="poison instruction executions (final population)",
  trans="pseudo_log",
  breaks=c(0,10,100,1000, 10000, 100000, 1000000),
  limits=c(-1,1000000)
) +
facet_wrap(
  ~POISON_PENALTY,
  labeller=label_both
) +
theme(
  legend.position="none"
) +
ggsave(
  paste0(working_directory, "plots/final-population-poison-log.pdf"),
  width=15,
  height=10
)

```



```

penalties <- levels(summary_data$POISON_PENALTY)
for (penalty in penalties) {
  stat_data <- filter(summary_data, POISON_PENALTY==penalty)
  print(
    paste0(
      "PENALTY: ", penalty
    )
  )
}

```

```

    )
  )
  kt <- kruskal.test(
    formula=final_population_poison~condition,
    data=stat_data
  )
  print(
    kt
  )
  if (is.na(kt$p.value)) { next }
  if (kt$p.value > 0.05) { next }
  print(
    pairwise.wilcox.test(
      x=stat_data$final_population_poison,
      g=stat_data$condition,
      p.adjust.method="bonferroni"
    )
  )
}

## [1] "PENALTY: 0"
##
## Kruskal-Wallis rank sum test
##
## data: final_population_poison by condition
## Kruskal-Wallis chi-squared = 43.589, df = 2, p-value = 3.426e-10
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$final_population_poison and stat_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC 8.7e-07      -
## STATIC  9.8e-07      0.00074
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.03"
##
## Kruskal-Wallis rank sum test
##
## data: final_population_poison by condition
## Kruskal-Wallis chi-squared = 20.74, df = 2, p-value = 3.136e-05
##
##

```


```
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$final_population_poison and stat_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC 0.003          -
## STATIC  1e-04          1.000
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.1"
##
## Kruskal-Wallis rank sum test
##
## data: final_population_poison by condition
## Kruskal-Wallis chi-squared = 20.608, df = 2, p-value = 3.35e-05
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$final_population_poison and stat_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC 0.0093          -
## STATIC  4.9e-05          1.0000
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.3"
##
## Kruskal-Wallis rank sum test
##
## data: final_population_poison by condition
## Kruskal-Wallis chi-squared = 3.3994, df = 2, p-value = 0.1827
```

#### 5.5.4 Hitchhiker instruction execution along final dominant lineage (cummulative)

```
ggplot(summary_data, aes(x=condition, y=dominant_lineage_times_poison_executed, fill=condition))
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
```

```
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  scale_y_continuous(
    name="poison instruction executions (dominant lineage)",
    trans="pseudo_log",
    breaks=c(0,10,100,1000,10000,100000),
    limits=c(-1,100000)
  ) +
  facet_wrap(
    ~POISON_PENALTY,
    labeller=label_both
  ) +
  theme(
    legend.position="none"
  ) +
  ggsave(
    paste0(working_directory, "plots/final-dominant-lineage-poison-log.pdf"),
    width=15,
    height=10
  )
)
```





```

penalties <- levels(summary_data$POISON_PENALTY)
for (penalty in penalties) {
  stat_data <- filter(summary_data, POISON_PENALTY==penalty)
  print(
    paste0(
      "PENALTY: ", penalty
    )
  )
  kt <- kruskal.test(
    formula=dominant_lineage_times_poison_executed~condition,
    data=stat_data
  )
  print(
    kt
  )
  if (is.na(kt$p.value)) { next }
  if (kt$p.value > 0.05) { next }
  print(
    pairwise.wilcox.test(
      x=stat_data$dominant_lineage_times_poison_executed,
      g=stat_data$condition,
      p.adjust.method="bonferroni"
    )
  )
}

```

```

## [1] "PENALTY: 0"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_times_poison_executed by condition
## Kruskal-Wallis chi-squared = 178.84, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_lineage_times_poison_executed and stat_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.0018
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.03"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_times_poison_executed by condition
## Kruskal-Wallis chi-squared = 178.62, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_lineage_times_poison_executed and stat_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.011
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.1"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_times_poison_executed by condition
## Kruskal-Wallis chi-squared = 184.83, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_lineage_times_poison_executed and stat_data$condition
##

```

```
##          NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.21
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.3"
##
## Kruskal-Wallis rank sum test
##
## data:  dominant_lineage_times_poison_executed by condition
## Kruskal-Wallis chi-squared = 149.48, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data:  stat_data$dominant_lineage_times_poison_executed and stat_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC 4.4e-16      -
## STATIC  < 2e-16      0.84
##
## P value adjustment method: bonferroni
```

## 5.6 Characterizing mutations that increase hitchhiker instruction execution


### 5.6.1 Number of offspring along dominant lineage with increase in hitchhiker instruction execution

```
ggplot(summary_data, aes(x=condition, y=dominant_lineage_num_times_hitchhike_inst_exec_increases,
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
```

```

) +
scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
facet_wrap(
  ~POISON_PENALTY,
  labeller=label_both,
  scales="free_y"
) +
theme(
  legend.position="none"
) +
ggsave(
  paste0(working_directory, "plots/final-dominant-lineage-poison-increase-num-mutant"),
  width=15,
  height=10
)

```



```

penalties <- levels(summary_data$POISON_PENALTY)
for (penalty in penalties) {
  stat_data <- filter(summary_data, POISON_PENALTY==penalty)
  print(
    paste0(
      "PENALTY: ", penalty
    )
  )
}

```

```

    )
  )
  kt <- kruskal.test(
    formula=dominant_lineage_num_times_hitchhike_inst_exec_increases~condition,
    data=stat_data
  )
  print(
    kt
  )
  if (is.na(kt$p.value)) { next }
  if (kt$p.value > 0.05) { next }
  print(
    pairwise.wilcox.test(
      x=stat_data$dominant_lineage_num_times_hitchhike_inst_exec_increases,
      g=stat_data$condition,
      p.adjust.method="bonferroni"
    )
  )
}

```

```

## [1] "PENALTY: 0"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_num_times_hitchhike_inst_exec_increases by condition
## Kruskal-Wallis chi-squared = 179.79, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_lineage_num_times_hitchhike_inst_exec_increases and stat_data$condit
##
##      NON-PLASTIC PLASTIC
## PLASTIC < 2e-16      -
## STATIC  < 2e-16      0.00046
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.03"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_num_times_hitchhike_inst_exec_increases by condition
## Kruskal-Wallis chi-squared = 179.35, df = 2, p-value < 2.2e-16
##
##

```

```

## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_lineage_num_times_hitchhike_inst_exec_increases and stat_c
##
##          NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.03
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.1"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_num_times_hitchhike_inst_exec_increases by condition
## Kruskal-Wallis chi-squared = 185.34, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_lineage_num_times_hitchhike_inst_exec_increases and stat_c
##
##          NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.27
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.3"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_num_times_hitchhike_inst_exec_increases by condition
## Kruskal-Wallis chi-squared = 146.35, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_lineage_num_times_hitchhike_inst_exec_increases and stat_c
##
##          NON-PLASTIC PLASTIC
## PLASTIC 7.8e-16      -
## STATIC  < 2e-16      0.86
##
## P value adjustment method: bonferroni
sum(filter(summary_data, condition=="NON-PLASTIC" & POISON_PENALTY==0.1)$dominant_lineage

```

## 5.6. CHARACTERIZING MUTATIONS THAT INCREASE HITCHHIKER INSTRUCTION EXECUTION119

```
## [1] 1916
```

```
sum(filter(summary_data, condition=="PLASTIC" & POISON_PENALTY==0.1)$dominant_lineage_num_times_h
```

```
## [1] 18
```


```
sum(filter(summary_data, condition=="STATIC" & POISON_PENALTY==0.1)$dominant_lineage_num_times_hi
```

```
## [1] 58
```

```
# sum(filter(summary_data, condition=="NON-PLASTIC" & POISON_PENALTY==0.1)$dominant_lineage_trait_vol  
# sum(filter(summary_data, condition=="PLASTIC" & POISON_PENALTY==0.1)$dominant_lineage_trait_vol  
# sum(filter(summary_data, condition=="STATIC" & POISON_PENALTY==0.1)$dominant_lineage_trait_vol
```

### 5.6.1.1 Normalized by generations

```
summary_data$dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation <- summary_d  
ggplot(summary_data, aes(x=condition, y=dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation))  
  geom_flat_violin(  
    position = position_nudge(x = .2, y = 0),  
    alpha = .8  
  ) +  
  geom_point(  
    mapping=aes(color=condition),  
    position = position_jitter(width = .15),  
    size = .5,  
    alpha = 0.8  
  ) +  
  geom_boxplot(  
    width = .1,  
    outlier.shape = NA,  
    alpha = 0.5  
  ) +  
  scale_x_discrete(  
    name="Condition",  
    limits=condition_order  
  ) +  
  facet_wrap(  
    ~POISON_PENALTY,  
    labeller=label_both,  
    scales="free_y"  
  ) +  
  theme(  
    legend.position="none"  
  )
```



```

penalties <- levels(summary_data$POISON_PENALTY)
for (penalty in penalties) {
  stat_data <- filter(summary_data, POISON_PENALTY==penalty)
  print(
    paste0(
      "PENALTY: ", penalty
    )
  )
  kt <- kruskal.test(
    formula=dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation ~ condition,
    data=stat_data
  )
  print(
    kt
  )
  if (is.na(kt$p.value)) { next }
  if (kt$p.value > 0.05) { next }
  print(
    pairwise.wilcox.test(
      x=stat_data$dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation,
      g=stat_data$condition,
      p.adjust.method="bonferroni"
    )
  )
}

```



## 5.6. CHARACTERIZING MUTATIONS THAT INCREASE HITCHHIKER INSTRUCTION EXECUTION121

```
## [1] "PENALTY: 0"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation by condition
## Kruskal-Wallis chi-squared = 180.05, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation and s
##
##      NON-PLASTIC PLASTIC
## PLASTIC < 2e-16      -
## STATIC  < 2e-16      7.8e-05
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.03"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation by condition
## Kruskal-Wallis chi-squared = 176.25, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation and s
##
##      NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.019
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.1"
##
## Kruskal-Wallis rank sum test
##
## data: dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation by condition
## Kruskal-Wallis chi-squared = 184.17, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation and s
##
```


```
##          NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.2
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.3"
##
## Kruskal-Wallis rank sum test
##
## data:  dominant_lineage_num_times_hitchhike_inst_exec_increases_per_generation by c
## Kruskal-Wallis chi-squared = 140.99, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data:  stat_data$dominant_lineage_num_times_hitchhike_inst_exec_increases_per_genera
##
##          NON-PLASTIC PLASTIC
## PLASTIC 2.2e-15      -
## STATIC  < 2e-16      0.79
##
## P value adjustment method: bonferroni
```

### 5.6.2 What fraction of mutations that increase hitchhiker instruction execution co-occur with base trait changes?

```
# Fraction of unexpressed vs expressed increases in hitchhiker instructions
summary_data$frac_hitchhiking_linked_trait_change <- summary_data$dominant_lineage_num_


ggplot(filter(summary_data, dominant_lineage_num_times_hitchhike_inst_exec_increases>0)
  geom_density() +
  facet_grid(
    condition~POISON_PENALTY,
    labeller=label_both,
    scales="free_y"
  ) +
  theme(
    legend.position="none"
  ) +
  ggsave(
    paste0(working_directory, "plots/dominant-lineage-frac_hitchhiking_linked_trait_cha
    width=15,
    height=10
  )
```

## 5.6. CHARACTERIZING MUTATIONS THAT INCREASE HITCHHIKER INSTRUCTION EXECUTION123



```
ggplot(filter(summary_data, dominant_lineage_num_times_hitchhike_inst_exec_increases>0 ), aes(x=frac_hitchhiking_linked_trait_change)) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order
  ) +
  facet_wrap(
    ~POISSON_PENALTY,
    labeller=label_both,
    scales="free_y"
  ) +
```

```
theme(
  legend.position="none"
)
```



```
penalties <- levels(summary_data$POISSON_PENALTY)
for (penalty in penalties) {
  stat_data <- filter(summary_data, POISSON_PENALTY==penalty & dominant_lineage_num_time > 0)
  print(
    paste0(
      "PENALTY: ", penalty
    )
  )
  kt <- kruskal.test(
    formula=frac_hitchhiking_linked_trait_change~condition,
    data=stat_data
  )
  print(
    kt
  )
  if (is.na(kt$p.value)) { next }
  if (kt$p.value > 0.05) { next }
  print(
    pairwise.wilcox.test(
      x=stat_data$frac_hitchhiking_linked_trait_change,
      g=stat_data$condition,
```

## 5.6. CHARACTERIZING MUTATIONS THAT INCREASE HITCHHIKER INSTRUCTION EXECUTION125

```

    p.adjust.method="bonferroni",
    exact=FALSE
  )
}

```

```

## [1] "PENALTY: 0"
##
## Kruskal-Wallis rank sum test
##
## data: frac_hitchhiking_linked_trait_change by condition
## Kruskal-Wallis chi-squared = 211.29, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$frac_hitchhiking_linked_trait_change and stat_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC <2e-16      -
## STATIC  <2e-16      0.031
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.03"
##
## Kruskal-Wallis rank sum test
##
## data: frac_hitchhiking_linked_trait_change by condition
## Kruskal-Wallis chi-squared = 186.88, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$frac_hitchhiking_linked_trait_change and stat_data$condition
##
##      NON-PLASTIC PLASTIC
## PLASTIC 2.9e-16      -
## STATIC  < 2e-16      -
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.1"
##
## Kruskal-Wallis rank sum test
##
## data: frac_hitchhiking_linked_trait_change by condition

```

```

## Kruskal-Wallis chi-squared = 113.72, df = 2, p-value < 2.2e-16
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$frac_hitchhiking_linked_trait_change and stat_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC 3.3e-08      -
## STATIC  < 2e-16      -
##
## P value adjustment method: bonferroni
## [1] "PENALTY: 0.3"
##
## Kruskal-Wallis rank sum test
##
## data: frac_hitchhiking_linked_trait_change by condition
## Kruskal-Wallis chi-squared = 34.791, df = 2, p-value = 2.788e-08
##
##
## Pairwise comparisons using Wilcoxon rank sum test with continuity correction
##
## data: stat_data$frac_hitchhiking_linked_trait_change and stat_data$condition
##
##          NON-PLASTIC PLASTIC
## PLASTIC 0.26          -
## STATIC  2.4e-08      0.18
##
## P value adjustment method: bonferroni
denom <- sum(filter(summary_data, condition=="NON-PLASTIC" & POISON_PENALTY==0.1)$dominant_1)
num <- sum(filter(summary_data, condition=="NON-PLASTIC" & POISON_PENALTY==0.1)$dominant_0)
paste0("NON-PLASTIC: ", num/denom, " (", num, "/", denom, ")")

## [1] "NON-PLASTIC: 0.498956158663883(956/1916)"

denom <- sum(filter(summary_data, condition=="PLASTIC" & POISON_PENALTY==0.1)$dominant_1)
num <- sum(filter(summary_data, condition=="PLASTIC" & POISON_PENALTY==0.1)$dominant_0)
paste0("PLASTIC: ", num/denom, " (", num, "/", denom, ")")

## [1] "PLASTIC: 0 (0/18)"

denom <- sum(filter(summary_data, condition=="STATIC" & POISON_PENALTY==0.1)$dominant_1)
num <- sum(filter(summary_data, condition=="STATIC" & POISON_PENALTY==0.1)$dominant_0)
paste0("STATIC: ", num/denom, " (", num, "/", denom, ")")


## [1] "STATIC: 0 (0/58)"

```

### 5.6.3 What about unexpressed vs expressed trait changes in plastic populations?

```
summary_data$frac_unexpressed_hitchhiker_inc <- summary_data$dominant_lineage_num_times_hitchhike_inst_exec_increases>0 & condition == "PLASTIC"
summary_data$frac_expressed_hitchhiker_inc <- summary_data$dominant_lineage_num_times_hitchhike_inst_exec_increases>0 & condition == "PLASTIC"

ggplot(filter(summary_data, dominant_lineage_num_times_hitchhike_inst_exec_increases>0 & condition == "PLASTIC")) +
  geom_density() +
  facet_grid(
    condition~POISON_PENALTY,
    labeller=label_both,
    scales="free_y"
  ) +
  theme(
    legend.position="none"
  )
```




```
# ggplot(filter(summary_data, dominant_lineage_num_times_hitchhike_inst_exec_increases>0 & condition == "PLASTIC")) +
#   geom_density() +
#   facet_grid(
#     condition~POISON_PENALTY,
#     labeller=label_both,
#     scales="free_y"
#   ) +
#   theme(
```

```
#     legend.position="none"
#   )
```

```
hitchiker_penalty <- 0.1
5.7 Manuscript Figures
ggplot(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="NON-PLASTIC")) +
  geom_bar() +
  geom_text(
    stat="count",
    mapping=aes(label=..count..),
    position=position_dodge(0.9),
    vjust=0
  ) +
  scale_y_continuous(
    name="Number of replicates where poison is executed by dominant genotype",
    limits=c(0, replicates+2)
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order,
    labels=condition_order,
    breaks=condition_order
  ) +
  scale_fill_brewer(
    palette="Paired"
  ) +
  scale_color_brewer(
    palette="Paired"
  ) +
  coord_flip() +
  theme(
    legend.position="none"
  )
```






```
fig_data <- data.frame(
  num_execute_poison=c(
    length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="NON-PLASTIC" & domina
    length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="PLASTIC" & domina
    length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="STATIC" & domina
  ),
  num_execute_no_poison=c(
    length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="NON-PLASTIC" & do
    length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="PLASTIC" & domina
    length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="STATIC" & domina
  ),
  total=c(
    length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="NON-PLASTIC")$RANDOM
    length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="PLASTIC")$RANDOM
    length(filter(summary_data, POISON_PENALTY==hitchiker_penalty & condition=="STATIC" )$RANDOM
  ),
  condition=c(
    "NON-PLASTIC",
    "PLASTIC",
    "STATIC"
  )
)

ggplot(fig_data, aes(x=condition, y=num_execute_poison, fill=condition)) +
  geom_col(position=position_dodge(0.9)) +
```

```

geom_text(aes(label=num_execute_poison, y=num_execute_poison+0.25)) +
scale_x_discrete(
  name="Condition",
  limits=condition_order
) +
scale_fill_brewer(
  palette="Paired"
) +
scale_color_brewer(
  palette="Paired"
) +
theme(
  legend.position="none"
)

```




Number of poison increases along lineage.

```

poison_increases_fig <- ggplot(
  filter(summary_data, POISON_PENALTY==hitchiker_penalty),
  aes(x=condition, y=dominant_lineage_num_times_hitchhike_inst_exec_increases, fill=
) +
geom_flat_violin(
  position = position_nudge(x = .2, y = 0),
  alpha = .8
) +

```

```
geom_point(  
  mapping=aes(color=condition),  
  position = position_jitter(width = .15),  
  size = .5,  
  alpha = 0.8  
) +  
geom_boxplot(  
  width = .1,  
  outlier.shape = NA,  
  alpha = 0.5  
) +  
scale_x_discrete(  
  name="Condition",  
  limits=condition_order,  
  labels=condition_order  
) +  
scale_y_continuous(  
  name="Increases in poison instruction execution along lineage",  
) +  
scale_fill_brewer(  
  palette="Paired"  
) +  
scale_color_brewer(  
  palette="Paired"  
) +  
theme(  
  legend.position="none"  
) +  
coord_flip()  
  
poison_increases_fig
```




```
poison_increases_per_gen_fig <- ggplot(
  filter(summary_data, POISON_PENALTY==hitchiker_penalty),
  aes(x=condition, y=dominant_lineage_num_times_hitchhike_inst_exec_increases_per_gen)
) +
  geom_flat_violin(
    position = position_nudge(x = .2, y = 0),
    alpha = .8
  ) +
  geom_point(
    mapping=aes(color=condition),
    position = position_jitter(width = .15),
    size = .5,
    alpha = 0.8
  ) +
  geom_boxplot(
    width = .1,
    outlier.shape = NA,
    alpha = 0.5
  ) +
  scale_x_discrete(
    name="Condition",
    limits=condition_order,
    labels=condition_order
  ) +
  scale_y_continuous(
```

```

    name="Increases in poison instruction execution along lineage (per generation)",
  ) +
  scale_fill_brewer(
    palette="Paired"
  ) +
  scale_color_brewer(
    palette="Paired"
  ) +
  theme(
    legend.position="none"
  ) +
  coord_flip()
poison_increases_per_gen_fig

```



Pull it all together.

```

# grid <- plot_grid(
#   dominant_extra_tasks_fig,
#   lineage_extra_tasks_discovered_fig + theme(axis.ticks.y=element_blank(),axis.text.y=element_blank()),
#   lineage_extra_tasks_lost_fig + theme(axis.ticks.y=element_blank(),axis.text.y=element_blank()),
#   nrow=1,
#   align="v",
#   labels="auto"
# )

```

```
# save_plot(  
#   paste0(working_directory, "plots/", "complex-traits-panel.pdf"),  
#   grid,  
#   base_height=6,  
#   base_asp=2.5  
# )  
# grid
```