

R version 4.0.2 (2020-06-22) -- "Taking Off Again"
 Copyright (C) 2020 The R Foundation for Statistical Computing
 Platform: x86_64-w64-mingw32/x64 (64-bit)

R is free software and comes with ABSOLUTELY NO WARRANTY.
 You are welcome to redistribute it under certain conditions.
 Type 'license()' or 'licence()' for distribution details.

Natural language support but running in an English locale

R is a collaborative project with many contributors.
 Type 'contributors()' for more information and
 'citation()' on how to cite R or R packages in publications.

Type 'demo()' for some demos, 'help()' for on-line help, or
 'help.start()' for an HTML browser interface to help.
 Type 'q()' to quit R.

[Previously saved workspace restored]

```
> #Experiment 10
> #Date 1-12-2020
> #Slot lab L1
> #Title Test on Single Proportion
> #Name Amlan Nayak
> #Reg No. 19BCD7143
> ##Course Applied statistics
> #Code MAT1011
> #Suppose 60% of citizens voted in last election
> #In a telephonic survey, they said that 85 out of #148 people voted in current election
> #At 5% level of significance, can we reject the null #hypothesis that the proportion of voters in
> #the population is above 60% this year.
> #Null hypothesis H0:p0.6
> #Alternate hypothesis H1:p<=0.6
> #Left tailed test
> pbar=85/148
> pbar
[1] 0.5743243
> [1] 0.5743243
Error: unexpected '[' in "["
> p0=0.6
> q0=1-p0
> q0
[1] 0.4
> [1] 0.4
Error: unexpected '[' in "["
> n=148
> z=(pbar-p0)/(sqrt((p0*q0)/n))
> z
[1] -0.6375983
> [1] -0.6375983
Error: unexpected '[' in "["
> # z0 is negative and left tailed test
> #Pvalue=P(Z<=z0)
> Pvalue=pnorm(z)
> Pvalue
[1] 0.2618676
> [1] 0.2618676
Error: unexpected '[' in "["
> #Rejected area=significance level=0.05=alpha
> #Pvalue is much greater than alpha
> #Accept null hypothesis
> #Suppose that 12% of apples harvested in an orchard #last year.30 out of 214 apples in a harvest sample #this year turns to be rotten. At 5% level of significance #can we reject null hypothesis that the proportion #of rotten apples in harvest stays below 12% this year. #Null hypothesis H0:p<0.12
> #Alternate hypotheiss H1:p=0.12 Right tailed test pbar=30/214
> pbar
[1] 0.5743243
> [1] 0.1401869
Error: unexpected '[' in "["
> p0=0.12
```

```

> q0=1-p0
> q0
[1] 0.88
> [1] 0.88
Error: unexpected '[' in "["
> z=(pbar-p0)/(sqrt((p0*q0)/214))
> z
[1] 20.45224
> R Console Page 2
Error: unexpected symbol in "R Console"
> [1] 0.908751
Error: unexpected '[' in "["
> #positive and Right tailed test Pvalue=P(Z=z0)=1-P(Z<=z0) Pvalue=1-pnorm(z)
> Pvalue
[1] 0.2618676
> [1] 0.1817408
Error: unexpected '[' in "["
> #Alpha=0.05
> #pvaluealpha Accept H0.
> #Two tailed test problem
> #Suppose a coin toss turns up 12 heads out of
> #20 trials. At 5% level of significance, can we
> #reject Null hypothesis the coin toss is fair
> #Null hypothesis H0: p=0.5
> #Alternate hypothesis H1:p not equal to 0.5
> pbar=12/20
> pbar
[1] 0.6
> [1] 0.6
Error: unexpected '[' in "["
> p0=0.5
> q0=1-p0
> q0
[1] 0.5
> [1] 0.5
Error: unexpected '[' in "["
> z=(pbar-p0)/(sqrt((p0*q0)/20))
> z
[1] 0.8944272
> [1] 0.8944272
Error: unexpected '[' in "["
> #Since it is a two tailed test
> # and z0 is positive ,Pvalue is 2*P(Z=z0)
> #Pvalue=2*(1-P(Z<=z0))
> Pvalue=2*(1-pnorm(z))
> Pvalue
[1] 0.3710934
> [1] 0.3710934
Error: unexpected '[' in "["
> alpha=0.05
> #Pvaluealpha
> #Accept hypothesis
> #Accept Null Hypothesis H0.
> #Assignment problem
> #z for the data set specified in a, with a mean equal to 13
> # and a variance equal to 16.
> a <- c(10, 8, 7, 12, 9, 6, 7, 8)
> mu <- 13
> var <- 16
> z.test = function(a, mu, var){
+   zeta = (mean(a) - mu) / (sqrt(var / length(a)))
+   return(zeta)
+ }
> z.test(a, mu, var)
[1] -3.270369
> #Group A with an early morning class of 400 students with 342 female students.
> #Group B with a late class of 400 students with 290 female students.
> #Use a 5% alpha level. We want to know, whether the proportions
> #of females are the same in the two groups of the student?
> prop.test(x = c(342, 290),
+           n = c(400, 400))

```

2-sample test for equality of proportions with continuity correction

```
data:  c(342, 290) out of c(400, 400)
X-squared = 19.598, df = 1, p-value = 9.559e-06
alternative hypothesis: two.sided
95 percent confidence interval:
 0.07177443 0.18822557
sample estimates:
prop 1 prop 2
 0.855  0.725
```

```
> prop.test(x = c(342, 290),
+           n = c(400, 400),
+           alternative = "less")
```

2-sample test for equality of proportions with continuity correction

```
data:  c(342, 290) out of c(400, 400)
X-squared = 19.598, df = 1, p-value = 1
alternative hypothesis: less
95 percent confidence interval:
 -1.0000000 0.1792664
sample estimates:
prop 1 prop 2
 0.855  0.725
```

```
> prop.test(x = c(342, 290),
+           n = c(400, 400),
+           alternative = "greater")
```

2-sample test for equality of proportions with continuity correction

```
data:  c(342, 290) out of c(400, 400)
X-squared = 19.598, df = 1, p-value = 4.779e-06
alternative hypothesis: greater
95 percent confidence interval:
 0.08073363 1.00000000
sample estimates:
prop 1 prop 2
 0.855  0.725
```

```
>
```