

R version 4.0.2 (2020-06-22) -- "Taking Off Again"
 Copyright (C) 2020 The R Foundation for Statistical Computing
 Platform: x86_64-w64-mingw32/x64 (64-bit)

R is free software and comes with ABSOLUTELY NO WARRANTY.
 You are welcome to redistribute it under certain conditions.
 Type 'license()' or 'licence()' for distribution details.

Natural language support but running in an English locale

R is a collaborative project with many contributors.
 Type 'contributors()' for more information and
 'citation()' on how to cite R or R packages in publications.

Type 'demo()' for some demos, 'help()' for on-line help, or
 'help.start()' for an HTML browser interface to help.
 Type 'q()' to quit R.

[Previously saved workspace restored]

```
> #name-Amlan Shivam Nayak
> #Reg no.-19BCD7143
> #date-7/12/20
> #Applied statistics mat1011
> #Lab-L1
> #An outbreak of Salmonella related illness was attributed to ice produced at
> #a certain factory. Scientists measured the level of Salmonella in 9
> #randomly sampled batches ice cream. The levels in mpn.gm were :
> #0.593 0.142 0.329 0.691 0.231 0.793 0.519 0.392 0.418
> #Is there any evidence that the mean level of salmonella in the ice cream
> #greater than 0.3 mpn/gm ?
> #H0:mu<=0.3
> #H1:mu0.3
> #Right tailed test
> x=c(0.593,0.142,0.329,0.691,0.231,0.793,0.519,0.392,0.418)
> t.test(x,mu=0.3,alternative="greater")
```

One Sample t-test

```
data: x
t = 2.2051, df = 8, p-value = 0.02927
alternative hypothesis: true mean is greater than 0.3
95 percent confidence interval:
 0.3245133      Inf
sample estimates:
mean of x
0.4564444
```

```
> One Sample t-test
Error: unexpected symbol in " One Sample"
> data: x
Error in data:x : NA/NaN argument
In addition: Warning message:
In data:x : numerical expression has 9 elements: only the first used
> t = 2.2051, df = 8, p-value = 0.02927
Error: unexpected ',' in "t = 2.2051,"
> alternative hypothesis: true mean is greater than 0.3
Error: unexpected symbol in "alternative hypothesis"
> 95 percent confidence interval:
Error: unexpected symbol in "95 percent"
> 0.3245133 Inf
Error: unexpected numeric constant in " 0.3245133 Inf"
> sample estimates:
Error: unexpected symbol in "sample estimates"
> mean of x
Error: unexpected symbol in "mean of"
> 0.4564444
[1] 0.4564444
> #pvalue<alpha=0.5
> #Reject H0
> #There is sufficient evidence that the mean level of Salmonella is greater
> #than 0.3
> #Suppose that 10 volunteers have taken an intelligence test. Here are the
```

```
> #results obtained. The average score of entire population is 75 in the same
> #test. Is there any significant difference between the sample and population
> #means, assuming that the variance of the population is not known.
> #Scores : 65,78,88,55,48,95,66,57,79,81
> #H0:mu=75
> #H1:mu is not equal to 75
> x=c(65,78,88,55,48,95,66,57,79,81)
> t.test(x,mu=75,alternative="two.sided")
```

One Sample t-test

```
data: x
t = -0.78303, df = 9, p-value = 0.4537
alternative hypothesis: true mean is not equal to 75
95 percent confidence interval:
 60.22187 82.17813
sample estimates:
mean of x
 71.2
```

```
> One Sample t-test
Error: unexpected symbol in " One Sample"
> data: x
Error in data:x : NA/NaN argument
In addition: Warning message:
In data:x : numerical expression has 10 elements: only the first used
> t = -0.78303, df = 9, p-value = 0.4537
Error: unexpected ',' in "t = -0.78303,"
> R Console Page 2
Error: unexpected symbol in "R Console"
> alternative hypothesis: true mean is not equal to 75
Error: unexpected symbol in "alternative hypothesis"
> 95 percent confidence interval:
Error: unexpected symbol in "95 percent"
> 60.22187 82.17813
Error: unexpected numeric constant in " 60.22187 82.17813"
> sample estimates:
Error: unexpected symbol in "sample estimates"
> mean of x
Error: unexpected symbol in "mean of"
> 71.2
[1] 71.2
> #pvaluealpha=0.05
> #Accept H0: There is no significant difference between sample mean and
> #population mean.
> #HW
> #One problem from Mayer book or some other sources
> #try left tailed test.
> mu=71.2
> sd(x)
[1] 15.34637
> [1] 15.34637
Error: unexpected '[' in "["
> x1<-rnorm(10,mean=71.2,sd=15.34637)
> x1
[1] 81.19945 83.04234 69.32159 52.49277 74.70743 56.13057 61.64727
[8] 80.39532 104.94237 72.73213
> [1] 75.96110 78.54377 85.72119 73.84867 96.69831 61.18057 75.62806
Error: unexpected '[' in "["
> [8] 108.39588 61.83122 84.77760
Error: unexpected '[' in "["
> t.test(x1,mu=75,alternative="two.sided")
```

One Sample t-test

```
data: x1
t = -0.27789, df = 9, p-value = 0.7874
alternative hypothesis: true mean is not equal to 75
95 percent confidence interval:
 62.76190 84.56035
sample estimates:
mean of x
 73.66112
```

```

> One Sample t-test
Error: unexpected symbol in " One Sample"
> data: x1
Error in data:x1 : NA/NaN argument
In addition: Warning message:
In data:x1 : numerical expression has 10 elements: only the first used
> t = 1.1446, df = 9, p-value = 0.2819
Error: unexpected ',' in "t = 1.1446,"
> alternative hypothesis: true mean is not equal to 75
Error: unexpected symbol in "alternative hypothesis"
> 95 percent confidence interval:
Error: unexpected symbol in "95 percent"
> 69.86554 90.65174
Error: unexpected numeric constant in " 69.86554 90.65174"
> sample estimates:
Error: unexpected symbol in "sample estimates"
> mean of x
Error: unexpected symbol in "mean of"
> 80.25864
[1] 80.25864
> #large health clinic and we are testing a new drug, Procardia,
> #whose work is to reduce hypertension. We find 13000
> #individuals with high systolic blood pressure ( $\bar{x}$ =150 mmHg, SD=10 mmHg)
> #and we provide them Procardia for a month, and then measure their blood pressure again.
> #We find that the mean systolic blood pressure has decreased
> #to 144 mmHg with a standard deviation of 9 mmHg.
> set.seed(2820)
> pre_Treatment <- c(rnorm(2000, mean = 150, sd = 10))
> post_Treatment <- c(rnorm(2000, mean = 144, sd = 9))
> t.test(pre_Treatment, post_Treatment, paired = TRUE)

      Paired t-test

data:  pre_Treatment and post_Treatment
t = 20.789, df = 1999, p-value < 2.2e-16
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 5.593910 6.759259
sample estimates:
mean of the differences
      6.176585

>

```

```

> #Experiment - 11
> #Date 8/12/2020
> #Slot Lab L1
> #Name: Amlan Nayak
> #Reg.No:19BCD7143
> #course: Applied statistics
> #Code: MAT1011
> #Problem 1
> #A school atheletics has taken a new instructor,want to test the effectiveness
> #of the new type of training proposed by hte new instructor comparing the
> #average time of 10 runners in the 100 meters.The results are given below.
> #(Time in seconds)
> #Before training
> #12.9 13.5 12.8 15.6 17.2 19.2 12.6 15.3 14.4 11.3
> #After training:
> #12.7 13.6 12.0 15.2 16.8 20.0 12.0 15.9 16.0 11.1
> # test the hypothesis that there is significant difference between before
> # and aftrt training
>
> x=c(12.9,13.5,12.8,15.6,17.2,19.2,12.6,15.3,14.4,11.3)
> y=c(12.7,13.6,12,15.2,16.8,20,12,15.9,16,11.1)
> #Null hypothesis H0:mu1=mu2
> #H1:mu1 is not equal to mu2
> t.test(x,y,mu=0,alternative="two.sided")

```

Welch Two Sample t-test

```

data:  x and y
t = -0.043323, df = 17.653, p-value = 0.9659
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 -2.478112  2.378112
sample estimates:
mean of x mean of y
 14.48    14.53

> t.test(x,y,mu=0,paired=TRUE,alternative="two.sided")

```

Paired t-test

```

data:  x and y
t = -0.21331, df = 9, p-value = 0.8358
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 -0.5802549  0.4802549
sample estimates:
mean of the differences
 -0.05

```

```

> #pvalue>alpha=0.05 , Accept H0 there is no improvement
>
> #Problem 2
> #Suppose now that manager of the team fired the coach and take another
> #promising. We report the times of atheletes after the second training.
> #Before training:
> #12.9,13.5,12.8,15.6,17.2,19.2,12.6,15.3,14.4,11.3
> #After training:
> #12.0,12.2,11.2,13.0,15.0,15.8,12.2,13.4,12.9,11.0
> #test the hypothesis that there is an improvement in training
> #Null hypothesis H0: mu1-mu2<=0
> #Alternative hypothesis : mu1-mu2>0
> x=c(12.9,13.5,12.8,15.6,17.2,19.2,12.6,15.3,14.4,11.3)
> y=c(12.0,12.2,11.2,13.0,15.0,15.8,12.2,13.4,12.9,11.0)
> #Right tailed test
> t.test(x,y,paired=TRUE,alternative="greater")

```

Paired t-test

```

data:  x and y
t = 5.2671, df = 9, p-value = 0.0002579

```

```

alternative hypothesis: true difference in means is greater than 0
95 percent confidence interval:
 1.049675      Inf
sample estimates:
mean of the differences
      1.61

```

```

> #pvalue=0.0002579<alpha=0.05=Area of rejection region
> #reject H0: there is an improvement in training
> #Left tailed test problem
> #Consider the paired data below that represents cholesterol levels in 10 men
> #before and after a certain medication.
> #Before (x)
> #237,289,257,228,303,275,262,304,244,233
> #After (y)
> #194,240,230,186,265,222,242,281,240,212
> # Test the claim that on average , the drug may not lower the cholesterol
> #in all men.
> #Test this at 5% level of significance
> #Null hypothesis H0:mu1>=mu2
> #Alternative hypothesis H1:mu1<=mu2
> #Left tailed test
> x=c(237,289,257,228,303,275,262,304,244,233)
> y=c(194,240,230,186,265,222,242,281,240,212)
> t.test(x,y,paired=TRUE,alternative="less")

```

Paired t-test

```

data:  x and y
t = 6.5594, df = 9, p-value = 0.9999
alternative hypothesis: true difference in means is less than 0
95 percent confidence interval:
 -Inf 40.94289
sample estimates:
mean of the differences
      32

```

```

> #pvalue>alpha=0.05 Accept H0 Medication lower the cholesterol
> #Experiment - 11 F-test
> #Try on problem paired t-test from mayer book
> #rnorm(n,mean=,sd=) generate n random samples
> #values for each sample
> #x=rnorm(n,mean=,sd=)
> #y=rnorm(n,mean=,sd=)
> #Five measurements of output of two units have given the following
> #results(in kilpgrams per one hour) of operation.Assume that both the samples
> #have been obtained from normal operations,test at 10% significance level if two
> #populations have same variance
> #UNIT A: 14.1 10.1 14.7 13.7 14.0
> #UNIT B: 14.0 14.5 13.7 12.7 14.1
> #H0:var1=var2
> #H1:var1 != var2
> x=c(14.1,10.1,14.7,13.7,14)
> y=c(14,14.5,13.7,12.7,14.1)
> var.test(x,y,alternative="two.sided")

```

F test to compare two variances

```

data:  x and y
F = 7.3304, num df = 4, denom df = 4, p-value = 0.07954
alternative hypothesis: true ratio of variances is not equal to 1
95 percent confidence interval:
 0.7632268 70.4053799
sample estimates:
ratio of variances
      7.330435

```

```

> #Alpha=0.1
> #pvalue<alpha
> #Reject H0 population variances are not equal

```

```
> #At 5% level pvalue=0.07954>0.05=alpha Accept H0  
> #Take on application problem from Mayer book on f-test  
> #Related to right tailed test  
>
```

```
R version 4.0.2 (2020-06-22) -- "Taking Off Again"
Copyright (C) 2020 The R Foundation for Statistical Computing
Platform: x86_64-w64-mingw32/x64 (64-bit)
```

```
R is free software and comes with ABSOLUTELY NO WARRANTY.
You are welcome to redistribute it under certain conditions.
Type 'license()' or 'licence()' for distribution details.
```

```
Natural language support but running in an English locale
```

```
R is a collaborative project with many contributors.
Type 'contributors()' for more information and
'citation()' on how to cite R or R packages in publications.
```

```
Type 'demo()' for some demos, 'help()' for on-line help, or
'help.start()' for an HTML browser interface to help.
Type 'q()' to quit R.
```

```
[Previously saved workspace restored]
```

```
> # Taking two random samples
> A = c(16, 17, 25, 26, 32,
+       34, 38, 40, 42)
> B = c(600, 590, 590, 630, 610, 630)
> # var test in R
> var.test(A, B, alternative = "two.sided")
```

```
F test to compare two variances
```

```
data: A and B
F = 0.27252, num df = 8, denom df = 5, p-value = 0.1012
alternative hypothesis: true ratio of variances is not equal to 1
95 percent confidence interval:
 0.04033118 1.31282683
sample estimates:
ratio of variances
 0.2725248
```

```
>
```

```
R version 4.0.3 (2020-10-10) -- "Bunny-Wunnies Freak Out"
Copyright (C) 2020 The R Foundation for Statistical Computing
Platform: i386-w64-mingw32/i386 (32-bit)
```

```
R is free software and comes with ABSOLUTELY NO WARRANTY.
You are welcome to redistribute it under certain conditions.
Type 'license()' or 'licence()' for distribution details.
```

```
Natural language support but running in an English locale
```

```
R is a collaborative project with many contributors.
Type 'contributors()' for more information and
'citation()' on how to cite R or R packages in publications.
```

```
Type 'demo()' for some demos, 'help()' for on-line help, or
'help.start()' for an HTML browser interface to help.
Type 'q()' to quit R.
```

```
[Previously saved workspace restored]
```

```
> #experiment 12
> #slot LAb L1
> #Date 12 th Dec 2020
> #name Amlan Nayak
> #REGD NO:19BCD7143
> #The below table gives the distribution of students
> #according to the family type and anxiety level
> #Family type Anxiety level
> # Low Normal High
> #
> #          Low      Normal      High
> #Joint Family      35      42      61
> #Nuclear family    48      51      68
> #test the null hypothesis that the family type
> #and the anxiety level are independent
> data<-matrix(c(35,42,61,48,51,68),ncol=3,byrow=T)
> data
      [,1] [,2] [,3]
[1,]   35   42   61
[2,]   48   51   68
> data<-matrix(c(35,42,61,48,51,68),ncol=3,nrow=2,byrow=T)
> data
      [,1] [,2] [,3]
[1,]   35   42   61
[2,]   48   51   68
> chisq.test(data)
```

```
Pearson's Chi-squared test
```

```
data: data
X-squared = 0.53441, df = 2, p-value = 0.7655
```

```
> #pvalue> alpha =0.05 Accept H0:Family type and anxiety
> #Level are independent
> #problem 2
> #test he null hpothesis that the type of drug and age grown
> #are independent for the following data from the excel
> data<-read.csv("C:\\Users\\HP\\Downloads\\chidata.csv")
> data
  PatientID Drug Age.Group
1         P1 Drug A Under 21
2         P2 Drug A Under 21
3         P3 Drug A Under 21
4         P4 Drug A Under 21
5         P5 Drug A 21 to 55
6         P6 Drug A 21 to 55
7         P7 Drug A 21 to 55
8         P8 Drug A 21 to 55
9         P9 Drug A Over 55
10        P10 Drug A Over 55
```



```
11      P11 Drug A    Over 55
12      P12 Drug A    Over 55
13      P13 Drug B    Under 21
14      P14 Drug B    Under 21
15      P15 Drug B    Under 21
16      P16 Drug B    Under 21
17      P17 Drug B    21 to 55
18      P18 Drug B    21 to 55
19      P19 Drug B    21 to 55
20      P20 Drug B    21 to 55
21      P21 Drug B    Over 55
22      P22 Drug B    Over 55
23      P23 Drug B    Over 55
24      P24 Drug B    Over 55
25      P25 Drug C    Under 21
26      P26 Drug C    Under 21
27      P27 Drug C    Under 21
28      P28 Drug C    Under 21
29      P29 Drug C    21 to 55
30      P30 Drug C    21 to 55
31      P31 Drug C    21 to 55
32      P32 Drug C    21 to 55
33      P33 Drug C    Over 55
34      P34 Drug C    Over 55
35      P35 Drug C    Over 55
36      P36 Drug C    Over 55
37          Drug A    21 to 55
38          Drug A    21 to 55
39          Drug A    21 to 55
40          Drug A    21 to 55
41          Drug A    Over 55
42          Drug A    Over 55
43          Drug A    Over 55
44          Drug A    Over 55
45          Drug A    Over 55
46          Drug A    Over 55
47          Drug A    Over 55
48          Drug A    Over 55
49          Drug B    Under 21
50          Drug B    Under 21
51          Drug B    Under 21
52          Drug B    Under 21
53          Drug B    Over 55
54          Drug B    Over 55
55          Drug B    Over 55
56          Drug B    Over 55
57          Drug B    Over 55
58          Drug B    Over 55
59          Drug B    Over 55
60          Drug B    Over 55
61          Drug B    Over 55
62          Drug B    Over 55
63          Drug B    Over 55
64          Drug B    Over 55
65          Drug B    Over 55
66          Drug B    Over 55
67          Drug C    Over 55
68          Drug C    Over 55
69          Drug C    Over 55
70          Drug C    Over 55
71          Drug C    Over 55
72          Drug C    Over 55
73          Drug C    Under 21
74          Drug C    Under 21
75          Drug C    Under 21
76          Drug C    Under 21
77          Drug C    Under 21
78          Drug C    Under 21
79          Drug C    Under 21
80          Drug C    Under 21
```

```
81      Drug C Under 21
82      Drug C Under 21
83      Drug C Under 21
84      Drug C Under 21
85      Drug C Under 21
86      Drug C Under 21
87      Drug C Under 21
88      Drug C Under 21
89      Drug C Under 21
90      Drug C Under 21
91      Drug C Under 21
92      Drug C Under 21
93      Drug C Over 55
94      Drug A Under 21
95      Drug A Under 21
96      Drug A Under 21
97      Drug A Under 21
98      Drug A Under 21
99      Drug A Under 21
100     Drug A Under 21
101     Drug A Under 21
102     Drug A Under 21
103     Drug A Under 21
104     Drug A Under 21
105     Drug A Under 21
106     Drug A Under 21
107     Drug A Under 21
108     Drug A Under 21
109     Drug A Under 21
110     Drug A Under 21
111     Drug A Under 21
112     Drug B 21 to 55
113     Drug B 21 to 55
114     Drug B 21 to 55
115     Drug B 21 to 55
116     Drug B 21 to 55
117     Drug B 21 to 55
118     Drug B 21 to 55
119     Drug B 21 to 55
120     Drug B 21 to 55
121     Drug B 21 to 55
122     Drug B 21 to 55
123     Drug B 21 to 55
124     Drug B 21 to 55
125     Drug B 21 to 55
126     Drug B 21 to 55
127     Drug B 21 to 55
128     Drug B 21 to 55
129     Drug B 21 to 55
130     Drug B 21 to 55
131     Drug B 21 to 55
132     Drug B 21 to 55
133     Drug B 21 to 55
134     Drug B 21 to 55
135     Drug B 21 to 55
136     Drug B 21 to 55
137     Drug B 21 to 55
138     Drug B 21 to 55
139     Drug B 21 to 55
140     Drug B 21 to 55
141     Drug B 21 to 55
142     Drug C 21 to 55
143     Drug C 21 to 55
144     Drug C 21 to 55
145     Drug C 21 to 55
146     Drug C 21 to 55
147     Drug C 21 to 55
148     Drug C 21 to 55
149     Drug C 21 to 55
150     Drug C 21 to 55
```

```

151      Drug C  21 to 55
152      Drug C  21 to 55
153      Drug C  21 to 55
154      Drug C  21 to 55
155      Drug C  21 to 55
156      Drug C  21 to 55
157      Drug C  21 to 55
158      Drug C  21 to 55
159      Drug C  21 to 55
160      Drug C  21 to 55
161      Drug C  21 to 55
162      Drug C  21 to 55
163      Drug C  21 to 55
164      Drug C  21 to 55
165      Drug C  21 to 55
166      Drug C  21 to 55
167      Drug C  21 to 55
> tab2<-table(data$Drug,data$Age.Group)
> tab2

      21 to 55 Over 55 Under 21
Drug A      8     12      22
Drug B     34     18       8
Drug C     30     11      24
> chisq.test(tab2)

      Pearson's Chi-squared test

data:  tab2
X-squared = 23.223, df = 4, p-value = 0.0001143

> #pvalue=0.0001143<alpha=0.05 Reject H0
> #type of drug and age group are dependent
> #fitting of binomial distribution
> #A surver of 320 families with 5 children each
> #revealed the following distribution
> #no of boys :5      4      3      2      1      0
> #no of girls:0      1      2      3      4      5
> #is this result consistent with the hypothesis
> #the male and female births are equally possible.
> n=5
> N=320
> x=0:5
> obsf<-c(14,56,110,88,40,12)
> p=0.5
> x=0:5
> exf<-dbinom(x,n,p)*320
> exf
[1] 10 50 100 100 50 10
> chisq<-sum((obsf-exf)^2/exf)
> chisq
[1] 7.16
> qchisq(0.95,5)
[1] 11.0705
> #caluculated value<tabulated value
> #Accept H0 male and female births are equal
> #Fitting of Poisson distribution
> #Fit the Poisson distribution and test the godness of fit
> #x:0 1 2 3 4 5 6
> #f:275 72 30 7 5 2 1
> x<-c(275,72,30,7,5,2,1)
> lambda<-sum(f*x)/sum(f)
Error: object 'f' not found
> f<-c(275 72 30 7 5 2 10
Error: unexpected numeric constant in "f<-c(275 72"
> f<-c(275 72 30 7 5 2 10 )
Error: unexpected numeric constant in "f<-c(275 72"
> f<-c(275 ,72, 30, 7, 5, 2, 10 )
> lambda<-sum(f*x)/sum(f)
> lambda

```

```
[1] 203.9825
> exf<-dpois(x,lambda)*sum(f)
> exf
[1] 1.407119e-04 3.298767e-24 7.564518e-50 3.015586e-74 3.043933e-77
[6] 2.151827e-82 2.109814e-84
> exf=round(exf)
> exf1=round(exf)
> obsf<-c(275,72,30,15)
> exf2<-c(242,117,28,6)
> chisq<-sum((obsf-exf2)^2/exf)
Warning message:
In (obsf - exf2)^2/exf :
  longer object length is not a multiple of shorter object length
> exf2
[1] 242 117 28 6
> obsf1
Error: object 'obsf1' not found
>
> chisq<-sum((obsf-exf2)^2/exf2)
> chisq
[1] 35.45055
> qchisq(0.95,2)
[1] 5.991465
> #caluculated value >tabulated value
> #caluculated value >tabulated value Reject HO
> #poission distribution is not a good fit
> #Fit the normal distribution.
```