

In [?], Lipyanskiy uses a different notion of co-orientation from the one we have used to define geometric cochains. We here discuss Lipyanskiy's co-orientation, which he initially refers to as orientations of maps, and show that when we have smooth maps $f : M \rightarrow N$, it is equivalent to our definition, up to possible sign conventions. In other words, we show that a map is co-orientable in our sense if and only if it is co-orientable in Lipyanskiy's sense. We will not explore the precise differences between the specific co-orientation conventions.

To define, co-orientations, Lipyanskiy utilizes the determinant line bundles of Donaldson and Kronheimer in [?, Section 5.2.1]. A key point throughout our discussion will be the following lemma, which is presented without proof in [?]:

Lemma 0.1. *Given an exact sequence of vector bundles*

$$0 \rightarrow V_1 \rightarrow \cdots \rightarrow V_m \rightarrow 0,$$

there is a canonical isomorphism

$$\bigotimes_{i \text{ odd}} \text{Det}(V_i) \cong \bigotimes_{i \text{ even}} \text{Det}(V_i).$$

Proof. UGH - TRY TO FIND THIS SOMEWHERE We first consider the case of a short exact sequence $0 \rightarrow V_1 \xrightarrow{d_1} V_2 \xrightarrow{d_2} V_3 \rightarrow 0$. As short exact sequences of vector bundles always split (REF!), we have a map $e : V_3 \rightarrow V_2$ such that $d_1 \oplus e : V_1 \oplus V_3 \rightarrow V_2$ is an isomorphism. So $\text{Det}(V_1 \oplus V_3) \xrightarrow{\text{Det}(d_1 \oplus e)} \text{Det}(V_2)$ is an isomorphism, as is the canonical map (REF OR PROVE) $\text{Det}(V_1) \otimes \text{Det}(V_3) \rightarrow \text{Det}(V_1 \oplus V_3 \rightarrow V_2)$. The composite isomorphism appears *a priori* to depend on the splitting e , but if e' is another such splitting then the image of $e - e'$ lies in $\ker(d_2) = \text{im}(d_1)$, so $\text{Det}(d_1 \otimes e) - \text{Det}(d_1 \otimes e') = \text{Det}(d_1 \otimes (e - e')) = 0$ as the exterior product of a top form of $\text{im}(d_1)$ with anything else in $\text{im}(d_1)$ must be 0. REF TO THESE MECHANICS So the isomorphism is independent of the choice of splitting. THIS PROOF NEEDS A LOT OF FIXING. \square

We can now define the determinant line bundles as in [?, Section 5.2.1]. Donaldson and Kronheimer work in a more general setting, but we will confine ourselves to considering a map of vector bundles $F : E \rightarrow E'$ over M . At first, we also assume that $\ker(F)$ and $\text{cok}(F)$ are well-defined vector bundles. Then, in our notation from D: det bundle, the Donaldson-Kronheimer determinant line bundle is defined to be

$$\text{Det}(\ker(F)) \otimes \text{Det}(\text{cok}(F))^*,$$

where the $*$ over $\text{Det}(\text{cok}(F))$ denotes the dual bundle. Below we will consider that $\ker(F)$ and $\text{cok}(F)$ are not always vector bundles, but for now we see that the determinant bundle is morally related to the index of an operator. We refer to [?, Section 5.2.1] for a more precise statement of the relationship.

To relate the Donaldson-Kronheimer determinant line bundle to our notion of co-orientation, consider the exact sequence of vector bundles

$$0 \rightarrow \ker(F) \rightarrow E \rightarrow E' \rightarrow \text{cok}(F) \rightarrow 0.$$

Applying L: det sequence, we have $\text{Det}(\ker(F)) \otimes \text{Det}(E') \cong \text{Det}(E) \otimes \text{Det}(\text{cok}(F))$. Next we use that for a line bundle L we have $L \otimes L^* \cong \underline{\mathbb{R}}$, the trivial line bundle. So multiplying both sides by $\text{Det}(\text{cok}(F))^*$ and $\text{Det}(E')^*$, we get

$$\text{Det}(\ker(F)) \otimes \text{Det}(\text{cok}(F))^* \cong \text{Det}(E) \otimes \text{Det}(E')^*.$$

The latter is isomorphic to $\text{Hom}(\text{Det}(E'), \text{Det}(E))$, which is dual to $\text{Hom}(E, E')$. In particular, $\text{Hom}(E, E')$ is trivial, and so admits a non-zero section, if and only if the Donaldson-Kronheimer determinant bundle $\text{Det}(\ker(F)) \otimes \text{Det}(\text{cok}(F))^*$ is trivial.

In the setting of a smooth map $f : M \rightarrow N$, we can think of the derivative Df as a map $Df : TM \rightarrow f^*(TN)$, and then the above demonstrates that $\text{Hom}(TM, f^*(TN))$ is trivial if and only if the determinant bundle $\text{Det}(\ker(dF)) \otimes \text{Det}(\text{cok}(dF))^*$ is trivial. We recall that the triviality of $\text{Hom}(TM, f^*(TN))$ is the condition for co-orientability of f in the sense of D: co-orientations. So, up to the actual co-orientation conventions, the two notions of co-orientability coincide. We leave it to the reader to define the isomorphisms in sufficient detail to carry a particular co-orientation as defined in S: co-orientations to one as defined here.

The problem with the preceding analysis is that in general $\ker(Df)$ and $\text{cok}(Df)$ do not necessarily have the same dimensions from fiber to fiber, and so $\ker(Df)$ and $\text{cok}(Df)$ are not necessarily well defined as vector bundles. The solution is to reframe the definition of the determinant line bundle as in [?] so that it is always well defined and such that it is isomorphic to $\text{Det}(\ker(F)) \otimes \text{Det}(\text{cok}(F))^*$ when it is also well defined.

For this, let $\underline{\mathbb{R}}^n$ be the trivial \mathbb{R}^n bundle over M , and suppose we have a map $\psi : \underline{\mathbb{R}}^n \rightarrow E'$ such that $F \oplus \psi : E \oplus \underline{\mathbb{R}}^n$ is surjective¹. This will always be true in our setting, as we observed in Section ?? that work of Joyce and Melrose implies that smooth manifolds with corners can always be embedded in finite dimensional Euclidean space. Hence tangent bundles are subbundles of trivial bundles and so the images of projections of trivial bundles (or, up to isomorphism, quotients of the trivial bundle by their orthogonal complements after endowing the trivial bundle with a Riemannian structure). The gain is that $F \oplus \psi$ now has trivial cokernel and a kernel that is a vector bundle, as now the fibers of the kernel have a fixed dimension. We then define the determinant line bundle to be

$$\mathcal{L} = \text{Det}(\ker(F \oplus \psi)) \otimes \text{Det}(\underline{\mathbb{R}}^n)^* \cong \text{Det}(\ker(F \oplus \psi)).$$

In the case where $\ker(F)$ and $\text{cok}(F)$ were already vector bundles, \mathcal{L} is isomorphic to the determinant line bundle of F using L: det sequence and the following lemma:

Lemma 0.2. *If $F : E \rightarrow E'$ and $\psi : \underline{\mathbb{R}}^n \rightarrow E'$ are bundle maps with $F \oplus \psi : E \oplus \underline{\mathbb{R}}^n$ surjective and $\ker(F)$ and $\text{cok}(F)$ well-defined vector bundles, then the following sequence is exact²:*

$$0 \longrightarrow \ker(F) \longrightarrow \ker(F \oplus \psi) \longrightarrow \underline{\mathbb{R}}^n \longrightarrow \text{cok}(F) \longrightarrow 0.$$

¹Donaldson and Kronheimer work with complex vector bundles, so [?] features $\underline{\mathbb{C}}^n$ rather than $\underline{\mathbb{R}}^n$.

²This exact sequence appears incorrectly in [?] with the ψ in place of F in the first and last terms.

Proof. This exact sequence is simply the serpent lemma exact sequence obtained from the commutative diagram of exact sequences

$$\begin{array}{ccccccccc}
0 & \longrightarrow & E & \longrightarrow & E \oplus \underline{\mathbb{R}}^n & \longrightarrow & \underline{\mathbb{R}}^n & \longrightarrow & 0 \\
& & \downarrow F & & \downarrow F \oplus \psi & & \downarrow & & \\
0 & \longrightarrow & E' & \xrightarrow{=} & E' & \longrightarrow & 0 & \longrightarrow & 0.
\end{array}$$

The category of vector bundles over a space is not technically an abelian category, but one can check by hand for this diagram that, with our assumptions, all the maps of the exact sequence are well defined and the exactness then holds fiberwise by the classical serpent lemma. In particular, the map $\underline{\mathbb{R}}^n \rightarrow \text{cok}(F)$ is the composition of the splitting map $\underline{\mathbb{R}}^n \rightarrow E \oplus \underline{\mathbb{R}}^n$, the map $F \oplus \psi$, and the projection E' to $\text{cok}(F)$. \square

Combining this lemma with L: det sequence gives us an isomorphism

$$\text{Det}(\ker(F)) \otimes \text{Det}(\underline{\mathbb{R}}^n) \cong \text{Det}(\ker(F \oplus \psi)) \otimes \text{Det}(\text{cok}(F)).$$

Multiplying both sides by $\text{Det}(\underline{\mathbb{R}}^n)^* \otimes \text{Det}(\text{cok}(F))^*$ and using again that for a line bundle L we have $L \otimes L^* \cong \underline{\mathbb{R}}$, we obtain

$$\text{Det}(\ker(F)) \otimes \text{Det}(\text{cok}(F))^* \cong \text{Det}(\ker(F \oplus \psi)) \otimes \text{Det}(\underline{\mathbb{R}}^n)^*.$$

So, as promised, the two definitions agree (up to canonical isomorphisms) when $\ker(F)$ and $\text{cok}(F)$ are defined.

Finally, we note that the definition of \mathcal{L} is clearly independent of n up to canonical isomorphism as $\det(\underline{\mathbb{R}}^n) \cong \det(\underline{\mathbb{R}}^n)^* \cong \underline{\mathbb{R}}$ for all n (for the isomorphisms to be canonical, we may assume $\underline{\mathbb{R}}^n$ to be given its standard oriented basis and standard inner product).