KINKAJOU

Potos flavus · Rainforest canopy — southern Mexico to Brazil
Frugivore · Wild-range diet

3-SECOND CHECK Do I see fruit — dominating the bowl?

THE RULE What they would naturally come across in their wild range, by percentage of natural diet. Nothing else.
Mid- to upper-canopy — fruit dominates (especially figs); nectar, flowers, occasional insects supplement.

≈90% BASE always visible in the bowl

≈10% TREATS hand-fed or scattered — never in the bowl

BASE
Wild figs
primary canopy staple

TREAT
Nectar flowers
important pollinator role

TREAT
Mixed fruit
mango, banana, papaya

TREAT
Flowers
rainforest blooms

TREAT
Occasional insects
minor supplement

Plus other plants and prey from their wild range — pesticide-free only.

NO PROCESSED SUGAR No candy, no processed sugar, no dairy. Whole or chunked fruit must dominate.

HOW WE FEED

- Base must be visible in the bowl.
- Treats are hand-fed or scattered, never in the bowl.
- Baseline before guest hours — do not feed to fullness before guests arrive.
- All items pesticide- and herbicide-free.

MATCH THE BOWL — DON'T INTERPRET.

If unsure: Pause → Ask → Escalate. No substitutions without approval.

What a living thing can sense becomes its reality. Environment shapes design.