RED RUFFED LEMUR

Varecia rubra · Rainforest canopy — northeast Madagascar
Frugivore · Wild-range diet

3-SECOND CHECK Do I see fruit — dominating the bowl?

THE RULE What they would naturally come across in their wild range, by percentage of natural diet. Nothing else.
Rainforest canopy — ripe fruit dominates (75-95%); nectar, leaves, seeds supplement.

≈90% **BASE** always visible in the bowl

≈10% **TREATS** hand-fed or scattered — never in the bowl

BASE
Ripe rainforest fruit
figs, native fruits

TREAT
Wild figs
favored canopy fruit

TREAT
Nectar flowers
Traveler's palm, fig

TREAT
Canopy leaves
dry-season fallback

TREAT
Young shoots
tender new growth

Plus other plants and prey from their wild range — pesticide-free only.

NO ORANGES No oranges, no citrus, no romaine-heavy bowls. Whole or chunked fruit must dominate.

HOW WE FEED

- Base must be visible in the bowl.
- Treats are hand-fed or scattered, never in the bowl.
- Baseline before guest hours — do not feed to fullness before guests arrive.
- All items pesticide- and herbicide-free.

MATCH THE BOWL — DON'T INTERPRET.

If unsure: Pause → Ask → Escalate. No substitutions without approval.

What a living thing can sense becomes its reality. Environment shapes design.