PREHENSILE-TAILED PORCUPINE

Coendou prehensilis · Rainforest canopy — northern South America
Root + leaf herbivore · Wild-range diet

3-SECOND CHECK Do I see leaves AND a root in the bowl?

THE RULE What they would naturally come across in their wild range, by percentage of natural diet. Nothing else.
Rainforest canopy — leaves and stems dominate; a visible root is required; flowers and fruit supplement.

≈90% **BASE** always visible in the bowl

≈10% **TREATS** hand-fed or scattered — never in the bowl

BASE
Canopy leaves & stems
primary forage

TREAT
Root (required)
yam or sweet potato

TREAT
Flowers & shoots
tender new growth

TREAT
Wild figs
seasonal fruit

TREAT
Guava
occasional native fruit

Plus other plants and prey from their wild range — pesticide-free only.

ROOT IS REQUIRED If there is no visible root, the bowl is wrong. No salad-only, no pellet-only.

HOW WE FEED

- Base must be visible in the bowl.
- Treats are hand-fed or scattered, never in the bowl.
- Baseline before guest hours — do not feed to fullness before guests arrive.
- All items pesticide- and herbicide-free.

MATCH THE BOWL — DON'T INTERPRET.

If unsure: Pause → Ask → Escalate. No substitutions without approval.

What a living thing can sense becomes its reality. Environment shapes design.