[photo]

# KEEL-BILLED TOUCAN

Ramphastos sulfuratus

**FRUGIVORE — Low Iron, Lower Sugar**

WHAT THIS ANIMAL IS: Fruit + low iron + sugar-sensitive

SITES: Houston

**3-SEC CHECK** Do I see berries/pear — not banana?

## STANDARD DIET — HEALTHY ANIMALS ONLY

Excludes sick / vet-care / special-diet animals

Covers ~95% of daily feedings. For the remaining 5%, use training and observation within sound science. Unsure — Pause. Ask. Escalate.

### MORNING FEED-OUT

Fruit in 2" chunks, lower-sugar bias (berries/apple/pear lead). Softbill chow in separate bowl. Fruit visible and dominant.

APPROX DAILY: ~120–180 g fruit + chow (keeper posts exact portion)

### ENCOUNTER / ENRICHMENT PORTION

1/2" cubes of approved lower-sugar fruits for tong delivery. Never hand-feed banana or mango — even a small piece can cause loose stool.

## CORRECT

Fruit dominant · softbill · lower sugar

Berries

Apple

[photo]

Pear

Melon

Papaya

Leafy greens

Low-iron softbill chow

## WRONG — FIX IMMEDIATELY

High-sugar fruit heavy · Iron-rich foods · Pellet heavy

Banana

Mango

Orange

Herbivore pellets

Primate chow

### SIMPLE BUILD

- 60–70% fruit (lower-sugar bias)
- Low-iron softbill chow
- Small leafy greens portion

### PREP RULE

Chunked fruit (2" pieces). Low-iron softbill chow only. Limit banana/mango/papaya to trace amounts — these cause loose stool in keel-billed.

**MATCH THE BOWL — DO NOT INTERPRET.**

No photo = feeding not complete. · No substitutions without approval. · Unsure → Pause → Ask → Escalate.

Source: DIETS-steward-export PART 7 §1C (frugivore low-iron) + Mia Ketsa field note (keel-billed sugar sensitivity) · Body-text extraction
LIFE Diet Governance System · v2026.04.23 · Match the bowl. Don't interpret.