RED RUFFED LEMUR

FRUGIVORE — Fruit Dominant

WHAT THIS ANIMAL IS: Fruit specialist — Madagascar rainforest

SITES: Houston · SAA · Other

3-SEC CHECK Is fruit dominant?

STANDARD DIET — HEALTHY ANIMALS ONLY

Excludes sick / vet-care / special-diet animals

Covers ~95% of daily feedings. For the remaining 5%, use training and observation within sound science. Unsure — Pause. Ask. Escalate.

MORNING FEED-OUT

Fruit in 2-3 inch chunks. Softbill/primate chow as prescribed.

APPROX DAILY: Per posted portion chart

ENCOUNTER / ENRICHMENT PORTION

Bite-sized cubes (1/2 inch) for hand or tong delivery. Part of daily allocation, not extra.

CORRECT

Fruit dominant · leaves · small chow

Mango

Papaya

Berries

Melon

Kale

Primate chow

WRONG — FIX IMMEDIATELY

Oranges (not natural) · Romaine · Starch vegetables

Orange

Romaine

Sweet potato / yam

Carrot

SIMPLE BUILD

- Fruit dominant
- leaves secondary
- small chow

PREP RULE

Chunked fruit. Whole leaves. No citrus.

MATCH THE BOWL — DO NOT INTERPRET.

No photo = feeding not complete. · No substitutions without approval. · Unsure → Pause → Ask → Escalate.