BLACK & WHITE RUFFED LEMUR

Varecia variegata · Rainforest canopy — eastern Madagascar
Frugivore · Wild-range diet

3-SECOND CHECK Do I see fruit — dominating the bowl?

THE RULE What they would naturally come across in their wild range, by percentage of natural diet. Nothing else.
Rainforest canopy — ripe fruit dominates (≈92%); nectar, pollen, leaves, seeds supplement.

≈90% BASE always visible in the bowl

≈10% TREATS hand-fed or scattered — never in the bowl

BASE
Ripe rainforest fruit
figs, native fruits

TREAT
Wild figs
favored canopy fruit

TREAT
Nectar & pollen
flowers as nectar source

TREAT
Canopy leaves
dry-season fallback

TREAT
Wild berries
seasonal fruit

Plus other plants and prey from their wild range — pesticide-free only.

NO ORANGES No oranges, no citrus, no romaine-heavy bowls. Whole or chunked fruit must dominate.

HOW WE FEED

- Base must be visible in the bowl.
- Treats are hand-fed or scattered, never in the bowl.
- Baseline before guest hours — do not feed to fullness before guests arrive.
- All items pesticide- and herbicide-free.

MATCH THE BOWL — DON'T INTERPRET.

If unsure: Pause → Ask → Escalate. No substitutions without approval.

What a living thing can sense becomes its reality. Environment shapes design.