MONKEY-TAILED SKINK

Corucia zebrata · Solomon Islands rainforest canopy
Folivore · Wild-range diet

3-SECOND CHECK Do I see leaves — dominating the bowl?

THE RULE What they would naturally come across in their wild range, by percentage of natural diet. Nothing else.
Rainforest canopy — leaves of Ficus and Epipremnum dominate; flowers, shoots, and fruit supplement.

≈90% BASE always visible in the bowl

≈10% TREATS hand-fed or scattered — never in the bowl

BASE
Epipremnum / Ficus leaves
their main wild diet

TREAT
Strangler fig leaves
Ficus species

TREAT
Flowers
canopy blooms

TREAT
Young shoots
tender new growth

TREAT
Wild figs
occasional fruit

Plus other plants and prey from their wild range — pesticide-free only.

NO CHOPPED SALAD No chopped salad bowl, no fruit-dominant bowls. Whole leaves dominate.

HOW WE FEED

- Base must be visible in the bowl.
- Treats are hand-fed or scattered, never in the bowl.
- Baseline before guest hours — do not feed to fullness before guests arrive.
- All items pesticide- and herbicide-free.

MATCH THE BOWL — DON'T INTERPRET.

If unsure: Pause → Ask → Escalate. No substitutions without approval.

What a living thing can sense becomes its reality. Environment shapes design.