TWO-TOED SLOTH

Choloepus didactylus · Rainforest canopy — northern South America
Folivore · Wild-range diet

3-SECOND CHECK Do I see leaves — dominating the bowl?

THE RULE What they would naturally come across in their wild range, by percentage of natural diet. Nothing else.
Rainforest canopy — leaves dominate; tender shoots, blossoms, and occasional fruit supplement.

≈90% **BASE** always visible in the bowl

≈10% **TREATS** hand-fed or scattered — never in the bowl

BASE
Canopy leaves
Cecropia, mixed browse

TREAT
Young shoots
tender new growth

TREAT
Blossoms
rainforest flowers

TREAT
Wild figs
seasonal fruit

TREAT
Mixed browse
varied canopy leaves

Plus other plants and prey from their wild range — pesticide-free only.

NO CHOPPED SALAD No chopped salad bowl, no fruit-dominant bowls. Whole leaves dominate.

HOW WE FEED

- Base must be visible in the bowl.
- Treats are hand-fed or scattered, never in the bowl.
- Baseline before guest hours — do not feed to fullness before guests arrive.
- All items pesticide- and herbicide-free.

MATCH THE BOWL — DON'T INTERPRET.

If unsure: Pause → Ask → Escalate. No substitutions without approval.

What a living thing can sense becomes its reality. Environment shapes design.