BENNETT'S WALLABY

Notamacropus rufogriseus · Tasmania and southeastern Australia
Grazer · Wild-range diet

3-SECOND CHECK Do I see grass / hay — dominating the bowl?

THE RULE What they would naturally come across in their wild range, by percentage of natural diet. Nothing else.
Eucalyptus forest edge and grasslands — grasses dominate (≈74%); herbs, juicy roots, leaves supplement.

≈90% BASE always visible in the bowl

≈10% TREATS hand-fed or scattered — never in the bowl

BASE
Grass hay
timothy, orchard, meadow

TREAT
Fresh grass
primary wild forage

TREAT
Herbs & forbs
broadleaf plants

TREAT
Juicy roots
dry-season water source

TREAT
Young shoots
tender new growth

Plus other plants and prey from their wild range — pesticide-free only.

NO STARCHY VEG No corn, peas, beans, or carrot-heavy bowls. Grass or hay must always be visible.

HOW WE FEED

- Base must be visible in the bowl.
- Treats are hand-fed or scattered, never in the bowl.
- Baseline before guest hours — do not feed to fullness before guests arrive.
- All items pesticide- and herbicide-free.

MATCH THE BOWL — DON'T INTERPRET.

If unsure: Pause → Ask → Escalate. No substitutions without approval.

What a living thing can sense becomes its reality. Environment shapes design.