

Hands-on practice on the Reaction4Exp platform





Manuela Rodríguez-Gallardo
Antonio Moro Mario Gómez-Ramos
Carla Muñoz-Chimbo

8th July 2025

Elastic and Inelastic scattering

Hands-on practice
on the
Reaction4Exp
platform



Elastic scattering

Hands-on practice
on the
Reaction4Exp
platform

Woods-Saxon parameters for $^4\text{He} + ^{208}\text{Pb}$ optical potential

V_0 [MeV]	r_0 [fm]	a_0 [fm]	W_v [MeV]	r_i [fm]	a_i [fm]	r_c [fm]
96.44	1.085	0.625	32	0.958	0.42	1.2

Remember: absolute (physical) radii $R_x = r_x(A_p^{1/3} + A_t^{1/3})$.

Elastic scattering

Hands-on practice
on the
Reaction4Exp
platform

Woods-Saxon parameters for $^4\text{He} + ^{208}\text{Pb}$ optical potential

V_0 [MeV]	r_0 [fm]	a_0 [fm]	W_v [MeV]	r_i [fm]	a_i [fm]	r_c [fm]
96.44	1.085	0.625	32	0.958	0.42	1.2





Remember: absolute (physical) radii $R_x = r_x(A_p^{1/3} + A_t^{1/3})$.

🔍 We will try four incident energies:

E_{lab} (MeV)	E_{cm} (MeV)	k (fm ⁻¹)	η	r_{min} (fm)
10	9.81	1.36	16.3	24.1
22	21.6	2.01	11.0	10.9
27	26.5	2.23	9.94	8.9
60	58.9	3.32	6.67	4.0





Elastic scattering cross sections

Hands-on practice
on the
Reaction4Exp
platform



Elastic S-matrices

Hands-on practice
on the
Reaction4Exp
platform



Inelastic scattering

Hands-on practice
on the
Reaction4Exp
platform

Woods-Saxon parameters for ${}^4\text{He} + {}^{208}\text{Pb}(3^-)$ at 23.5 MeV

V_0	r_0	a_0	W_s	r_i	a_i	r_c
[MeV]	[fm]	[fm]	[MeV]	[fm]	[fm]	[fm]
92.5	1.384	0.625	22.24	1.265	0.592	1.2

Nuclear and Coulomb deformation parameters

β_N	β_C
0.103	0.113

Inelastic scattering

Hands-on practice
on the
Reaction4Exp
platform

Woods-Saxon parameters for $^4\text{He} + ^{208}\text{Pb}(3^-)$ at 23.5 MeV

V_0	r_0	a_0	W_s	r_i	a_i	r_c
[MeV]	[fm]	[fm]	[MeV]	[fm]	[fm]	[fm]
92.5	1.384	0.625	22.24	1.265	0.592	1.2

Nuclear and Coulomb deformation parameters


β_N	β_C
0.103	0.113

$$\delta_N = \beta_N R_N = 0.85 \text{ fm}$$

$$M_n(E3) = \beta_C 3 Ze R_C^3 / 4\pi = 795.08 e \text{ fm}^3$$

Elastic and inelastic cross sections using CC method


Hands-on practice
on the
Reaction4Exp
platform





EPM: inelastic and Coulomb breakup

Hands-on practice
on the
Reaction4Exp
platform

$^{11}\text{Be} + ^{197}\text{Au}$ at 39.6 and 31.9 MeV



$^{11}\text{Be} + ^{197}\text{Pb}$ at 69 MeV/u



Inelastic scattering to the excited bound state of ^{11}Be

Hands-on practice
on the
Reaction4Exp
platform

$^{11}\text{Be} + ^{197}\text{Au}$ at 39.6 and 31.9 MeV

$$B(E1; gs \rightarrow 1/2^-) = 0.116 \text{ e}^2\text{fm}^2$$

Remember: distribution type "discrete"



Inelastic scattering to the excited bound state of ^{11}Be

Hands-on practice
on the
Reaction4Exp
platform

$^{11}\text{Be} + ^{197}\text{Au}$ at 39.6 and 31.9 MeV

$$B(E1; gs \rightarrow 1/2^-) = 0.116 \text{ e}^2\text{fm}^2$$

Remember: distribution type "discrete"



Coulomb breakup for ^{11}Be

Hands-on practice
on the
Reaction4Exp
platform

$^{11}\text{Be} + ^{208}\text{Pb}$ at 69 MeV/u

We need a $dB(E1)/d\varepsilon$ distribution from a theor. model

Remember: distribution type "continous"


Coulomb breakup for ^{11}Be

Hands-on practice
on the
Reaction4Exp
platform

$^{11}\text{Be} + ^{208}\text{Pb}$ at 69 MeV/u

We need a $dB(E1)/d\varepsilon$ distribution from a theor. model

Remember: distribution type "continuous"




Coulomb breakup for ^{11}Be

Hands-on practice
on the
Reaction4Exp
platform

$^{11}\text{Be} + ^{208}\text{Pb}$ at 69 MeV/u


We need a $dB(E1)/d\varepsilon$ distribution from a theor. model

Remember: distribution type "continuous"



Transfer reaction

Hands-on practice
on the
Reaction4Exp
platform



Potentials needed

Hands-on practice
on the
Reaction4Exp
platform

For n-p ground state we use the Gaussian form:


$$V_{pn}(r) = -72.15 \exp[-(r/1.484)^2]$$

system	V_0 [MeV]	r_0 [fm]	a_0 [fm]	W_s [MeV]	r_i [fm]	a_i [fm]	r_c [fm]
$d+^{208}\text{Pb}$	112.0	1.25	0.682	19.4	1.25	0.783	1.30
$p+^{208(9)}\text{Pb}$	52.0	1.25	0.65	10.0	1.25	0.76	1.25
$n+^{208}\text{Pb}(\text{gs})$	adjust.	1.23	0.65				

Remember: $Q = Q_0 - E_x$

Transfer cross sections to several states of ^{209}Pb

Hands-on practice
on the
Reaction4Exp
platform



$$\theta_{\max} \approx \arcsin \left(\frac{\sqrt{\ell(\ell+1)}\hbar}{pR} \right)$$

Discussion and feedback on the platform

Hands-on practice
on the
Reaction4Exp
platform

- With Reaction4Exp we can do calculations for elastic, inelastic, transfer and Coulomb breakup reactions as seen.
- Soon, breakup using CDCC will be available.
- Also available it is the obtention of double-folding optical potentials using the SPP code.

Discussion and feedback on the platform

Hands-on practice
on the
Reaction4Exp
platform

- With Reaction4Exp we can do calculations for elastic, inelastic, transfer and Coulomb breakup reactions as seen.
- Soon, breakup using CDCC will be available.
- Also available it is the obtention of double-folding optical potentials using the SPP code.
- Feedback & suggestions



Hands-on practice
on the
Reaction4Exp
platform

To finish...