

CLINICAL CASE STUDY REPORT

AI Case Study Generator | Generated: 16 May 2026 19:03

PATIENT INFORMATION

Age: 55

Gender: Female

Marital Status: N/A

Occupation: N/A

Address: N/A

Mode of Admission: N/A

PRESENTING SYMPTOMS / CHIEF COMPLAINT

- The patient is a 55-year-old female presenting with right-sided neck pain and stiffness for the last 1 month. The pain aggravates with neck rotation and looking up, and is relieved by resting and hot pack application. There is no history of trauma or sudden jerk. The patient's past medical history reveals mild hypertension.
- On examination, the patient is conscious and hemodynamically stable. The vitals and systems review are unremarkable. The posture is observed to be within normal limits, although the patient exhibits tenderness over the right lower cervical paraspinal muscles on palpation.
- The musculoskeletal and neuro assessment reveals that the neck range of motion is restricted in extension and right rotation due to pain. However, the neuro assessment is intact, with no numbness or pain radiating to the arms, and reflexes are normal. The gait is observed to be normal.

PAIN ASSESSMENT (NUMERIC RATING SCALE 0-10)

Pain Intensity:  **6 / 10**

TREATMENT APPLIED

- The treatment applied during the session included heat therapy to relieve pain and stiffness, gentle cervical mobilizations to improve range of motion, and neck isometric exercises to strengthen the cervical muscles.

CLINICAL RECOMMENDATIONS & FOLLOW-UP

- The clinical impression is that the patient is suffering from Cervical Spondylosis with muscle spasm, which is causing the right-sided neck pain and stiffness.
- The ongoing treatment plan includes continued heat therapy, gentle cervical mobilizations, and neck isometric exercises. The patient is advised to continue with the home exercises and follow-up with the physiotherapist for further assessment and progression of the treatment plan.

Date of session: 16 May 2026

Physiotherapist Signature

DISCLAIMER: This document was generated by an AI assistant to assist clinical documentation. It must be reviewed and countersigned by the responsible physiotherapist before use as an official clinical record.