

# Singular Spacetimes: A Unified Computational and Theoretical Treatment of Black Holes, Wormholes, and White Holes

Amna Fatima

July 2026

## 1 Overview of Black Holes, Wormholes, and White Holes

### 1.1 Black Holes

A black hole is a region of spacetime where gravity is so extreme that nothing—not even light—can escape beyond the event horizon. Black holes emerge as exact solutions to Einstein’s field equations and represent the most extreme curvature of spacetime known in General Relativity.

### 1.2 White Holes

A white hole is the time-reversed counterpart of a black hole—mathematically valid but physically mysterious. Whereas nothing can escape from a black hole, nothing can enter a white hole. White holes appear naturally in the maximally extended Schwarzschild solution, yet they have never been observed.

### 1.3 Wormholes

A wormhole is a hypothetical tunnel connecting two separate regions of spacetime. The original Einstein–Rosen bridge connects a black hole to a white hole, making all three objects part of the same geometric family.

### 1.4 Why Study Them Together?

These three objects are not separate phenomena. They are different manifestations of the same mathematical framework—all arising as exact solutions to Einstein’s field equations and as consequences of extreme spacetime curvature. Studying them together provides a deeper understanding than studying each object in isolation.

This project is divided into two main parts: **Observed Physics** and **Theoretical Physics**.

### 1.5 A Brief Overview

Before understanding black holes, wormholes, and white holes, it is important to understand gravity itself. The first successful mathematical description of gravity was proposed by Isaac Newton, forming the foundation upon which later theories were built.

## 2 Newton’s Law of Gravitation

Newton stated that every object in the Universe attracts every other object with a gravitational force. The strength of this force depends on the masses of the objects and the distance between them.

$$F = G \frac{m_1 m_2}{r^2}$$

where

- $F$  = gravitational force
- $G$  = gravitational constant

- $m_1$  = mass of the first object
- $m_2$  = mass of the second object
- $r$  = distance between the centres of the two objects

The equation shows that:

- Greater mass produces a stronger gravitational force. For example, Earth exerts a stronger gravitational pull than the Moon because it has a much greater mass.
- Greater distance weakens the gravitational force. If the distance between two objects is doubled, the gravitational force becomes one-quarter of its original value because of the inverse-square dependence on  $r^2$ .

## 2.1 Why Is It Not Enough?

Although Newton's law accurately explains gravity in most everyday situations, it cannot fully describe extremely strong gravitational phenomena such as black holes or the behaviour of spacetime in their vicinity. Furthermore, Newton treated gravity as a force acting between masses.

Einstein later introduced the theory of General Relativity, which describes gravity as the curvature of spacetime rather than as a conventional force. Since black holes are predicted by General Relativity, and wormholes and white holes arise as theoretical solutions of Einstein's field equations, General Relativity provides the appropriate theoretical foundation for this project.

## 3 Einstein's General Relativity

While Newton described gravity as a force acting between masses, Einstein explained gravity not as a fundamental force but as the curvature of spacetime caused by mass and energy.

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}$$

where

- $G_{\mu\nu}$  = Einstein tensor, representing the curvature of spacetime.
- $\Lambda$  = Cosmological constant, associated with the energy density of empty space (dark energy).
- $g_{\mu\nu}$  = Metric tensor, describing the geometry of spacetime.
- $G$  = Newton's gravitational constant.
- $c$  = Speed of light in a vacuum.
- $T_{\mu\nu}$  = Stress-energy tensor, representing the distribution of matter and energy.

In simple terms, Einstein's field equations state:

- Matter tells spacetime how to curve.
- Curved spacetime tells matter how to move.
- Massive objects such as planets and stars bend spacetime, and that curvature determines the motion of other objects.

### 3.1 Connection

This new understanding of gravity enabled Einstein's theory to predict black holes and provided the mathematical framework from which wormholes and white holes also arise. Therefore, General Relativity forms the theoretical foundation of this project.

## 4 Observed Physics

### 4.1 Overview

This section focuses on black holes because they are predicted by General Relativity and supported by strong observational evidence. It covers:

1. Formation of black holes
2. Types of black holes
3. Structure of black holes
4. Observational evidence

### 4.2 Formation of Black Holes

Black holes form through the gravitational collapse of massive stars. Every star exists in a continuous balance between gravity, which pulls matter inward, and pressure, which pushes outward. This outward pressure is generated by nuclear fusion in the stellar core, where enormous amounts of energy are released, preventing the star from collapsing under its own gravity.

A star initially fuses hydrogen into helium and later continues fusing progressively heavier elements. Iron is unique because, unlike the earlier stages of fusion, fusing iron does not release energy; instead, it requires energy. Once the core becomes primarily composed of iron, the star can no longer generate enough energy to support itself against gravity. The outward pressure rapidly decreases, the core begins to collapse, and gravity compresses the matter into an increasingly smaller volume.

The final outcome depends on the mass of the collapsing core. If the core is relatively small, it becomes a white dwarf. If it is more massive, it forms a neutron star. However, if the core exceeds the maximum mass that neutron degeneracy pressure can support, gravity becomes so strong that no known force can halt the collapse. The core continues collapsing until it becomes so compact that not even light can escape. At this stage, an event horizon forms.

According to General Relativity, the collapse continues toward a singularity, where the current theory is expected to break down. General Relativity can be viewed as the mathematical framework in which black holes, wormholes, and white holes emerge as different solutions to Einstein's field equations. As an enormous amount of mass is compressed into an extremely small region, spacetime becomes increasingly curved, ultimately leading to the formation of a black hole.

During the collapse, the outer layers of the star fall inward and then rebound violently, producing an explosion known as a **core-collapse supernova**, while the remaining stellar core becomes a black hole.




Figure 1: Simplified sequence of black hole formation from the gravitational collapse of a massive star. After nuclear fusion ceases at an iron core, gravity causes the core to collapse. The resulting core-collapse supernova ejects the outer layers of the star, while the remnant core forms a black hole if its mass exceeds the maximum supported by neutron degeneracy pressure.

### 4.3 Astrophysical Classification

#### 4.3.1 Stellar-Mass Black Holes

Stellar-mass black holes form when the core of a massive star collapses under its own gravity. If the core is sufficiently massive, gravity overwhelms all known forms of pressure, resulting in the formation of a black hole. If the remaining core is less massive, it instead becomes a neutron star.

Some stellar-mass black holes can become significantly more massive if their parent stars have low metallicity. Such stars lose less mass through stellar winds before collapsing, leaving behind heavier black holes.

Their masses typically range from approximately **3 to 100 times the mass of the Sun**.

#### 4.3.2 Intermediate-Mass Black Holes

Intermediate-mass black holes occupy the mass range between stellar-mass and supermassive black holes. Although several candidates have been identified, strong observational evidence for their existence remains limited.

Several formation mechanisms have been proposed:

- **Runaway Collisions:** In dense star clusters, repeated collisions between massive stars produce an extremely massive star, which eventually collapses into an intermediate-mass black hole.
- **Black Hole Mergers:** Numerous stellar-mass black holes gradually merge over millions of years, producing a larger black hole.
- **Direct Collapse:** A massive cloud of gas in the early Universe may collapse directly into a black hole without first forming a star.

Their masses are estimated to lie between approximately **100 and 100,000 times the mass of the Sun**.

Intermediate-mass black holes are important because they may represent the missing evolutionary link between stellar-mass and supermassive black holes. However, observational evidence for their existence is still limited.

#### 4.3.3 Supermassive Black Holes

Almost every large galaxy is believed to contain a supermassive black hole at its centre. For example, **Sagittarius A\*** resides at the centre of the Milky Way, while **M87\*** became the first black hole to be directly imaged by the Event Horizon Telescope in 2019.

Scientists are still investigating how supermassive black holes formed. Two main formation models have been proposed.

**Light Seed Model** A relatively small black hole forms from the collapse of a massive early-generation star. Over billions of years, it grows by:

- Accreting gas and dust from its surroundings.
- Merging with other black holes.

**Heavy Seed Model** Instead of forming from a star, an enormous cloud of primordial gas collapses directly into a massive black hole, providing a much larger initial seed.

Supermassive black holes have masses ranging from approximately **100,000 to 10 billion times the mass of the Sun**.

They play a crucial role in the evolution of galaxies. Observations show that the mass of a galaxy's central black hole is closely correlated with the size and motion of stars in the galactic bulge, suggesting that galaxies and their central black holes evolve together.

**Observational Evidence** Evidence for supermassive black holes includes:

- Observations of stars orbiting **Sagittarius A\***.
- Powerful quasars and active galactic nuclei.
- Images obtained by the **Event Horizon Telescope**.

#### 4.3.4 Primordial Black Holes

Unlike other black holes, primordial black holes did not form from stars. Instead, they may have formed just fractions of a second after the Big Bang.

At that time, the Universe was incredibly hot and dense. Some regions may have been much denser than others. If one of these dense regions had enough gravity, it could collapse directly into a black hole.

They are important because, according to scientists, they might explain the mysterious nature of dark matter, how supermassive black holes formed so early in the Universe, and, if very small primordial black holes exist, they may eventually evaporate through Hawking radiation. Detecting such evaporation would provide evidence for Hawking radiation because very small primordial black holes would slowly evaporate over time.

Their masses could range from extremely tiny values to thousands of times the mass of the Sun, depending on when they formed.

No primordial black holes have been confirmed. They remain a theoretical possibility rather than a confirmed type of black hole.

Among these classifications, stellar-mass and supermassive black holes have strong observational evidence, whereas intermediate-mass and primordial black holes remain active areas of research.

### 4.4 General Relativity Classification

#### 4.4.1 Schwarzschild Black Hole (1916)

The Schwarzschild black hole was the first exact solution to Einstein's gravitational field equations, found by Karl Schwarzschild in 1916. It provides the mathematical description of spacetime around a spherical, non-rotating, uncharged mass.

Schwarzschild showed that every non-rotating black hole has a single event horizon. The solution predicts that time slows down near a black hole. The closer you are to a massive object such as the Sun, Earth, or a black hole, the more slowly time flows compared with someone farther away. This phenomenon is called **gravitational time dilation**.

The solution also predicts that light follows **null geodesics** because spacetime is curved. As a result, light appears to bend around massive objects such as stars. This prediction was confirmed during the 1919 solar eclipse.

When the Schwarzschild solution is examined, a special radius naturally appears. This is called the **Schwarzschild radius**. If all the mass of an object is compressed within this radius, an event horizon forms. In this way, black holes naturally emerge from the mathematics of General Relativity.

#### 4.4.2 The Schwarzschild Metric

The Schwarzschild metric is

$$ds^2 = -\left(1 - \frac{2GM}{rc^2}\right) c^2 dt^2 + \left(1 - \frac{2GM}{rc^2}\right)^{-1} dr^2 + r^2 d\Omega^2.$$

Every term in the metric has a physical meaning.

#### 4.4.3 The Time Component

The first term is

$$-\left(1 - \frac{2GM}{rc^2}\right) c^2 dt^2.$$

This term determines how time flows in the gravitational field.

Far from the black hole,

$$r \rightarrow \infty,$$

so

$$1 - \frac{2GM}{rc^2} \rightarrow 1.$$

Time flows at its normal rate for a distant observer.

## Schwarzschild Spacetime




Figure 2: Spherically symmetric geometry of a Schwarzschild black hole. Surfaces of constant radial coordinate  $r$  are concentric spheres centred on the black hole, reflecting the spherical symmetry of the Schwarzschild solution.

As one moves closer to the black hole, the quantity

$$1 - \frac{2GM}{rc^2}$$

decreases. Consequently, time passes more slowly compared with that experienced by a distant observer.

At the Schwarzschild radius,

$$r = \frac{2GM}{c^2},$$

the coefficient becomes zero. To a distant observer, a clock at the event horizon appears to slow down indefinitely and never quite crosses the horizon. This phenomenon is known as **gravitational time dilation**.

### 4.4.4 The Radial Component

The radial term is

$$\left(1 - \frac{2GM}{rc^2}\right)^{-1} dr^2.$$

This term determines how radial distances are measured in the gravitational field.

As one approaches the event horizon,

$$r \rightarrow \frac{2GM}{c^2},$$

the factor

$$\left(1 - \frac{2GM}{rc^2}\right)^{-1}$$

grows without bound. Consequently, the radial distance between nearby points becomes greatly stretched. This effect reflects the strong curvature of spacetime near the event horizon.

#### 4.4.5 The Angular Component

The angular part of the Schwarzschild metric is

$$r^2 (d\theta^2 + \sin^2 \theta d\phi^2).$$

This term describes the geometry of a sphere of radius  $r$ . It shows that every surface of constant  $r$  remains a perfect sphere, demonstrating that Schwarzschild spacetime possesses **spherical symmetry**.




Figure 3: Illustration of gravitational time dilation. Clocks nearer the Schwarzschild event horizon run more slowly relative to clocks located far from the black hole, as observed by a distant observer.

#### 4.4.6 Reissner–Nordström Black Hole (1916–1918)

The Reissner–Nordström black hole is the solution to Einstein’s field equations for a non-rotating, electrically charged black hole. It extends the Schwarzschild solution by introducing electric charge.

It is characterized by only two quantities:

- Mass ( $M$ )
- Electric charge ( $Q$ )

This solution was needed because the Schwarzschild solution assumes that the black hole possesses mass only, with no electric charge or rotation. However, Einstein’s equations also allow matter to possess electric charge. When the electromagnetic field is included through the Einstein–Maxwell equations, the field equations become

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}^{(\text{EM})},$$

where  $T_{\mu\nu}^{(\text{EM})}$  is the electromagnetic stress–energy tensor.

Outside the black hole, no ordinary matter exists, but the electromagnetic field stores energy. That energy contributes to the curvature of spacetime.

Unlike the Schwarzschild solution, the Reissner–Nordström solution contains two horizons:

- Outer event horizon
- Inner Cauchy horizon

If the electric charge becomes too large, the horizons disappear mathematically, leaving a naked singularity.

Unlike the Schwarzschild black hole, whose singularity is always in the future of any observer crossing the event horizon, the Reissner–Nordström singularity is timelike. This means the singularity is not necessarily the unavoidable final destination of every observer.

#### 4.4.7 Kerr Black Hole (1963)

Unlike the Schwarzschild solution, Kerr black holes are believed to represent what nature actually produces because astrophysical objects retain their angular momentum as they collapse. As a result, most black holes are expected to rotate.

## Reissner–Nordström Black Hole




Figure 4: Schematic structure of a Reissner–Nordström black hole. In contrast to the Schwarzschild solution, a charged black hole possesses two horizons: an outer event horizon and an inner Cauchy horizon. The electric field outside the black hole contributes to the curvature of spacetime through the Einstein–Maxwell equations.

### 4.4.8 Spin Parameter

Unlike a Schwarzschild black hole, a Kerr black hole rotates about an axis and is characterized by two fundamental quantities:

- Mass ( $M$ )
- Angular momentum ( $J$ )

One of the most important predictions of General Relativity is that a spinning black hole drags the surrounding spacetime with it. This phenomenon is known as **frame dragging**, or the **Lense–Thirring effect**. Objects near the black hole cannot remain completely stationary; they are forced to rotate along with the surrounding spacetime.

The spin parameter is defined as

$$a = \frac{J}{Mc},$$

where

- $a$  = spin parameter (angular momentum per unit mass),
- $J$  = angular momentum of the black hole,
- $M$  = mass of the black hole,
- $c$  = speed of light in a vacuum.

A larger value of  $a$  corresponds to a more rapidly rotating black hole. In geometrized units ( $G = c = 1$ ),

$$a \leq M.$$

If

$$a > M,$$

no event horizon exists, and the solution describes a naked singularity rather than a black hole.



#### 4.4.9 Event Horizon

The event horizon is the boundary beyond which nothing, not even light, can escape.

For a Kerr black hole, the radius of the outer event horizon is

$$r_+ = \frac{GM}{c^2} + \sqrt{\left(\frac{GM}{c^2}\right)^2 - a^2}.$$

where

- $r_+$  = radius of the outer event horizon,
- $G$  = gravitational constant,
- $M$  = mass of the black hole,
- $c$  = speed of light in a vacuum,
- $a$  = spin parameter, defined as  $a = \frac{J}{Mc}$ .

As the spin parameter  $a$  increases, the radius of the event horizon decreases. For an extremal Kerr black hole, where

$$a = \frac{GM}{c^2},$$

the inner and outer horizons coincide.

The Kerr black hole also possesses a unique region called the **ergosphere**, which lies outside the event horizon.

Unlike the Schwarzschild solution, whose singularity is treated as a point, the Kerr solution predicts a **ring singularity**. Its radius is equal to the spin parameter  $a$ . This ring singularity arises because the black hole is rotating.

Kerr black holes are important because most real black holes in the Universe are expected to rotate, making the Kerr solution the most realistic description of astrophysical black holes.




Figure 5: Schematic structure of a Kerr black hole. Rotation gives rise to an ergosphere outside the event horizon, where frame dragging forces nearby objects to co-rotate with the black hole. The central singularity takes the form of a ring rather than a point, distinguishing the Kerr solution from the Schwarzschild solution.

#### 4.4.10 Kerr–Newman Black Hole (1965)

The Kerr–Newman black hole is the most general stationary black hole solution of Einstein’s field equations with electromagnetism included. It describes a black hole with three physical properties:

- Mass ( $M$ )
- Angular momentum (rotation) ( $J$ )

- Electric charge ( $Q$ )

It combines all previous black hole solutions. In 1965, Ezra Newman and his colleagues extended the Kerr solution to include electric charge, producing the Kerr–Newman metric.

Its physical characteristics include:

- **Mass ( $M$ ):** Determines the strength of the gravitational field.
- **Spin ( $J$ ):** The black hole rotates, dragging spacetime around it through frame dragging.
- **Electric charge ( $Q$ ):** Creates an electromagnetic field around the black hole.

Unlike the Schwarzschild or Kerr solutions, the Kerr–Newman solution satisfies the Einstein–Maxwell equations because the black hole possesses both gravity and electromagnetism.

#### 4.4.11 Kerr–Newman Metric

In Boyer–Lindquist coordinates, the Kerr–Newman metric is

$$\begin{aligned}
 ds^2 = & - \left( 1 - \frac{2Mr - Q^2}{\Sigma} \right) c^2 dt^2 - \frac{2a(2Mr - Q^2) \sin^2 \theta}{\Sigma} c dt d\phi \\
 & + \frac{\Sigma}{\Delta} dr^2 + \Sigma d\theta^2 \\
 & + \left( r^2 + a^2 + \frac{(2Mr - Q^2)a^2 \sin^2 \theta}{\Sigma} \right) \sin^2 \theta d\phi^2.
 \end{aligned}$$

where

$$\Sigma = r^2 + a^2 \cos^2 \theta,$$

$$\Delta = r^2 - 2Mr + a^2 + Q^2,$$

and

$$a = \frac{J}{Mc}.$$

The quantities are defined as:

- $M$  = mass of the black hole,
- $J$  = angular momentum of the black hole,
- $Q$  = electric charge,
- $a$  = spin parameter (angular momentum per unit mass),
- $c$  = speed of light,
- $\Sigma$  and  $\Delta$  = auxiliary functions that simplify the metric.

## 4.5 Horizons

The horizon locations are

$$r_{\pm} = M \pm \sqrt{M^2 - a^2 - Q^2}.$$

where

- $r_+$  = outer event horizon,
- $r_-$  = inner Cauchy horizon.

A Kerr–Newman black hole exists only if

$$M^2 \geq a^2 + Q^2.$$

If

$$M^2 < a^2 + Q^2,$$

there is no event horizon, resulting in a **naked singularity**. According to the **Cosmic Censorship Conjecture**, such naked singularities are generally not expected to occur in nature.

Instead of a point singularity, the Kerr–Newman black hole contains a ring singularity located at

$$r = 0, \quad \theta = \frac{\pi}{2}.$$

Because the black hole is charged and rotating, it generates:

- An electric field.
- A magnetic field produced by the rotating charge.

However, astrophysical black holes are expected to be nearly electrically neutral because any net charge is quickly neutralized by surrounding accreting plasma.




Figure 6: Schematic structure of a Kerr–Newman black hole. In addition to mass and rotation, the black hole carries electric charge, producing an electromagnetic field. The solution contains an ergosphere, an outer event horizon, an inner Cauchy horizon, and a ring singularity.

#### 4.6 Why Do These Four Solutions Exist?

Einstein’s field equations are

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}.$$

The left-hand side,

$$G_{\mu\nu},$$

describes the curvature of spacetime.

The right-hand side,

$$T_{\mu\nu},$$

describes the distribution of matter and energy.

Different assumptions about matter, energy, and symmetry lead to different solutions of Einstein’s field equations. For example:

- Vacuum ( $T_{\mu\nu} = 0$ )  $\rightarrow$  Schwarzschild and Kerr black holes.
- Electrovacuum ( $T_{\mu\nu} = T_{\mu\nu}^{(\text{EM})}$ )  $\rightarrow$  Reissner–Nordström and Kerr–Newman black holes.

#### 4.6.1 Vacuum Solutions

Outside a star or black hole, there is essentially no matter or radiation. Therefore, the stress–energy tensor vanishes:

$$T_{\mu\nu} = 0.$$

The Einstein field equations then reduce to the vacuum field equations:

$$R_{\mu\nu} = 0.$$

where

- $R_{\mu\nu}$  = Ricci curvature tensor,
- $T_{\mu\nu}$  = stress–energy tensor.

Although the stress–energy tensor is zero, gravity does not disappear. The curvature of spacetime produced by the central mass remains, allowing gravity to exist even in empty space.

These vacuum field equations give rise to two important black hole solutions:

- **Schwarzschild solution** – describes a non-rotating, uncharged black hole.
- **Kerr solution** – describes a rotating, uncharged black hole.

#### 4.6.2 Electrovacuum Solutions

Suppose the black hole possesses a net electric charge.

Outside the black hole, there is no ordinary matter, but an electromagnetic field exists. The electromagnetic field carries energy and momentum, which contribute to the curvature of spacetime.

Therefore, the stress–energy tensor is no longer zero:

$$T_{\mu\nu}^{(\text{EM})} \neq 0.$$

Einstein’s field equations become

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}^{(\text{EM})},$$

where

- $G_{\mu\nu}$  = Einstein tensor,
- $T_{\mu\nu}^{(\text{EM})}$  = electromagnetic stress–energy tensor,
- $G$  = Newton’s gravitational constant,
- $c$  = speed of light in a vacuum.

The Einstein field equations must also be solved together with Maxwell’s equations, which describe the electromagnetic field. These coupled equations are known as the **Einstein–Maxwell equations**.

The Einstein–Maxwell equations give rise to two important charged black hole solutions:

- **Reissner–Nordström solution** – describes a non-rotating, charged black hole.
- **Kerr–Newman solution** – describes a rotating, charged black hole.

#### 4.6.3 Why Only Mass, Charge, and Spin?

Imagine two stars: one is blue, hot, magnetic, turbulent, and irregular, while the other is cold, red, and chemically different. After gravitational collapse, if they produce black holes with the same mass, charge, and angular momentum, they become identical.

Everything else disappears behind the event horizon.

This idea is summarized by the statement that black holes have **no hair**. The theorem states that any stationary black hole is completely described by only three quantities:

$$(M, Q, J).$$

#### 4.6.4 No-Hair Theorem

A stationary black hole in four-dimensional General Relativity is completely described by only three externally measurable quantities:

- **Mass ( $M$ ):** Determines how strongly spacetime is curved.
- **Electric Charge ( $Q$ ):** If the black hole carries electric charge, it produces an electric field outside the event horizon.
- **Angular Momentum ( $J$ ):** If the black hole rotates, it drags spacetime around it through frame dragging.

During gravitational collapse, the newly formed black hole is highly disturbed.

It emits gravitational waves and, if charged, electromagnetic radiation. This process causes the black hole to “ring down”, carrying away irregularities until it settles into a stable state.

This process is sometimes compared to striking a bell:

Immediately after the strike, the sound is complex. After a short time, only a few simple tones remain.

Similarly, after collapse, only three quantities remain measurable:

- Mass ( $M$ )
- Charge ( $Q$ )
- Spin ( $J$ )

Everything else has either fallen behind the event horizon or been radiated away.




Figure 7: Illustration of the No-Hair Theorem. A complex collapsing star emits gravitational radiation during the ringdown phase and settles into a stationary black hole completely characterised by only three externally measurable quantities: mass ( $M$ ), electric charge ( $Q$ ), and angular momentum ( $J$ ).

## 4.7 Anatomy of a Black Hole

### 4.7.1 Event Horizon – The Point of No Return

The event horizon is not a physical wall or a solid surface. It is an imaginary boundary in spacetime.

Once anything crosses this boundary, light, matter, or any form of information cannot escape. Everything inside becomes disconnected from the outside universe.

It is called the “point of no return” because near a black hole, gravity curves spacetime so severely that every possible future path points inward.

The event horizon marks the boundary where escape becomes impossible.

### 4.7.2 Photon Sphere

Imagine a photon. Normally, light always travels in straight lines through flat spacetime. Near a black hole, spacetime itself is curved. The photon follows the straightest possible path in that curved geometry—a null geodesic. At exactly

$$r=3M$$

gravity bends light just enough that it travels around the black hole in a circular orbit. This is called the photon sphere (for Schwarzschild). This unstable orbit is why the photon sphere helps determine the size of a black hole's observed shadow.

Kerr photon region change the picture due to rotation. Frame dragging helps photons moving with the spin (prograde) and hinders those moving against it (retrograde). Instead of a single spherical photon orbit, rotating black holes have a more complicated photon region containing different unstable light orbits. This asymmetry contributes to the slightly "D"-shaped appearance expected for rapidly spinning black holes.

### 4.7.3 Ergosphere

This exists only for rotating black holes. The black hole spins. Because mass curves spacetime, rotating mass twists spacetime. Imagine stirring coffee. The liquid near the spoon rotates. Farther away, rotation weakens. Now replace coffee with spacetime. Everything nearby is dragged around. Static limit

Outside the event horizon there exists a surface where

$$g_{tt} = 0.$$

This surface is called the **static limit** (or **stationary limit**).

The region between the static limit and the event horizon is known as the **ergosphere**.

Within the ergosphere, the intense rotation of the black hole drags spacetime around with it, a phenomenon known as **frame dragging**. As a result, no object can remain stationary with respect to a distant observer, although it is still possible to escape from this region.

The outer boundary of the ergosphere is located at

$$r_{\text{static}} = \frac{GM}{c^2} + \sqrt{\left(\frac{GM}{c^2}\right)^2 - a^2 \cos^2 \theta},$$

where

- $r_{\text{static}}$  = radius of the static limit,
- $G$  = gravitational constant,
- $M$  = mass of the black hole,
- $c$  = speed of light,
- $a = \frac{J}{Mc}$  = spin parameter,
- $\theta$  = polar angle.

Unlike event horizon, escape from ergosphere is possible because it lies outside the event horizon. Although the black hole's rotation drags spacetime so strongly that everything inside the ergosphere is forced to rotate with it, there are still paths that lead outward. Light and matter can still escape if they move in the right direction. Only after crossing the event horizon are all possible future paths directed inward, making escape impossible.

### 4.7.4 Singularity

A singularity is not a point made of infinitely dense matter in the ordinary sense. Mathematically, it is a place where the equations of General Relativity predict quantities such as spacetime curvature become infinite or geodesics cannot be extended further. This signals that the theory has reached the edge of its applicability.

In Schwarzschild

At

$$r = 0,$$

the Kretschmann scalar,

$$K = R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} = \frac{48G^2M^2}{c^4r^6},$$

diverges as

$$r \rightarrow 0.$$

This divergence confirms the presence of a true **curvature singularity**, where the curvature of spacetime becomes infinite.

Inside the event horizon, the radial coordinate  $r$  behaves like a time coordinate. Just as it is impossible to stop moving forward in time, it is impossible to avoid moving toward

$$r = 0,$$

once the event horizon of an ideal Schwarzschild black hole has been crossed.

In Kerr ring singularity

Rotation changes the nature of the singularity completely.

Instead of being a single point, the singularity becomes a ring.

The ring singularity is located at

$$r = 0, \quad \theta = \frac{\pi}{2}.$$

This corresponds to the equatorial plane of the black hole. Unlike the Schwarzschild singularity, which is a point, the Kerr singularity forms a circular ring of radius

$$a = \frac{J}{Mc},$$

where

- $a$  = spin parameter,
- $J$  = angular momentum of the black hole,
- $M$  = mass of the black hole,
- $c$  = speed of light.

The strange things predicted inside a Kerr black hole come from the ideal mathematical solution. We don't know if they actually exist in real black holes

#### 4.7.5 Accretion Disk

An accretion disk is a hot, rotating disk of gas, dust, plasma, and other matter that forms around a black hole (or other compact objects like neutron stars) as material falls inward under the influence of gravity. Instead of falling directly into the black hole, the matter usually has some angular momentum, causing it to spiral inward while orbiting the black hole. An accretion disk forms because gas and dust falling toward a black hole usually have angular momentum, so they cannot fall straight in. Instead, the material spirals around the black hole, forming a rotating disk. As particles in the disk collide and rub against each other, friction and turbulence convert gravitational energy into heat, raising the temperature to millions of degrees. This intense heat causes the accretion disk to emit large amounts of light and X-rays, making it much brighter than the black hole itself.

### 4.8 Quantum Black Hole Physics

#### 4.8.1 Hawking Radiation

For many years Nothing can escape a black hole. In 1974, Stephen Hawking showed that this isn't completely true. Because of quantum mechanics, black holes slowly emit radiation. This radiation is called Hawking radiation. Quantum mechanics says that empty space is never truly empty. Even a vacuum contains tiny fluctuations called virtual particle-antiparticle pairs. These pairs appear everywhere in space. A particle and antiparticle appear. They exist for an extremely short time. They annihilate each other. No energy is left behind. Near the event horizon, gravity is extremely strong.

Sometimes the two particles are pulled apart before they can annihilate. Imagine two people holding hands. A huge current suddenly pulls one into the black hole while the other escapes. The pair separates




Figure 8: Schematic anatomy of a rotating (Kerr) black hole. The central singularity is enclosed by the event horizon. Outside it lies the photon region, where light can follow unstable orbits, and the ergosphere, where frame dragging forces all objects to co-rotate with the black hole. The surrounding accretion disk consists of hot plasma emitting electromagnetic radiation.

forever. This picture is a useful way to visualize Hawking radiation, although Hawking's original derivation is based on quantum field theory in curved spacetime rather than particles literally being created at the event horizon

Now:

One particle falls into the black hole. The other escapes into space.

The escaping particle becomes real radiation.

The black hole loses mass, because energy cannot come from nowhere the scaping particle have positive energy, the enrgy come from black hole itself. The particle falling inward effectively reduces the black hole's mass So every escaping particle makes the black hole slightly lighter. Over an enormous amount of time, the black hole slowly evaporates.




Figure 9: Conceptual illustration of Hawking radiation. (a) Quantum fluctuations produce a virtual particle–antiparticle pair near the event horizon. (b) Strong spacetime curvature separates the pair before annihilation can occur. One particle falls into the black hole while the other escapes to infinity. (c) The escaping particle is observed as Hawking radiation, and the black hole loses a corresponding amount of mass, causing it to evaporate over extremely long timescales.

—Every object with a temperature emits thermal radiation. In 1974, Stephen Hawking showed that black holes also emit thermal radiation due to quantum effects near the event horizon.

The Hawking temperature is given by

$$T_H = \frac{\hbar c^3}{8\pi G M k_B},$$

where

- $T_H$  = Hawking temperature,
- $\hbar$  = reduced Planck constant,
- $c$  = speed of light,
- $G$  = gravitational constant,
- $M$  = mass of the black hole,
- $k_B$  = Boltzmann constant.



The Hawking temperature is inversely proportional to the mass of the black hole:

$$T_{\text{H}} \propto \frac{1}{M}.$$

This means

Large mass  $\longrightarrow$  Lower temperature,

Small mass  $\longrightarrow$  Higher temperature.

A rotating black hole also emits Hawking radiation.

Unlike a Schwarzschild black hole, its temperature depends on the **surface gravity** ( $\kappa$ ) of the event horizon rather than only on its mass.

As the black hole spins faster and approaches the extremal limit,

$$a \rightarrow M,$$

its surface gravity decreases:

$$\kappa \rightarrow 0.$$

Consequently, the Hawking temperature also approaches zero:

$$T_{\text{H}} \rightarrow 0.$$

An extremal Kerr black hole therefore has zero Hawking temperature.

The Hawking temperature of a Kerr black hole is given by

$$T_{\text{H}} = \frac{\hbar \kappa}{2\pi k_{\text{B}} c},$$

where

- $T_{\text{H}}$  = Hawking temperature,
- $\hbar$  = reduced Planck constant,
- $\kappa$  = surface gravity of the event horizon,
- $k_{\text{B}}$  = Boltzmann constant,
- $c$  = speed of light in a vacuum.

## 4.9 Black Hole Thermodynamics

Before Hawking, black holes looked completely different from ordinary objects. After Hawking, physicists realized they obey laws almost identical to thermodynamics. In thermodynamics, entropy measures the amount of disorder or, equivalently, the number of microscopic states corresponding to a given macroscopic state.

Jacob Bekenstein proposed that black holes also possess entropy, and Stephen Hawking later derived its exact expression.

The Bekenstein–Hawking entropy is

$$S_{\text{BH}} = \frac{k_{\text{B}} c^3 A}{4G\hbar},$$

where

- $S_{\text{BH}}$  = Bekenstein–Hawking entropy,
- $k_{\text{B}}$  = Boltzmann constant,
- $c$  = speed of light in a vacuum,
- $A$  = surface area of the event horizon,

- $G$  = gravitational constant,
- $\hbar$  = reduced Planck constant.

The entropy of a black hole is directly proportional to the area of its event horizon:

$$S_{\text{BH}} \propto A.$$

This remarkable result shows that a black hole's entropy depends on the area of its event horizon rather than its volume, a discovery that has had profound implications for black hole thermodynamics and quantum gravity.

#### 4.9.1 Four Laws of Black Hole Thermodynamics

The laws of black hole thermodynamics closely mirror the ordinary laws of thermodynamics.

##### Zeroth Law

The surface gravity is the same everywhere on the event horizon of a stationary black hole, just as a system in thermal equilibrium has a uniform temperature.

##### First Law

If a black hole gains or loses energy, its

- mass,
- horizon area,
- spin,
- charge

change according to a precise relationship, analogous to the conservation of energy in ordinary thermodynamics.

##### Second Law

Classically,

the event horizon area of a black hole never decreases.

Black holes can merge and grow, but the total horizon area cannot shrink.

However, Hawking radiation showed that black holes can lose mass, causing their horizon area to decrease. To resolve this, Jacob Bekenstein proposed that the entropy of a black hole must be included together with the entropy of the surrounding environment.

The total quantity, known as the **generalized entropy**, never decreases.

$$S_{\text{total}} = S_{\text{BH}} + S_{\text{outside}}$$

where

- $S_{\text{BH}}$  = black hole entropy,
- $S_{\text{outside}}$  = entropy outside the black hole.

##### Third Law

It is impossible to reach an extremal black hole, which has zero surface gravity and zero Hawking temperature, through any finite physical process.

This is similar to the third law of ordinary thermodynamics, where absolute zero temperature cannot be reached in a finite number of steps. As summarized in Figure 10, each law of black hole thermodynamics has a direct analogue in classical thermodynamics, reinforcing the interpretation of black holes as genuine thermodynamic systems.




Figure 10: Correspondence between the four laws of ordinary thermodynamics and the four laws of black hole thermodynamics. The close analogy played a central role in the development of black hole thermodynamics and ultimately led to Hawking’s prediction that black holes emit thermal radiation.

#### 4.9.2 Black Hole Evaporation

Black holes slowly lose mass through Hawking radiation, which carries energy away from the event horizon.

As the black hole loses mass, its temperature increases, causing it to emit radiation more rapidly.

Over an extremely long period of time, this process may cause the black hole to completely evaporate.

#### 4.10 The Information Paradox

The Black Hole Information Paradox is a major unresolved problem in theoretical physics that arises from a conflict between General Relativity and Quantum Mechanics.

According to quantum mechanics, information can never be permanently destroyed, a principle known as **unitarity**.

However, Hawking’s original theory predicts that black holes emit thermal Hawking radiation, which appears to contain no information about the matter that fell into the black hole.

If a black hole eventually evaporates completely, the information about everything it absorbed appears to disappear, violating the principles of quantum mechanics.

Resolving this paradox remains one of the greatest challenges in modern physics, and research into this problem is still ongoing.

## 5 Theoretical Physics

Unlike black holes, which have strong observational evidence, white holes and wormholes remain theoretical predictions of General Relativity. They arise as mathematical solutions to Einstein’s field equations, although no direct observational evidence currently confirms their existence.

#### 5.0.1 The Penrose Singularity Theorem (1965)

The theorem states that:

If General Relativity is correct, enough matter is present, and gravity creates a trapped surface, then spacetime becomes geodesically incomplete.

This means that some paths through spacetime suddenly end. Physically, this is what we call a singularity.

### 5.0.2 What is a Trapped Surface?

Imagine you are standing on a sphere around a star.

Normally:

- Light going outward moves away.
- Light going inward moves inward.

Gravity is not strong enough to stop outgoing light.

During gravitational collapse, gravity becomes extremely strong. Even light trying to escape gets pulled inward instead of moving outward. Both inward and outward light rays are directed toward the centre.

That surface is called a **trapped surface**.




Figure 11: Illustration of a trapped surface. Both inward- and outward-directed light rays converge toward the centre because the gravitational field is sufficiently strong. The formation of such a trapped surface is a key assumption in Penrose's singularity theorem.

Once a trapped surface forms, Penrose proved that nothing can prevent the collapse into a singularity.

This work also changed the understanding of black holes. Before Penrose, many scientists were unsure whether black holes were physically real or only mathematical solutions.

Penrose showed that black holes are a natural prediction of General Relativity. If massive stars collapse sufficiently, black holes should inevitably form.

Decades later, astronomers observed black holes, confirming the importance of Penrose's work. This contribution earned Roger Penrose the 2020 Nobel Prize in Physics.

### 5.0.3 Penrose Diagram

People often mistake the Penrose theorem and the Penrose diagram as the same thing.

The **Penrose theorem** is a mathematical proof that predicts singularity formation using topology.

The **Penrose diagram** is a method of representing spacetime, showing causal structure using conformal geometry.

### 5.0.4 Conformal Compactification

Conformal compactification means shrinking infinite distances into a finite diagram without changing the paths of light.

Imagine Google Maps. The Earth is enormous, yet your phone can display it on a small screen. The map does not preserve actual distances, but it preserves relationships. This is similar to what Penrose did with spacetime.

It preserves angles because light is fundamental in General Relativity. The speed of light remains constant, and in a Penrose diagram light always travels at  $45^\circ$  angles.

This makes causality easier to understand.

Causality simply means:

Can one event influence another?

If light can travel from event A to event B, then A can affect B. If not, they are causally disconnected.

Instead of solving equations directly, we draw lines at  $45^\circ$ . If the light path reaches another event, communication is possible. If it cannot, the events are causally separated.

For an eternal Schwarzschild black hole, spacetime is divided into four regions.

#### 5.0.5 Region I

This is our universe: stars, Earth, planets, and everything outside the event horizon.

#### 5.0.6 Region II

This is inside the event horizon.

Once you cross into this region, every possible future direction points toward the singularity. You cannot escape, not because you are moving too slowly, but because the geometry of spacetime itself directs all future paths inward.

#### 5.0.7 Region III

This represents a second universe.

It is not another galaxy, but a completely separate region of spacetime. In the idealised eternal Schwarzschild solution, it appears mathematically, but there is no evidence that such a second universe exists in nature.

#### 5.0.8 Region IV

This is the white hole region.

A white hole is the time-reversed version of a black hole. Things can come out, but nothing can enter.

Like Region III, it appears in the maximal mathematical extension of the eternal Schwarzschild solution and is generally not expected to exist for black holes formed from collapsing stars.




Figure 12: Penrose diagram of the maximally extended Schwarzschild spacetime. Region I represents our asymptotically flat universe, Region II is the black hole interior, Region III is a second asymptotically flat universe, and Region IV corresponds to the white hole. The dashed diagonal lines denote the event horizons. Light rays always travel at  $45^\circ$ , preserving the causal structure of spacetime under conformal compactification.

## 5.1 White Holes

White holes are the theoretical opposite of black holes.

A black hole allows matter and light to enter but prevents escape.

A white hole allows matter and light to escape but prevents anything from entering.

### 5.1.1 Where Did White Holes Come From?

Scientists did not discover white holes through observation. They appeared naturally while solving Einstein's equations.

Initially, scientists thought the Schwarzschild metric only described black holes. Later, using Kruskal–Szekeres coordinates, the solution was extended to describe the complete spacetime.

This is called the **maximal extension of Schwarzschild spacetime**.

Instead of only one region, the complete solution contains four regions:

- Region I  $\rightarrow$  Our universe
- Region II  $\rightarrow$  Black hole
- Region III  $\rightarrow$  Another universe
- Region IV  $\rightarrow$  White hole

The white hole is not a separate solution of Einstein's equations. It appears naturally as Region IV in the maximally extended Schwarzschild solution.

The original Schwarzschild coordinates described only part of spacetime. Kruskal coordinates revealed the complete structure.

### 5.1.2 Region IV

Region IV is called the white hole region.

Unlike Region II, where everything moves toward the singularity, everything in Region IV moves away from it.

The singularity can be imagined as the beginning of every path rather than the end.

### 5.1.3 Time Reversal

A white hole is the time-reversed version of a black hole.

Imagine recording a star collapsing into a black hole.

Now play the video backwards.

Instead of:

star  $\rightarrow$  collapse  $\rightarrow$  black hole

you get:

white hole  $\rightarrow$  matter explodes outward  $\rightarrow$  star.

This backward movie is essentially a white hole.

This is why physicists call it the time reverse of a black hole.

A white hole still has gravity. It still attracts nearby objects. However, something strange happens. As an object approaches the horizon, spacetime prevents it from crossing. The object is forced back outward.

Therefore, the causal structure of a white hole prevents anything from entering.

A black hole event horizon is called a **trapped surface**. A white hole has an **anti-trapped surface**. Every light ray, no matter where it is directed, moves outward. Nothing can remain inside, and nothing can enter.

Although white holes are mathematically valid solutions of Einstein's field equations, they are believed to be classically unstable.

In 1974, Douglas Eardley showed that even a tiny amount of infalling matter or radiation would become highly blueshifted near the white hole horizon. This causes energy to accumulate, destabilizing the horizon. As a result, the white hole horizon would rapidly collapse into a black hole.




Figure 13: Conceptual comparison between a black hole and its time-reversed counterpart, the white hole. A black hole permits matter and radiation to cross the event horizon inward but not outward. A white hole exhibits the opposite causal behaviour: matter and radiation emerge from the horizon, while nothing can enter.

Furthermore, unlike black holes, which can form naturally through the gravitational collapse of massive stars, there is no known physical mechanism that can create a white hole.

For matter to emerge from a singularity would require extremely unusual initial conditions and would contradict the observed arrow of time. Therefore, long-lived white holes are considered extremely unlikely to exist in the real universe.

#### 5.1.4 Entropy and the Arrow of Time

Another reason white holes are considered unlikely is their conflict with the arrow of time.

According to the second law of thermodynamics, the universe naturally evolves toward higher entropy, meaning that systems become more disordered over time.

The formation of a black hole is an entropy-increasing process, whereas a white hole would represent the reverse process, requiring matter to emerge from a singularity and entropy to decrease.

Such a process is considered extremely improbable, making the spontaneous formation of white holes inconsistent with the observed direction of time in our universe.

#### 5.1.5 Connection to Wormholes

White holes are closely related to wormholes through the maximally extended Schwarzschild solution.

In this geometry, the Einstein–Rosen bridge mathematically connects two separate exterior regions of spacetime, with the white hole (Region IV) and black hole (Region II) forming part of the complete structure.

However, this wormhole is non-traversable because it collapses too quickly for matter or information to pass through from one universe to the other.

#### 5.1.6 Quantum Gravity and White Holes

Although classical physics predicts that white holes are unstable, some theories of quantum gravity suggest that they may still exist.

One notable proposal is the **Planck Star model**, developed by Carlo Rovelli and collaborators.

According to this idea, quantum gravitational effects become dominant near the singularity, halting gravitational collapse and causing the black hole to eventually transition into a white hole.

This process could release the trapped matter and information, offering a possible solution to the black hole information paradox.

However, this remains a speculative hypothesis and has not yet been confirmed by observations or experiments.

#### 5.1.7 Einstein–Rosen Bridge

The concept of the **Einstein–Rosen bridge** was introduced in 1935 by Albert Einstein and Nathan Rosen in their paper *The Particle Problem in the General Theory of Relativity*.

At the time, they were investigating mathematical solutions to Einstein's field equations and searching for a way to describe elementary particles without the singularities that appeared in General Relativity.

While analyzing the Schwarzschild solution, they found that its maximally extended geometry could be interpreted as connecting two separate asymptotically flat regions of spacetime through a narrow bridge. This bridge appeared as a tunnel-like structure linking two distant regions, rather than ending abruptly at a singularity. Einstein and Rosen proposed this geometric connection, later named the **Einstein–Rosen bridge**, as a possible model for particles and as a way to avoid the problematic point singularities predicted by the theory.

Initially, the bridge suggested the fascinating possibility of a shortcut through spacetime. However, later work by physicists, particularly through the development of **Kruskal–Szekeres coordinates**, showed that the Einstein–Rosen bridge in the Schwarzschild solution is **non-traversable**. The bridge exists only momentarily and collapses too quickly for light or matter to cross from one side to the other. As a result, it cannot be used for interstellar travel or communication.

Although the original Einstein–Rosen bridge is not physically traversable, its discovery marked the birth of wormhole physics. It demonstrated that General Relativity naturally permits geometries that connect distant regions of spacetime, inspiring decades of research into traversable wormholes, exotic matter, quantum gravity, and modern ideas such as the

$$ER = EPR$$

conjecture, which explores a possible connection between wormholes and quantum entanglement.

### 5.1.8 What is the Einstein–Rosen Bridge?

The **Einstein–Rosen bridge** is the original theoretical concept of a wormhole proposed within General Relativity. It appears in the maximally extended Schwarzschild solution, where two separate asymptotically flat regions of spacetime are mathematically connected by a bridge-like geometry. Although this structure resembles a tunnel linking two distant regions, it is not a physical passage through which humans or objects could travel. Instead, it is an idealized mathematical feature of the Schwarzschild spacetime.

### 5.1.9 Non-Traversability

The **Einstein–Rosen bridge** is non-traversable because its throat collapses too quickly for light or matter to pass through. The bridge exists only momentarily and pinches off before anything entering one side can emerge from the other. As a result, it cannot be used for travel or communication.




Figure 14: Schematic illustration of an Einstein–Rosen bridge connecting two asymptotically flat regions of spacetime. In the Schwarzschild solution, the bridge is non-traversable because its throat collapses before light or matter can cross it.

## 5.2 Worm Holes

A wormhole is a hypothetical tunnel-like structure in spacetime that acts as a shortcut connecting two separate regions of the universe or even different universes. It consists of two mouths joined by a central



passage called the throat. Unlike a black hole, which traps anything that crosses its event horizon, a traversable wormhole is theoretically designed to allow matter and light to pass through from one mouth to the other. Although wormholes are valid mathematical solutions to Einstein's General Relativity, no observational evidence has yet confirmed their existence.




Figure 15: Conceptual illustration of a traversable wormhole. Two mouths are connected by a throat through which matter and light could theoretically travel if the throat remained stable.

### 5.2.1 Morris–Thorne Traversable Wormholes (1988)

In 1988, Michael S. Morris and Kip S. Thorne proposed a theoretical model for a traversable wormhole—a wormhole that could, in principle, allow matter and light to travel safely from one region of spacetime to another. Unlike the Einstein–Rosen bridge, which is non-traversable, the Morris–Thorne wormhole was specifically designed to remain open and stable.

The geometry of a Morris–Thorne wormhole is described by the metric

$$ds^2 = -e^{2\Phi(r)} dt^2 + \frac{dr^2}{1 - \frac{b(r)}{r}} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2).$$

Although the equation appears complex, each term has a clear physical meaning:

- **Redshift Function,  $\Phi(r)$ :** Determines how gravity affects the passage of time. To keep the wormhole traversable, this function must remain finite everywhere so that no event horizon forms and travellers can pass through without becoming trapped.
- **Shape Function,  $b(r)$ :** Describes the spatial geometry of the wormhole and determines how the tunnel curves through spacetime. At the wormhole's narrowest point, called the throat, the condition

$$b(r) = r$$

is satisfied.

- **Wormhole Throat:** The throat is the narrowest part of the tunnel connecting the two mouths of the wormhole. It is the critical region through which matter or light must pass when travelling between the two connected regions of spacetime.

### 5.2.2 Morris–Thorne Traversable Wormholes (1988)

In 1988, Michael S. Morris and Kip S. Thorne proposed a theoretical model for a traversable wormhole—a wormhole that could, in principle, allow matter and light to travel safely from one region of spacetime to another. Unlike the Einstein–Rosen bridge, which is non-traversable, the Morris–Thorne wormhole was specifically designed to remain open and stable.

The geometry of a Morris–Thorne wormhole is described by the metric

$$ds^2 = -e^{2\Phi(r)} dt^2 + \frac{dr^2}{1 - \frac{b(r)}{r}} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2).$$

Although the equation appears complex, each term has a clear physical meaning:

- **Redshift Function,  $\Phi(r)$ :** Determines how gravity affects the passage of time. To keep the wormhole traversable, this function must remain finite everywhere so that no event horizon forms and travellers can pass through without becoming trapped.

- **Shape Function,  $b(r)$ :** Describes the spatial geometry of the wormhole and determines how the tunnel curves through spacetime. At the wormhole's narrowest point, called the throat, the condition

$$b(r) = r$$

is satisfied.

- **Wormhole Throat:** The throat is the narrowest part of the tunnel connecting the two mouths of the wormhole. It is the critical region through which matter or light must pass when travelling between the two connected regions of spacetime.

The Morris–Thorne model showed that a traversable wormhole is mathematically consistent within General Relativity. However, maintaining the throat requires **exotic matter** with negative energy density to prevent the wormhole from collapsing under its own gravity. Since no confirmed source of macroscopic exotic matter has been discovered, traversable wormholes remain theoretical and have not been observed in nature.

### 5.2.3 Wormhole Stability

A key challenge in wormhole physics is stability. Even if a traversable wormhole could exist, it must remain open long enough for matter or light to pass through. According to General Relativity, the intense gravitational attraction would naturally cause the wormhole throat to collapse. To prevent this collapse, the wormhole must be supported by exotic matter, a hypothetical form of matter with negative energy density that counteracts gravity. The stability of traversable wormholes remains an active area of research, and no observational evidence has yet confirmed that such stable wormholes exist in nature.




Figure 16: Embedding diagram illustrating the spatial geometry of a Morris–Thorne wormhole. The narrow central region represents the throat connecting two asymptotically flat regions of spacetime.

### 5.2.4 Exotic Matter and Energy Conditions

One of the greatest challenges in wormhole physics is the requirement for **exotic matter**, a hypothetical form of matter with negative energy density. In General Relativity, the relationship between matter, energy, and the curvature of spacetime is described by the stress–energy tensor,  $T_{\mu\nu}$ , which acts as the source of gravity in Einstein’s field equations. While ordinary matter produces positive energy density and attractive gravity, exotic matter is predicted to possess negative energy density, creating a repulsive gravitational effect that can prevent the wormhole throat from collapsing.

A traversable wormhole requires a violation of the **Null Energy Condition (NEC)**, which is expressed as

$$T_{\mu\nu}k^{\mu}k^{\nu} \geq 0,$$

where  $T_{\mu\nu}$  is the stress–energy tensor and  $k^{\mu}$  is any null (light-like) vector. Ordinary matter satisfies this condition, meaning its energy density measured along light rays is always non-negative. However, to keep a wormhole throat open against gravitational collapse, the energy density near the throat must become negative, violating the NEC. This violation allows spacetime to remain open rather than pinching shut.

Although no macroscopic form of exotic matter has been observed, quantum physics provides a small-scale example through the **Casimir effect**. When two conducting plates are placed extremely close together in a vacuum, quantum fluctuations create a region with negative energy density between the plates. While the Casimir effect demonstrates that negative energy is physically possible, the amount produced is extraordinarily small and far below what would be required to stabilize a traversable wormhole. Consequently, exotic matter remains one of the greatest theoretical obstacles to the existence of traversable wormholes.

The Morris–Thorne model showed that a traversable wormhole is mathematically consistent within General Relativity. However, maintaining the throat requires **exotic matter** with negative energy density to prevent the wormhole from collapsing under its own gravity. Since no confirmed source of macroscopic exotic matter has been discovered, traversable wormholes remain theoretical and have not been observed in nature.

### 5.2.5 Wormhole Stability

A key challenge in wormhole physics is stability. Even if a traversable wormhole could exist, it must remain open long enough for matter or light to pass through. According to General Relativity, the intense gravitational attraction would naturally cause the wormhole throat to collapse. To prevent this collapse, the wormhole must be supported by exotic matter, a hypothetical form of matter with negative energy density that counteracts gravity. The stability of traversable wormholes remains an active area of research, and no observational evidence has yet confirmed that such stable wormholes exist in nature.

### 5.2.6 Exotic Matter and Energy Conditions

One of the greatest challenges in wormhole physics is the requirement for **exotic matter**, a hypothetical form of matter with negative energy density. In General Relativity, the relationship between matter, energy, and the curvature of spacetime is described by the stress–energy tensor,  $T_{\mu\nu}$ , which acts as the source of gravity in Einstein’s field equations. While ordinary matter produces positive energy density and attractive gravity, exotic matter is predicted to possess negative energy density, creating a repulsive gravitational effect that can prevent the wormhole throat from collapsing.

A traversable wormhole requires a violation of the **Null Energy Condition (NEC)**, which is expressed as

$$T_{\mu\nu}k^{\mu}k^{\nu} \geq 0,$$

where  $T_{\mu\nu}$  is the stress–energy tensor and  $k^{\mu}$  is any null (light-like) vector. Ordinary matter satisfies this condition, meaning its energy density measured along light rays is always non-negative. However, to keep a wormhole throat open against gravitational collapse, the energy density near the throat must become negative, violating the NEC. This violation allows spacetime to remain open rather than pinching shut.

Although no macroscopic form of exotic matter has been observed, quantum physics provides a small-scale example through the **Casimir effect**. When two conducting plates are placed extremely close together in a vacuum, quantum fluctuations create a region with negative energy density between the plates. While the Casimir effect demonstrates that negative energy is physically possible, the amount produced is extraordinarily small and far below what would be required to stabilize a traversable wormhole. Consequently, exotic matter remains one of the greatest theoretical obstacles to the existence of traversable wormholes.





Figure 17: Schematic representation of exotic matter concentrated near the wormhole throat. Negative energy density is required to counteract gravitational collapse and keep the throat open in the Morris–Thorne model.

### 5.3 Connection Between Black Holes, White Holes, and Wormholes

Black holes, white holes, and the **Einstein–Rosen bridge** are all part of the same mathematical solution of Einstein’s field equations—the **maximally extended Schwarzschild solution**. Rather than being separate objects, they represent different regions of the same spacetime geometry, each with a distinct physical interpretation. The black hole is a region from which nothing can escape, the white hole is its time-reversed counterpart from which nothing can enter, and the Einstein–Rosen bridge mathematically connects two asymptotically flat regions of spacetime. Although this bridge is non-traversable, it inspired the concept of traversable wormholes developed later by Morris and Thorne.



While these structures belong to the same mathematical family, only black holes have strong observational evidence. White holes and wormholes remain theoretical predictions of General Relativity, with no direct observational confirmation to date.

## 6 Future Research Questions

Throughout this project, several conceptual questions arose while studying black holes, spacetime, and general relativity. Some of these questions have established answers within modern physics, while others remain active areas of theoretical research.

1. If gravity depends on mass, why does a black hole not pull the entire Universe into itself?
2. Why do objects orbit a black hole instead of immediately falling into it?
3. Is a black hole truly like a cosmic vacuum cleaner, or is this a common misconception?
4. If black holes can possess electric charge, why do they not produce strong electrical effects on their surroundings?
5. What is the true nature of the singularity? Does it physically exist, or does it indicate the breakdown of General Relativity?
6. Could certain black holes contain or connect to traversable wormholes?
7. If a wormhole exists, where could its other end lead?
8. Could one mouth of a wormhole emerge as a white hole?
9. What type of matter or energy would be required to stabilize a traversable wormhole?
10. Could a stable wormhole function as a time machine?
11. Why does gravity slow the passage of time?
12. Why does relativity appear to permit travel into the future but not into the past?
13. Does nature contain a mechanism that prevents backward time travel?
14. If light has no rest mass, why is its path bent by gravity?
15. If spacetime is curved, what is it actually curved into, if anything?
16. Can spacetime tear, break, or possess defects?

17. If spacetime is curved, why are planets and satellites able to maintain stable orbits?
18. How does Earth's rotation continue while moving through curved spacetime?
19. How do rotating black holes drag spacetime around them (frame dragging)?
20. Could microscopic black holes exist within the quantum world?
21. Could quantum-scale wormholes exist?
22. Why are General Relativity and Quantum Mechanics currently incompatible?
23. What is quantum foam, and does it influence spacetime?
24. Can a neutron star fall into a black hole?
25. What happens to neutron-star matter after crossing the event horizon?
26. If a neutron star is tidally disrupted before entering a black hole, why do its fragments not collapse into smaller black holes?
27. Could smaller black holes exist inside a larger black hole?
28. If space is expanding, what is it expanding into?
29. Is spacetime fundamental, or does it emerge from a deeper physical structure?
30. Could extra spatial dimensions exist beyond those we directly observe?
31. Could black holes provide the key to unifying gravity with quantum mechanics?
32. Is there any theoretical relationship between black hole singularities, Einstein–Rosen bridges, and traversable wormholes?

The following questions were developed during the course of this project. They represent areas of curiosity inspired by the study of black hole physics. Some have established answers within current physical theories, while others remain open problems in modern theoretical physics.