

Influence functions and functional derivatives

Anders Munch

May 11, 2021

Outline

Setting

Motivation

Functional derivatives

Canonical gradient / efficient influence function

Summary of main results

Next step – constructing estimators

Disclaimer about the note

- ▶ The note is work in progress, and we have not used it before – you are very welcome to comment on weird/unclear passages.

Disclaimer about the note

- ▶ The note is work in progress, and we have not used it before – you are very welcome to comment on weird/unclear passages.
- ▶ You should see it as a service – some exact mathematical statements are collected there if you care about it, but the important part is the intuition which we talk about today.

Disclaimer about the note

- ▶ The note is work in progress, and we have not used it before – you are very welcome to comment on weird/unclear passages.
- ▶ You should see it as a service – some exact mathematical statements are collected there if you care about it, but the important part is the intuition which we talk about today.
- ▶ Do NOT write like this in your report!

A statistical problem

We call a collection of probability measures \mathcal{P} together with a functional $\Psi: \mathcal{P} \rightarrow \mathbb{R}$ a *statistical problem*.

A statistical problem

We call a collection of probability measures \mathcal{P} together with a functional $\Psi: \mathcal{P} \rightarrow \mathbb{R}$ a *statistical problem*.

Example (Average treatment effect)

We are given n iid. sample of $O \sim P$, with $P \in \mathcal{P}$ and where $O = (X, A, Y)$, with $X \in \mathbb{R}^d$, $A \in \{0, 1\}$, and $Y \in \{0, 1\}$. We want to estimate the average treatment effect

$$\mathbb{E}_P [f(1, X) - f(0, X)],$$

with $f(a, x) := \mathbb{E}_P [Y \mid A = a, X = x]$. The target parameter is

$$\Psi(P) = \mathbb{E}_P [f_P(1, X) - f_P(0, X)].$$

A statistical problem

We call a collection of probability measures \mathcal{P} together with a functional $\Psi: \mathcal{P} \rightarrow \mathbb{R}$ a *statistical problem*.

Example (Average treatment effect)

We are given n iid. sample of $O \sim P$, with $P \in \mathcal{P}$ and where $O = (X, A, Y)$, with $X \in \mathbb{R}^d$, $A \in \{0, 1\}$, and $Y \in \{0, 1\}$. We want to estimate the average treatment effect

$$\mathbb{E}_P [f(1, X) - f(0, X)],$$

with $f(a, x) := \mathbb{E}_P [Y \mid A = a, X = x]$. The target parameter is

$$\Psi(P) = \mathbb{E}_P [f_P(1, X) - f_P(0, X)].$$

Target and nuisance parameters

Target and nuisance parameters

Target parameter Low-dimensional, scientifically meaningful.

Target and nuisance parameters

Target parameter Low-dimensional, scientifically meaningful.

Nuisance parameters Needed to express the target parameter.

Target and nuisance parameters

Target parameter Low-dimensional, scientifically meaningful.

Nuisance parameters Needed to express the target parameter.

Example (ATE)

The ATE can be written as $\Psi(P) = P[\varphi_1] = P[\varphi_2] = P[\varphi_3]$, for

$$\varphi_1(o; f) := f(1, x) - f(0, x),$$

$$\varphi_2(o; \pi) := \frac{a y}{\pi(x)} - \frac{(1 - a) y}{1 - \pi(x)},$$

$$\varphi_3(o; f, \pi) := \varphi_1(o; f) + \varphi_2(o; \pi) - \frac{a f(1, x)}{\pi(x)} + \frac{(1 - a) f(0, x)}{1 - \pi(x)},$$

with $f(a, x) := \mathbb{E}_P [Y \mid A = a, X = x]$, $\pi(x) := P(A = 1 \mid X = x)$.

$P[\varphi]$ means

$$P[\varphi] = \mathbb{E}_P [\varphi(O)] = \int \varphi(o) dP(o).$$

Infinite-dimensional nuisance parameters

A parametric setting means that \mathcal{P} is finite-dimensional. We are interested in *nonparametric* or *semiparametric* settings which mean that \mathcal{P} is infinite-dimensional.

Infinite-dimensional nuisance parameters

A parametric setting means that \mathcal{P} is finite-dimensional. We are interested in *nonparametric* or *semiparametric* settings which mean that \mathcal{P} is infinite-dimensional.

Having our data set and scientific question in mind, why would it be of interest to use infinite-dimensional nuisance parameters?

Infinite-dimensional nuisance parameters

A parametric setting means that \mathcal{P} is finite-dimensional. We are interested in *nonparametric* or *semiparametric* settings which mean that \mathcal{P} is infinite-dimensional.

Having our data set and scientific question in mind, why would it be of interest to use infinite-dimensional nuisance parameters?

Trying to control for confounding \implies nice to have:

- ▶ flexible model
- ▶ many covariates

Toy example: Integrated kernel density

\mathcal{P} consist all probability measures with continuous Lebesgue-density (this is an infinite-dimensional space). We want to estimate $F(x) = P(X \leq x)$ for unknown $P \in \mathcal{P}$.

Toy example: Integrated kernel density

\mathcal{P} consist all probability measures with continuous Lebesgue-density (this is an infinite-dimensional space). We want to estimate $F(x) = P(X \leq x)$ for unknown $P \in \mathcal{P}$. Our target parameter is then $\theta = \Psi(P) = F_P(x)$ which we can express as

$$\Psi(P) = \Psi_0(f) := \int_{-\infty}^x f(z) \, dz, \quad \text{for } P = f \cdot \lambda,$$

because of our assumption about \mathcal{P} .

Toy example: Integrated kernel density

\mathcal{P} consist all probability measures with continuous Lebesgue-density (this is an infinite-dimensional space). We want to estimate $F(x) = P(X \leq x)$ for unknown $P \in \mathcal{P}$. Our target parameter is then $\theta = \Psi(P) = F_P(x)$ which we can express as

$$\Psi(P) = \Psi_0(f) := \int_{-\infty}^x f(z) \, dz, \quad \text{for } P = f \cdot \lambda,$$

because of our assumption about \mathcal{P} . We want to use **machine learning** (!) for this problem, so use a kernel estimator, i.e.,

$$\hat{f}_n(x) = \hat{\mathbb{P}}_n[k_h(X, x)] = \frac{1}{n} \sum_{i=1}^n k_h(X_i, x),$$

where k_h is, e.g. $k_h(x, y) = \frac{1}{h} k\left(\frac{x-y}{h}\right)$, with k the density for the standard Gaussian distribution, and the bandwidth h is chosen using cross-validation.

Toy example: Integrated kernel density

\mathcal{P} consist all probability measures with continuous Lebesgue-density (this is an infinite-dimensional space). We want to estimate $F(x) = P(X \leq x)$ for unknown $P \in \mathcal{P}$. Our target parameter is then $\theta = \Psi(P) = F_P(x)$ which we can express as

$$\Psi(P) = \Psi_0(f) := \int_{-\infty}^x f(z) \, dz, \quad \text{for } P = f \cdot \lambda,$$


because of our assumption about \mathcal{P} . We want to use **machine learning** (!) for this problem, so use a kernel estimator, i.e.,

$$\hat{f}_n(x) = \hat{\mathbb{P}}_n[k_h(X, x)] = \frac{1}{n} \sum_{i=1}^n k_h(X_i, x),$$

where k_h is, e.g. $k_h(x, y) = \frac{1}{h} k\left(\frac{x-y}{h}\right)$, with k the density for the standard Gaussian distribution, and the bandwidth h is chosen using cross-validation. We then obtain the target estimator $\hat{\theta}_n = \Psi_0(\hat{f}_n)$.

How does this work in practice?

How does this work in practice?



What happened?

What happened?

Consider a general problem (\mathcal{P}, Ψ) for which we can write $\Psi(P) = \Psi_0(P, \nu) = P[\varphi(O, \nu)]$.

What happened?

Consider a general problem (\mathcal{P}, Ψ) for which we can write $\Psi(P) = \Psi_0(P, \nu) = P[\varphi(O, \nu)]$. We have

$$\begin{aligned}\sqrt{n}(\hat{\theta}_n - \theta) &= \sqrt{n}(\Psi_0(\hat{\mathbb{P}}_n, \hat{\nu}_n) - \Psi_0(P, \nu)) \\ &= \sqrt{n}(\hat{\mathbb{P}}_n[\varphi(O, \hat{\nu}_n)] - P[\varphi(O, \nu)]) \\ &= \sqrt{n}(\hat{\mathbb{P}}_n[\varphi(O, \hat{\nu}_n)] \pm P[\varphi(O, \hat{\nu}_n)] - P[\varphi(O, \nu)]) \\ &= \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] + \sqrt{n}\{\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu)\},\end{aligned}$$

with $\mathbb{G}_n := \sqrt{n}(\hat{\mathbb{P}}_n - P)$ the empirical process.

What happened?

Consider a general problem (\mathcal{P}, Ψ) for which we can write $\Psi(P) = \Psi_0(P, \nu) = P[\varphi(O, \nu)]$. We have

$$\begin{aligned}\sqrt{n}(\hat{\theta}_n - \theta) &= \sqrt{n}(\Psi_0(\hat{\mathbb{P}}_n, \hat{\nu}_n) - \Psi_0(P, \nu)) \\&= \sqrt{n}(\hat{\mathbb{P}}_n[\varphi(O, \hat{\nu}_n)] - P[\varphi(O, \nu)]) \\&= \sqrt{n}(\hat{\mathbb{P}}_n[\varphi(O, \hat{\nu}_n)] \pm P[\varphi(O, \hat{\nu}_n)] - P[\varphi(O, \nu)]) \\&= \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] + \sqrt{n}\{\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu)\},\end{aligned}$$

with $\mathbb{G}_n := \sqrt{n}(\hat{\mathbb{P}}_n - P)$ the empirical process.

$\mathbb{G}_n[\varphi(O, \hat{\nu}_n)]$ determines the (main) variance

What happened?

Consider a general problem (\mathcal{P}, Ψ) for which we can write $\Psi(P) = \Psi_0(P, \nu) = P[\varphi(O, \nu)]$. We have

$$\begin{aligned}\sqrt{n}(\hat{\theta}_n - \theta) &= \sqrt{n}(\Psi_0(\hat{\mathbb{P}}_n, \hat{\nu}_n) - \Psi_0(P, \nu)) \\ &= \sqrt{n}(\hat{\mathbb{P}}_n[\varphi(O, \hat{\nu}_n)] - P[\varphi(O, \nu)]) \\ &= \sqrt{n}(\hat{\mathbb{P}}_n[\varphi(O, \hat{\nu}_n)] \pm P[\varphi(O, \hat{\nu}_n)] - P[\varphi(O, \nu)]) \\ &= \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] + \sqrt{n}\{\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu)\},\end{aligned}$$

with $\mathbb{G}_n := \sqrt{n}(\hat{\mathbb{P}}_n - P)$ the empirical process.

$\mathbb{G}_n[\varphi(O, \hat{\nu}_n)]$ determines the (main) variance

$\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu)$ is bias!

What to do? – Taylor expansion

What to do? – Taylor expansion

Assume we could make a Taylor expansion of $\nu \mapsto \Psi_0(P, \nu)$, so that

$$\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu) = D_\nu \Psi_0[\hat{\nu}_n - \nu] + \mathcal{O}_P(\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2).$$

What to do? – Taylor expansion

Assume we could make a Taylor expansion of $\nu \mapsto \Psi_0(P, \nu)$, so that

$$\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu) = D_\nu \Psi_0[\hat{\nu}_n - \nu] + \mathcal{O}_P(\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2).$$

The decomposition then becomes

$$\sqrt{n}(\hat{\theta}_n - \theta) = \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] \tag{1}$$

$$+ D_\nu \Psi_0[\sqrt{n}(\hat{\nu}_n - \nu)] \tag{2}$$

$$+ \mathcal{O}_P(\sqrt{n}\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2). \tag{3}$$

What to do? – Taylor expansion

Assume we could make a Taylor expansion of $\nu \mapsto \Psi_0(P, \nu)$, so that

$$\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu) = D_\nu \Psi_0[\hat{\nu}_n - \nu] + \mathcal{O}_P(\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2).$$

The decomposition then becomes

$$\sqrt{n}(\hat{\theta}_n - \theta) = \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] \tag{1}$$

$$+ D_\nu \Psi_0[\sqrt{n}(\hat{\nu}_n - \nu)] \tag{2}$$

$$+ \mathcal{O}_P(\sqrt{n}\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2). \tag{3}$$

(1) can be handled by empirical process theory or sample splitting

What to do? – Taylor expansion

Assume we could make a Taylor expansion of $\nu \mapsto \Psi_0(P, \nu)$, so that

$$\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu) = D_\nu \Psi_0[\hat{\nu}_n - \nu] + \mathcal{O}_P(\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2).$$

The decomposition then becomes

$$\sqrt{n}(\hat{\theta}_n - \theta) = \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] \tag{1}$$

$$+ D_\nu \Psi_0[\sqrt{n}(\hat{\nu}_n - \nu)] \tag{2}$$

$$+ \mathcal{O}_P(\sqrt{n}\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2). \tag{3}$$

(1) can be handled by empirical process theory or sample splitting

(2) is our focus! \rightarrow make sense of this

What to do? – Taylor expansion

Assume we could make a Taylor expansion of $\nu \mapsto \Psi_0(P, \nu)$, so that

$$\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu) = D_\nu \Psi_0[\hat{\nu}_n - \nu] + \mathcal{O}_P(\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2).$$

The decomposition then becomes

$$\sqrt{n}(\hat{\theta}_n - \theta) = \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] \tag{1}$$

$$+ D_\nu \Psi_0[\sqrt{n}(\hat{\nu}_n - \nu)] \tag{2}$$

$$+ \mathcal{O}_P(\sqrt{n}\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2). \tag{3}$$

- (1) can be handled by empirical process theory or sample splitting
- (2) is our focus! \rightarrow make sense of this
- (3) is specific to the functional Ψ , but importantly the rate $\sqrt{n}\|\hat{\nu}_n - \nu\|_{\mathcal{V}} = \mathcal{O}_P(n^{-1/4})$ is sufficient; whether this holds then depends on the specific nuisance estimator.

What to do? – Taylor expansion

Assume we could make a Taylor expansion of $\nu \mapsto \Psi_0(P, \nu)$, so that

$$\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu) = D_\nu \Psi_0[\hat{\nu}_n - \nu] + \mathcal{O}_P(\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2).$$

The decomposition then becomes

$$\sqrt{n}(\hat{\theta}_n - \theta) = \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] \tag{1}$$

$$+ D_\nu \Psi_0[\sqrt{n}(\hat{\nu}_n - \nu)] \tag{2}$$

$$+ \mathcal{O}_P(\sqrt{n}\|\hat{\nu}_n - \nu\|_{\mathcal{V}}^2). \tag{3}$$

- (1) can be handled by empirical process theory or sample splitting
- (2) **is our focus!** \rightarrow **make sense of this**
- (3) is specific to the functional Ψ , but importantly the rate $\sqrt{n}\|\hat{\nu}_n - \nu\|_{\mathcal{V}} = \mathcal{O}_P(n^{-1/4})$ is sufficient; whether this holds then depends on the specific nuisance estimator.

Defining a functional derivative

What is a derivative?

Defining a functional derivative

What is a derivative?

A linear approximation $\dot{\Psi}_x$ to the map Ψ at $x \in \mathcal{M}$, i.e.,

$$\left\| \Psi(x + \varepsilon_n h_n) - \Psi(x) - \dot{\Psi}_x(\varepsilon_n h_n) \right\| = \mathcal{O}(\varepsilon_n),$$

when $\varepsilon_n \rightarrow 0$.

Defining a functional derivative

What is a derivative?

A linear approximation $\dot{\Psi}_x$ to the map Ψ at $x \in \mathcal{M}$, i.e.,

$$\left\| \Psi(x + \varepsilon_n h_n) - \Psi(x) - \dot{\Psi}_x(\varepsilon_n h_n) \right\| = \mathcal{O}(\varepsilon_n),$$

when $\varepsilon_n \rightarrow 0$.

This expression also makes sense for functionals (or operators) Ψ .

Defining a functional derivative

What is a derivative?

A linear approximation $\dot{\Psi}_x$ to the map Ψ at $x \in \mathcal{M}$, i.e.,

$$\left\| \Psi(x + \varepsilon_n h_n) - \Psi(x) - \dot{\Psi}_x(\varepsilon_n h_n) \right\| = \mathcal{O}(\varepsilon_n),$$

when $\varepsilon_n \rightarrow 0$.

This expression also makes sense for functionals (or operators) Ψ .

- For which h_n should this hold? Along “lines”, “paths”, or “uniformly” (h_n fixed, converging, or bounded)?

Defining a functional derivative

What is a derivative?

A linear approximation $\dot{\Psi}_x$ to the map Ψ at $x \in \mathcal{M}$, i.e.,

$$\left\| \Psi(x + \varepsilon_n h_n) - \Psi(x) - \dot{\Psi}_x(\varepsilon_n h_n) \right\| = \mathcal{O}(\varepsilon_n),$$

when $\varepsilon_n \rightarrow 0$.

This expression also makes sense for functionals (or operators) Ψ .

- ▶ For which h_n should this hold? Along “lines”, “paths”, or “uniformly” (h_n fixed, converging, or bounded)?
- ▶ Which norm on \mathcal{M} should we use?

Defining a functional derivative

What is a derivative?

A linear approximation $\dot{\Psi}_x$ to the map Ψ at $x \in \mathcal{M}$, i.e.,

$$\left\| \Psi(x + \varepsilon_n h_n) - \Psi(x) - \dot{\Psi}_x(\varepsilon_n h_n) \right\| = \mathcal{O}(\varepsilon_n),$$


when $\varepsilon_n \rightarrow 0$.

This expression also makes sense for functionals (or operators) Ψ .

- ▶ For which h_n should this hold? Along “lines”, “paths”, or “uniformly” (h_n fixed, converging, or bounded)?
- ▶ Which norm on \mathcal{M} should we use?
- ▶ In which space should we represent \mathcal{P} ?


Pathwise Hadamard differentiability

Think of the gradient of a function defined on a manifold (surface).




Pathwise Hadamard differentiability

Think of the gradient of a function defined on a manifold (surface).



Pathwise Hadamard differentiability

Think of the gradient of a function defined on a manifold (surface).



Canonical gradient

Definition (Canonical gradient)

Let (\mathcal{P}, Ψ) be a statistical problem, with $\mathcal{P} \subset \mathcal{M}_\mu$, and $\dot{\mathcal{P}}_P$ the tangent space of \mathcal{P} at $P \in \mathcal{P}$. If $\Psi: \mathcal{P} \rightarrow \mathbb{R}$ is Hadamard differentiable at P tangential to $\dot{\mathcal{P}}_P$, we refer to the Hadamard derivative $\dot{\Psi}_P$ as the *canonical gradient of the statistical problem*.

Canonical gradient

Definition (Canonical gradient)

Let (\mathcal{P}, Ψ) be a statistical problem, with $\mathcal{P} \subset \mathcal{M}_\mu$, and $\dot{\mathcal{P}}_P$ the tangent space of \mathcal{P} at $P \in \mathcal{P}$. If $\Psi: \mathcal{P} \rightarrow \mathbb{R}$ is Hadamard differentiable at P tangential to $\dot{\mathcal{P}}_P$, we refer to the Hadamard derivative $\dot{\Psi}_P$ as the *canonical gradient of the statistical problem*.

Characterizing property

With $\Gamma_P := \overline{\text{span}}\{\dot{\ell}_0\} \subset \mathcal{L}_P^2$, where $\dot{\ell}_0 = \partial_0 \log(p_\varepsilon)$ is the score function of the sub-model P_ε , there exists a unique element $\varphi_P \in \Gamma_P$ such that

$$\partial_0 \Psi(P_\varepsilon) = \langle \varphi_P, \dot{\ell}_0 \rangle_P$$

holds for any differentiable submodel P_ε with score function $\dot{\ell}_0$.

Canonical gradient for the ATE

Example (ATE)

When we make no assumptions about \mathcal{P} , the canonical gradient for the ATE problem

$$\begin{aligned}\varphi_{\mathcal{P}}(o; f, \pi) &:= f(1, x) - f(0, x) \\ &\quad + \frac{a y}{\pi(x)} - \frac{(1-a) y}{1 - \pi(x)} \\ &\quad - \frac{a f(1, x)}{\pi(x)} + \frac{(1-a) f(0, x)}{1 - \pi(x)} \\ &\quad - \Psi(\mathcal{P})\end{aligned}$$

Canonical gradient for the ATE

Example (ATE)

When we make no assumptions about \mathcal{P} , the canonical gradient for the ATE problem

$$\begin{aligned}\varphi_{\mathcal{P}}(o; f, \pi) &:= f(1, x) - f(0, x) \\ &\quad + \frac{a y}{\pi(x)} - \frac{(1-a) y}{1 - \pi(x)} \\ &\quad - \frac{a f(1, x)}{\pi(x)} + \frac{(1-a) f(0, x)}{1 - \pi(x)} \\ &\quad - \Psi(\mathcal{P})\end{aligned}$$

One way to show this is to first show that the tangent space $\Gamma_{\mathcal{P}}$ is the full subset $\mathbb{H}_0 \subset \mathcal{L}_{\mathcal{P}}^2$ of zero-mean functions, and then show that $\partial_0 \Psi(\mathcal{P}_{\varepsilon}) = \langle \varphi_{\mathcal{P}}, \dot{\ell}_0 \rangle_{\mathcal{P}}$ for all $\mathcal{P}_{\varepsilon}$ (see for instance Kennedy [2016]).

Neyman orthogonality

Theorem (Neyman orthogonality)

If $\Psi(P) = \Psi_0(P, \nu) = P[\varphi(O, \nu(P))]$ and $\varphi(\cdot, \nu) - P[\varphi(O, \nu)]$ is the canonical gradient of (\mathcal{P}, Ψ) then $D_\nu \Psi_0 = 0$.

Neyman orthogonality

Theorem (Neyman orthogonality)

If $\Psi(P) = \Psi_0(P, \nu) = P[\varphi(O, \nu(P))]$ and $\varphi(\cdot, \nu) - P[\varphi(O, \nu)]$ is the canonical gradient of (\mathcal{P}, Ψ) then $D_\nu \Psi_0 = 0$.

Debiasing

The *first order* bias, coming from $\Psi_0(P, \hat{\nu}_n) - \Psi_0(P, \nu)$, is removed.

Efficiency

Efficiency

Definition (RAL estimators)

An estimator $\hat{\theta}_n$ of the parameter $\theta = \Psi(P)$ under the model \mathcal{P} , is called *asymptotically linear* with *influence function* $IF(\cdot, P) \in \mathcal{L}_P^2$, if $P[IF(O, P)] = 0$ for all $P \in \mathcal{P}$, and

$$\hat{\theta}_n - \theta = \hat{\mathbb{P}}_n[IF(O, P)] + o_P(n^{-1/2}).$$

Efficiency

Definition (RAL estimators)

An estimator $\hat{\theta}_n$ of the parameter $\theta = \Psi(P)$ under the model \mathcal{P} , is called *asymptotically linear* with *influence function* $IF(\cdot, P) \in \mathcal{L}_P^2$, if $P[IF(O, P)] = 0$ for all $P \in \mathcal{P}$, and

$$\hat{\theta}_n - \theta = \hat{\mathbb{P}}_n[IF(O, P)] + o_P(n^{-1/2}).$$

Theorem (Efficient influence function)

The RAL estimator with lowest possible asymptotic variance has the canonical gradient as its influence function.

Constructing estimators 1: Solve the efficient score equation

Constructing estimators 1: Solve the efficient score equation

Find a parametrization $\Psi(P) = P[\varphi(O, \nu)]$ such that φ is the (canonical) gradient.

Constructing estimators 1: Solve the efficient score equation

Find a parametrization $\Psi(P) = P[\varphi(O, \nu)]$ such that φ is the (canonical) gradient. Then by Neyman orthogonality and assumptions we can write

$$\begin{aligned}\sqrt{n}(\hat{\theta}_n - \theta) &= \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] \\ &\quad + D_\nu \Psi_0[\sqrt{n}(\hat{\nu}_n - \nu)] \quad = 0 \\ &\quad + \mathcal{O}_P(\sqrt{n}\|\hat{\nu}_n - \nu\|_\nu^2)\end{aligned}$$

Constructing estimators 1: Solve the efficient score equation

Find a parametrization $\Psi(P) = P[\varphi(O, \nu)]$ such that φ is the (canonical) gradient. Then by Neyman orthogonality and assumptions we can write

$$\begin{aligned}\sqrt{n}(\hat{\theta}_n - \theta) &= \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] &&= \mathbb{G}_n[\varphi(O, \nu)] \\ &+ D_\nu \Psi_0[\sqrt{n}(\hat{\nu}_n - \nu)] &&= 0 \\ &+ \mathcal{O}_P(\sqrt{n}\|\hat{\nu}_n - \nu\|_\nu^2) &&= \mathcal{O}_P(1)\end{aligned}$$

Constructing estimators 1: Solve the efficient score equation

Find a parametrization $\Psi(P) = P[\varphi(O, \nu)]$ such that φ is the (canonical) gradient. Then by Neyman orthogonality and assumptions we can write

$$\begin{aligned}\sqrt{n}(\hat{\theta}_n - \theta) &= \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] &&= \mathbb{G}_n[\varphi(O, \nu)] \\ &\quad + D_\nu \Psi_0[\sqrt{n}(\hat{\nu}_n - \nu)] &&= 0 \\ &\quad + \mathcal{O}_P(\sqrt{n}\|\hat{\nu}_n - \nu\|_\nu^2) &&= \mathcal{O}_P(1) \\ &= \mathbb{G}_n[\varphi(O, \nu)] + \mathcal{O}_P(1).\end{aligned}$$

Constructing estimators 1: Solve the efficient score equation

Find a parametrization $\Psi(P) = P[\varphi(O, \nu)]$ such that φ is the (canonical) gradient. Then by Neyman orthogonality and assumptions we can write

$$\begin{aligned}\sqrt{n}(\hat{\theta}_n - \theta) &= \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] &&= \mathbb{G}_n[\varphi(O, \nu)] \\ &\quad + D_\nu \Psi_0[\sqrt{n}(\hat{\nu}_n - \nu)] &&= 0 \\ &\quad + \mathcal{O}_P(\sqrt{n}\|\hat{\nu}_n - \nu\|_\nu^2) &&= \mathcal{O}_P(1) \\ &= \mathbb{G}_n[\varphi(O, \nu)] + \mathcal{O}_P(1).\end{aligned}$$

Hence $\hat{\theta}_n$ is a RAL estimator, and if $\varphi - P[\varphi]$ is the canonical gradient it will be *asymptotically efficient*.

Constructing estimators 1: Solve the efficient score equation

Find a parametrization $\Psi(P) = P[\varphi(O, \nu)]$ such that φ is the (canonical) gradient. Then by Neyman orthogonality and assumptions we can write

$$\begin{aligned}\sqrt{n}(\hat{\theta}_n - \theta) &= \mathbb{G}_n[\varphi(O, \hat{\nu}_n)] &&= \mathbb{G}_n[\varphi(O, \nu)] \\ &\quad + D_\nu \Psi_0[\sqrt{n}(\hat{\nu}_n - \nu)] &&= 0 \\ &\quad + \mathcal{O}_P(\sqrt{n}\|\hat{\nu}_n - \nu\|_\nu^2) &&= \mathcal{O}_P(1) \\ &= \mathbb{G}_n[\varphi(O, \nu)] + \mathcal{O}_P(1).\end{aligned}$$

Hence $\hat{\theta}_n$ is a RAL estimator, and if $\varphi - P[\varphi]$ is the canonical gradient it will be *asymptotically efficient*.

This is the approach taken in Chernozhukov et al. [2018]. See also Example 4.1 of the note.

Constructing estimators 2: TMLE

References

- V. Chernozhukov, D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins. Double/debiased machine learning for treatment and structural parameters, 2018.
- E. H. Kennedy. Semiparametric theory and empirical processes in causal inference. In *Statistical causal inferences and their applications in public health research*, pages 141–167. Springer, 2016.