EXPERIMENT-1


AIM: (i) To determine the acceleration due to gravity (g) by means of a compound pendulum.

(ii) To determine radius of gyration about an axis through the center of gravity for the compound pendulum.

APPARATUS: (i) A bar pendulum, (ii) a knife—edge with a platform, (iii) a sprit level, (iv) a precision stop watch, (v) a meter scale

THEORY: If the bar be allowed to oscillate about horizontal knife edge passing in each of the holes in turn, beginning with the nearest to center of gravity, and a graph plotted taking time period of oscillations as ordinates and corresponding distances of the axis of suspension from the end of the bar as abscissa, the graph will consist of two symmetrical branches.

FORMULA USED:




OBSERVATION:

Number of	Distance from	Time (for 10	Time Period (in
holes from	COG (in cm)	oscillations in	seconds)
center		seconds)	
1	5	25.16	2.516
2	10	18.87	1.887
3	15	16.42	1.642
4	20	15.30	1.530
5	25	15.28	1.528
6	30	15.02	1.502
7	35	15.20	1.520
8	40	15.40	1.540
9	45	16.10	1.610

For Inverted Pendulum:

Number of	Distance from	Time (for 10	Time Period (in
holes from	COG (in cm)	oscillations in	seconds)
center		seconds)	
1	5	25.88	2.588
2	10	18.90	1.890
3	15	15.34	1.534
4	20	15.18	1.518
5	25	15.10	1.510
6	30	15.20	1.520
7	35	15.34	1.534
8	40	15.97	1.597
9	45	16.13	1.613

GRAPH:



CALCULATIONS:

$$J = 57.5 \text{ cm} \qquad J_2 = 58.5 \text{ cm}$$

$$J = J_1 + J_2 = 0.58 \text{ m}$$

$$T = 0R = 1.65 = 2\pi / J_q$$

$$= g = 4\pi^2 J_z = 8.935 \text{ m/s}^2$$

$$K = \sqrt{J_1 \times J_2} = \sqrt{3363.75}$$

$$= 57.997 \text{ cm}$$

$$= 57.997 \text{ cm}$$

$$= 6.82 \text{ m}$$

$$= 8.82 \text{ m}$$

RESULT: 1) Acceleration due to gravity is 8.935 m/s²

2) Radius of gyration is 57.997cm

SOURCES OF ERROR:1) The knife edge may not be horizontal.

- 2) Air resistance could damp the oscillation.
- 3)Incorrect measurement of time.
- 4) Graph may be incorrectly plotted.

By: Amogh Garg

2020UCO1688

COE(Section-3)