Gemeinsamer Bundesausschuss

Resolution

of the Federal Joint Committee on an Amendment of the Pharmaceuticals Directive:
Annex XII – Benefit Assessment of Medicinal Products with New Active Ingredients according to Section 35a SGB V
Tiratricol (peripheral thyrotoxicosis with monocarboxylate transporter 8 (MCT8) deficiency (Allan-Herndon-Dudley Syndrome))

of 16 October 2025

At their session on 16 October 2025, the Federal Joint Committee (G-BA) resolved to amend the Pharmaceuticals Directive (AM-RL) in the version dated 18 December 2008 / 22 January 2009 (Federal Gazette, BAnz. No. 49a of 31 March 2009), as last amended by the publication of the resolution of D Month YYYY (Federal Gazette, BAnz AT DD.MM.YYYY BX), as follows:

I. Annex XII shall be amended in alphabetical order to include the active ingredient Tiratricol as follows:

Courtesy translation – only the German version is legally binding.

Tiratricol

Resolution of: 16 October 2025
Entry into force on: 16 October 2025
Federal Gazette, BAnz AT DD. MM YYYY Bx

Therapeutic indication (according to the marketing authorisation of 12 February 2025):

Emcitate is indicated for the treatment of peripheral thyrotoxicosis in patients with monocarboxylate transporter 8 (MCT8) deficiency (Allan-Herndon-Dudley Syndrome), from birth.

Therapeutic indication of the resolution (resolution of 16 October 2025):

See therapeutic indication according to marketing authorisation.

1. Extent of the additional benefit and significance of the evidence

Tiratricol is approved as a medicinal product for the treatment of rare diseases in accordance with Regulation (EC) No. 141/2000 of the European Parliament and the Council of 16 December 1999 on orphan drugs. In accordance with Section 35a, paragraph 1, sentence 11, 1st half of the sentence SGB V, the additional medical benefit is considered to be proven through the grant of the marketing authorisation.

The G-BA determine the extent of the additional benefit for the number of patients and patient groups for which there is a therapeutically significant additional benefit in accordance with Chapter 5 Section 12, paragraph 1, number 1, sentence 2 of its Rules of Procedure (VerfO) in conjunction with Section 5, paragraph 8 Ordinance on the Benefit Assessment of Pharmaceuticals (AM-NutzenV), indicating the significance of the evidence. This quantification of the additional benefit is based on the criteria laid out in Chapter 5 Section 5, paragraph 7, numbers 1 to 4 of the Rules of Procedure (VerfO).

Subjects with monocarboxylate transporter 8 (MCT8) deficiency (Allan-Herndon-Dudley Syndrome), for the treatment of peripheral thyrotoxicosis from birth

Extent of the additional benefit and significance of the evidence of tiratricol:

Hint for a non-quantifiable additional benefit since the scientific data does not allow quantification.

Study results according to endpoints:[1]

Subjects with monocarboxylate transporter 8 (MCT8) deficiency (Allan-Herndon-Dudley Syndrome), for the treatment of peripheral thyrotoxicosis from birth

[1] Data from the dossier assessment of the G-BA (published on 1. August 2025), unless otherwise indicated.

Courtesy translation – only the German version is legally binding.

Summary of results for relevant clinical endpoints

Endpoint category	Direction of effect/ risk of bias	Summary
Mortality	n.a.	The data are not assessable.
Morbidity	n.a.	The data are not assessable.
Health-related quality of life	∅	No data available.
Side effects	n.a.	The data are not assessable.

Explanations:
↑: statistically significant and relevant positive effect with low/unclear reliability of data
↓: statistically significant and relevant negative effect with low/unclear reliability of data
↑↑: statistically significant and relevant positive effect with high reliability of data
↓↓: statistically significant and relevant negative effect with high reliability of data
↔: no statistically significant or relevant difference
∅: No data available.
n.a.: not assessable

TRIAC I study: single-arm Phase IIb study over 12 months

Mortality

Endpoint	Tiratricol	
	N[a]	Patients with event n (%)
Overall mortality		
	46	1 (2.2)

Morbidity

Endpoint	Tiratricol			
	N[b]	Baseline value MV (SD)	Value in month 12 MV (SD)	Change from baseline to month 12 MV (SD)
T3 concentration in serum in nmol/l (presented additionally; primary endpoint)				
	45	4.97 (1.55)	1.82 (0.69)	-3.15 (1.56)
Physical development, z-score[c]				
Body weight	45	-2.85 (1.90)	-2.63 (1.74)	0.22 (0.77)
Body height	45	-1.96 (1.58)	-1.98 (1.47)	-0.02 (0.53)

Courtesy translation – only the German version is legally binding.

Side effects

Endpoint MedDRA system organ classes/ AEs of special interest	**Active ingredient**	
	N[a]	**Patients with event n (%)**
Total adverse events (presented additionally)	46	43 (93.5)
Severe AEs (presented additionally)	46	12 (26.1)
Serious adverse events (SAE)	46	18 (39.1)
Therapy discontinuation due to adverse events	46	1 (2.2)
Severe adverse events according to MedDRA system organ class (with an incidence ≥ 10%) (presented additionally)		
Infections and infestations	46	6 (13.0)
SAEs according to MedDRA system organ class (with an incidence ≥ 10%)		
Infections and infestations	46	10 (21.7)
a. Safety population b. Intention-to-treat population c. In relation to the WHO healthy reference population Abbreviations used: MedDRA = Medical Dictionary for Regulatory Activities; MV = mean value; N = number of patients evaluated; n = number of patients with (at least one) event; n.c. = not calculable; SD = standard deviation; SAE = serious adverse event; AE = adverse event		

2. Number of patients or demarcation of patient groups eligible for treatment

Subjects with monocarboxylate transporter 8 (MCT8) deficiency (Allan-Herndon-Dudley Syndrome), for the treatment of peripheral thyrotoxicosis from birth

Approx. 40 – 110 patients

Courtesy translation – only the German version is legally binding.

3. Requirements for a quality-assured application

The requirements in the product information are to be taken into account. The European Medicines Agency (EMA) provides the contents of the product information (summary of product characteristics, SmPC) for Emcitate (active ingredient: tiratricol) at the following publicly accessible link (last access: 24 July 2025):

https://www.ema.europa.eu/en/documents/product-information/emcitate-epar-product-information_en.pdf

Treatment with tiratricol should only be initiated and monitored by specialists who are experienced in the treatment of subjects with rare genetic disorders such as MCT8 deficiency.

4. Treatment costs

Annual treatment costs:

Subjects with monocarboxylate transporter 8 (MCT8) deficiency (Allan-Herndon-Dudley Syndrome), for the treatment of peripheral thyrotoxicosis from birth

Designation of the therapy	Annual treatment costs/ patient
Medicinal product to be assessed:	
Tiratricol *Additionally required SHI services*	€ 44,216.13 - € 972,754.87 *Different from patient to patient*

Costs after deduction of statutory rebates (LAUER-TAXE® as last revised: 15 August 2025)

5. Designation of medicinal products with new active ingredients according to Section 35a, paragraph 3, sentence 4 SGB V that can be used in a combination therapy with the assessed medicinal product

In the context of the designation of medicinal products with new active ingredients pursuant to Section 35a, paragraph 3, sentence 4 SGB V, the following findings are made:

Subjects with monocarboxylate transporter 8 (MCT8) deficiency (Allan-Herndon-Dudley Syndrome), for the treatment of peripheral thyrotoxicosis from birth

– No medicinal product with new active ingredients that can be used in a combination therapy that fulfils the requirements of Section 35a, paragraph 3, sentence 4 SGB V.

The designation of combinations exclusively serves the implementation of the combination discount according to Section 130e SGB V between health insurance funds and pharmaceutical companies. The findings made neither restrict the scope of treatment required to fulfil the medical treatment mandate, nor do they make statements about expediency or economic feasibility.

Courtesy translation – only the German version is legally binding.

6. Percentage of study participants at study sites within the scope of SGB V in accordance with Section 35a, paragraph 3, sentence 5 SGB V

The medicinal product Emcitate is a medicinal product placed on the market from 1 January 2025.

The percentage of study participants in the clinical studies of the medicinal product conducted or commissioned by the pharmaceutical company in the therapeutic indication to be assessed who participated at study sites within the scope of SGB V (German Social Security Code) is ≥ 5% of the total number of study participants.

The clinical studies of the medicinal product in the therapeutic indication to be assessed were therefore conducted to a relevant extent within the scope of SGB V.

II. The resolution will enter into force on the day of its publication on the website of the G-BA on 16 October 2025.

The justification to this resolution will be published on the website of the G-BA at www.g-ba.de.

Berlin, 16 October 2025

Federal Joint Committee (G-BA)
in accordance with Section 91 SGB V
The Chair

Prof. Hecken

Courtesy translation – only the German version is legally binding.