

# Implementation of poloidal density variation in collision operator

Aylwin Iantchenko

December 5, 2017

The aim of this document is to explain the modifications in **SFINCS** that have been done, in order to include poloidal density variation in the collision operator (in both full Fokker-Planck and pure pitch angle scattering operator options).

As is explained in Ref. [1] poloidal density variation can be included by modifying the lowest order distribution function

$$f_{Ms}(\psi) \rightarrow f_{0s} = f_{Ms}(\psi) e^{-Z_s e \Phi_1(\theta, \zeta) / T_s},$$

where  $f_M(\psi)$  is the flux-function Maxwellian,  $\Phi_1$  is first order variation of the electrostatic potential, and  $Z_s, T_s$  are the charge and temperature of species  $s$  respectively. To implement this change one may simply modify the species density accordingly

$$n_s \rightarrow n_s e^{-Z e \Phi_1(\theta, \zeta) / T}, \quad (1)$$

in the original equations for the various terms in the collision operator, which are described in Ref. [3]. The linearised collision operator takes the new form

$$\begin{aligned} C_{ab}^{L:f0} = & C_{ab} \{f_{aM}, f_{bM}\} e^{-(Z_a/T_a + Z_b/T_b) e \Phi_1(\theta, \zeta)} + C_{ab} \{f_{a1}, f_{bM}\} e^{-Z_b/T_b e \Phi_1(\theta, \zeta)} + \\ & + C_{ab} \{f_{aM}, f_{b1}\} e^{-Z_a/T_a e \Phi_1(\theta, \zeta)}, \end{aligned} \quad (2)$$

which describes collisions between species  $a$  and  $b$ . In normalised **SFINCS** units this becomes

$$\begin{aligned} \hat{C}_{ab}^{L:f0} = & \hat{C}_{ab} \{f_{aM}, f_{bM}\} e^{-(\hat{Z}_a/\hat{T}_a + \hat{Z}_b/\hat{T}_b) \alpha \hat{\Phi}_1(\theta, \zeta)} + \hat{C}_{ab} \{f_{a1}, f_{bM}\} e^{-\hat{Z}_b/\hat{T}_b \alpha \hat{\Phi}_1(\theta, \zeta)} + \\ & + \hat{C}_{ab} \{f_{aM}, f_{b1}\} e^{-\hat{Z}_a/\hat{T}_a \alpha \hat{\Phi}_1(\theta, \zeta)}. \end{aligned} \quad (3)$$

The first term on the RHS of Eq. (3) is the *temperature equilibration term*. Since this term does not include  $f_{1s}$  it has to be treated differently compared to the other two cases, as will be seen shortly. In the code, the calculation of the residual (see Ref. [2] for definition of the residual and Jacobian in this context) has to be modified to include the factors specified in Eq. (3).

The contribution to the Jacobian are  $\delta \hat{C}_{ab}^{L:f0} / \delta \hat{f}_{1s}$  and  $\delta \hat{C}_{ab}^{L:f0} / \delta \hat{\Phi}_1$ . The first contribution is the same as in Eq. (3) (without the temperature equilibration term) except that the collision operator does not act on the distribution function. In the second case we get

$$\begin{aligned} \frac{\delta \hat{C}_{ab}^{L:f0}}{\delta \hat{\Phi}_1} = & - \left( \hat{Z}_a / \hat{T}_a + \hat{Z}_b / \hat{T}_b \right) \alpha \hat{C}_{ab} \{f_{aM}, f_{bM}\} e^{-\left( \hat{Z}_a / \hat{T}_a + \hat{Z}_b / \hat{T}_b \right) \alpha \hat{\Phi}_1(\theta, \zeta)} \\ & - \hat{Z}_b / \hat{T}_b \alpha \hat{C}_{ab} \{f_{a1}, f_{bM}\} e^{-\hat{Z}_b / \hat{T}_b \alpha \hat{\Phi}_1(\theta, \zeta)} - \hat{Z}_a / \hat{T}_a \alpha \hat{\Phi}_1(\theta, \zeta) \hat{C}_{ab} \{f_{aM}, f_{b1}\} e^{-\hat{Z}_a / \hat{T}_a \alpha \hat{\Phi}_1(\theta, \zeta)}. \end{aligned} \quad (4)$$

Apart from the different pre-factor, the term in Eq. (4) differs from term in  $\delta \hat{C}_{ab}^{L:f0} / \delta \hat{f}_{1s}$  by acting on the first order distribution function  $f_{1s}$ . For example, in the first case, terms such as  $\partial f_{1s} / \partial x_s$  become

$$\frac{\partial}{\partial f_{1s}} \frac{\partial f_{1s}}{\partial x_s} = \frac{\partial}{\partial x_s},$$

whereas in the second case, when we take the derivative with respect to  $\Phi_1$ , we have to include the derivative  $\partial f_{1s} / \partial x_s$  (and  $\partial f_{0s} / \partial x_s$  for temperature equilibration, which again is not present in the first contribution), as well.

## Implementation in the Code

To implement Eq. (3) and Eq. (4) we need to modify the evaluation of the residual and the Jacobian. The residual is evaluated in `evaluateResidual.F90` and the Jacobian in `evaluateJacobian.F90`. Both these two routines call the functions in `PopulateMatrix.F90`, where the actual assembly of the matrices used to compute the residual (by multiplying the matrix with the state-vector) and Jacobian, is done. As a consequence of this structure, we only need to modify `PopulateMatrix.F90` in order to include poloidal density variation in the collision operator.

## Residual

In the residual we have to include the extra pre-factor appearing in Eq. (3). Since this factor is species specific, we have to keep track of which species density is appearing in the equations. The pitch-angle scattering and the  $\hat{C}_{ab}^E \{f_{a1}\}$  parts of the collision operator (see Ref. [3]) both uses the density `nHats(iSpeciesB)` while the other contributions to the total collision operator `CHat` uses `nHats(iSpeciesA)` instead.

Rather than creating many if-statements in the current implementation of the collision operator, for sake of clarity we separate the two implementations with one if-statement directly before the various contributions to the collision operator are calculated. If `Phi1` is included and `poloidalVariationInCollisionOperator = .true.` the contribution from the collision operator is calculated taking poloidal density variation into account.

```
if (poloidalVariationInCollisionOperator .and. includePhi1 .and.
    includePhi1InKineticEquation) then
```

Otherwise the second block is executed, which contains the original code. Since we now will have a `Phi1Hat` dependence in the various terms of the collision operator we will have to iterate over `itheta` and `izeta` when calculating the various contributions. Before calculating each term, we define a pre-factor (`PreFactor`) as is specified in Eq. (3), taking the correct species index into account.

```
do itheta=ithetaMin,ithetaMax
  do izeta=izetaMin,izetaMax
    ! Generate preFactor for nHats(iSpeciesA) terms
    preFactor =
      exp(-Zs(iSpeciesA)*alpha*Phi1Hat(itheta,izeta)/Thats(iSpeciesA))
```

When required, this pre-factor is overwritten using the charge and temperature of `iSpeciesB` instead. Because of the `itheta`, `izeta` dependence we have replaced `CECD` and `nuDHat` to their equivalents `CECDpol` and `nuDHatpol` with space for their `itheta` and `izeta` components.

The rest of this block is the same as in the original code, except that we do the second iteration over `itheta`, `izeta` slightly earlier, and include the appropriate `PreFactor` in front of additional contributions to the total collision operator `CHat`.

### Temperature equilibration term

In the original version of the code, the temperature equilibration term is calculated in `evaluateResidual.F90` by using `whichMatrix = 2` when calling `PopulateMatrix.F90`. The output is then multiplied with  $f_{0s}$ . Since the original version of `SFINCS` already includes  $\Phi_1$  in the definition of  $f_{0s}$  (in the subroutine `init_f0()` inside `PopulateMatrix.F90`), the same routines as were presented above can be used also if `includeTemperatureEquilibrationTerm = .true..`

### Jacobian

If the modifications explained in the previous section are implemented, the  $\delta C / \delta f_{1s}$  contribution to the Jacobian is at this point already taking the density variations into account. We need to add the calculation of  $\delta C / \delta \Phi_1$ .

#### The $\delta C / \delta \Phi_1$ term

We begin with treating the two latter terms appearing in Eq. (4) and will then look at the temperature equilibration term.

To implement the derivative with respect to  $\Phi_1$  we need to redo the calculation of the various contributions to the collision operator, but including a new pre-factor, `PreFactorJ` as specified in Eq. (4). This should only be done when the Jacobian (`whichmatrix = 1`) or the preconditioner (`whichmatrix = 0`) is calculated.

```
if (whichMatrix == 1 .or. whichMatrix == 0) then
  ! Generate preFactorJ for nHats(iSpeciesB) terms
  preFactorJ = (-Zs(iSpeciesA)*alpha/Thats(iSpeciesA)) &
```

```
*exp(-Zs(iSpeciesA)*alpha*Phi1Hat(itheta,izeta)/Thats(iSpeciesA))
```

Again, the same is done using the charge and temperature of `iSpeciesB` instead, whenever required. The result is saved in the new matrices `nuDHatpolJ`, `CECDpolJ` etc. Same calculations are then repeated for these terms (except for the different pre-factor), and the result is saved in `CHatJ`.

In contrary to the residual (and the  $\delta C/\delta f_{1s}$  terms), when calculating  $\delta C/\delta \Phi_1$  we have to include the first order distribution function before saving the result into the main matrix. For this purpose we first have to initiate this distribution function, by reading the appropriate terms in the current state vector.

```
do ix= max(ixMinCol,min_x_for_L(L)),Nx
  ! Generate flb from state vector
  index = getIndex(iSpeciesB,ix,L+1,itheta,izeta,BLOCK_F)
  flb(ix) = stateArray(index + 1)
```

We multiply the terms in the collision operator with this distribution function.

```
CHatTimesf = matmul(CHatJ,flb)
```

The result is then saved into the  $\delta \hat{C}/\delta \hat{\Phi}_1$  block of the main matrix,

```
do ix_row=max(ixMin,min_x_for_L(L)),Nx
  rowIndex=getIndex(iSpeciesA,ix_row,L+1,itheta,izeta,BLOCK_F)
  ! Get column index for the d/dPhi1 terms
  colIndex=getIndex(1,1,1,itheta,izeta,BLOCK_QN)
  ! Save into the main matrix, note that here we only use ix_row since
    CHatTimesf is now a vector
  call MatSetValue(matrix, rowIndex, colIndex, &
    -nu_n*CHatTimesf(ix_row), ADD_VALUES, ierr)
  ! need to use MatSetValue, otherwise petsc gives error
end do ! ix_row
```

## Temperature equilibration

If the temperature equilibration terms should be included, we have to add additional terms into the main matrix, when calculating  $\delta \hat{C}/\delta \hat{\Phi}_1$ . These terms are different from the other contributions since now we need to multiply the output with  $f_{0s} = f_{Ms}(\psi)e^{-Z_s e \Phi_1(\theta, \zeta)/T_s}$  (instead of the first order distribution function  $f_{1s}$ ). Because of this, we need to generate the Maxwellian distribution function for `iSpeciesB`.

```
if (includeTemperatureEquilibrationTerm .and. L==0) then
  fM(ix) =
    sqrt(mhats(iSpeciesB)/Thats(iSpeciesB))*mhats(iSpeciesB)/Thats(iSpeciesB)&
    * nhats(iSpeciesB)/(pi*sqrt(pi)*expb2(ix)
end if
```

Because of the `Phi1Hat` dependence in the zeroth order distribution function we get a different contribution to the Jacobian, as specified in the first term of Eq. (4). In the

code, we need to use both the residual term `CHat` and the Jacobian terms `CHatJ` to obtain the correct result.

```
CHatTimesf = matmul((CHatJ + CHat*(-Zs(iSpeciesB)*alpha/Thats(iSpeciesB))&
    )*exp(-Zs(iSpeciesB)*alpha*Phi1Hat(itheta,izeta)/Thats(iSpeciesB)),fM)
```

Written in terms of operators we have that `CHat` is  $\hat{C}_{ab}\{f_{aM}, \cdot\} e^{-\hat{Z}_a/\hat{T}_a \alpha \hat{\Phi}_1(\theta, \zeta)}$  and `CHatJ` is  $\hat{C}_{ab}\{f_{aM}, \cdot\} \left(-\hat{Z}_a/\hat{T}_a \alpha\right) e^{-\hat{Z}_a/\hat{T}_a \alpha \hat{\Phi}_1(\theta, \zeta)}$ . Here  $\cdot$  means that the operator does not act on anything. What is left is to add this term into the main matrix, using the the same procedure as was explained in the previous section.

## Pure pitch angle scattering operator

The implementation of poloidal density variation in the case the pure pitch angle scattering collision operator is used (corresponding to the case `collisionoperator = 1`), is slightly easier because of the existing loop over `itheta`, `izeta` and since we are working with a scalar `CHat_element`. Also, since  $L > 0$  always for this term, no temperature equilibration has to be included. The only block we need to change is when we save the collision operator into the main matrix.

We introduce the factor `preFactor` which is in normal cases equal to one, but when `poloidalVariationInCollisionOperator = .true.` it is equal to the exponential pre-factor containing  $\Phi_1$  in Eq. (3).

Before saving into the main matrix, we multiply the collision operator with this factor.

```
call MatSetValue(matrix, index, colIndex, &
    -nu_n*CHat_element*preFactor, ADD_VALUES, ierr)
```

## Jacobian, the $\delta C/\delta \Phi_1$ term

Just as before, the  $\delta C/\delta f_1$  terms are the same as in the residual. What is different is the  $\delta C/\delta \Phi_1$  term. The procedure to calculate this term is however very similar, to the case with the residual. The only difference is the different pre-factor which now also is multiplied with the distribution function from the state vector.

```
preFactor = -Zs(iSpeciesA)*alpha/Thats(iSpeciesA)*exp(-Zs(iSpeciesA) &
    *alpha*Phi1Hat(itheta,izeta)/Thats(iSpeciesA))*stateArray(index + 1)
```

## Current state of the project

Presently, the modifications discussed in the previous section have been implemented into the “`poloidalVariationInCollisionOperator`” branch of `SFINCS`. What remains is to benchmark the output with theoretical predictions, using for example the model described

in Ref. [4]<sup>1</sup>. Note that in this case only tokamak cases are considered.

The first step is to test the model with the output from the original **SFINCS** code, without including poloidal density variation. By setting the temperature gradient to zero we remove the effect from the collision operator and expect therefore the output from **SFINCS** to agree reasonably well with the predictions from the model. Example of typical **SFINCS**-input used to get the best agreement so far is:

```
&speciesParameters
Zs = 1.0, 20.0
mHats = 1.0, 20.0
nHats = 4.0, 0.012
THats = 2.0, 2.0
dnHatdrNs = -40.0, -0.12
dTHatdrNs = -0.0, -0.0

&geometryParameters
inputRadialCoordinate = 3
inputRadialCoordinateForGradients = 3

rN_wish = 0.3

geometryScheme = 1
B0OverBBar = 60.0
GHat = 180.0
IHat = 0.9
iota = 0.5
epsilon_t = 0.1
epsilon_h = 0
epsilon_antisymm = 0
helicity_l = 1
helicity_n = 1

psiAHat = 30.0
aHat = 1.0
/
```

For this input, the orderings of various parameters described in Ref. [4] are reasonably satisfied (we have  $\Delta \sim 0.3$ ,  $\delta \sim 0.15$ ,  $\epsilon \sim 0.1$ ,  $n_z Z^2/n_i \sim 1.2$  and  $\nu^* \sim 0.11$ ). If we use this input for varying magnetic field strength  $B_0$  (as an example) and calculate the sine and cosine parts of the perturbed impurity density, we get Fig. 1 and Fig. 2 respectively. Solid lines correspond to the case when **SFINCS** output is used to calculate the perturbed density while dashed lines correspond to the analytical model. The two outputs show good agreement for the cosine parts while the sine is slightly deviating. Same scan using the new code (where poloidal density variation is included) gives exactly the same result.

---

<sup>1</sup>Note that in this model the collision operator is not linearised around a poloidally varying zeroth order distribution function (as is done here), but the poloidal variation enters through the perturbed distribution function in the ion-impurity collision operator, when computing the parallel friction force.

Some convergence issues appear when using the new implementation with high impurity charge (in this regime for  $Z_s > 20$ ). No poloidal asymmetries have been observed so far, when using input to **SFINCS** which satisfies the orderings described in Ref. [4] (which is required if we want to compare the **SFINCS** output with the model).




Figure 1: Sine part of the perturbed density for a scan in the magnetic field strength  $B_0$ . Solid line shows result calculated from **SFINCS** output while the dashed line is the prediction from the model.




Figure 2: Cosine part of the perturbed density for a scan in the magnetic field strength  $B_0$ . Solid line shows result calculated from **SFINCS** output while the dashed line is the prediction from the model

## References

- [1] S. Buller, *Poloidal variation in collision operator* (2017).
- [2] A. Mollén, *Implementation of  $\Phi_1$  in SFINCS* (2016).
- [3] M. Landreman, *Technical Documentation for version 3 of SFINCS* (2014).
- [4] T. Fülöp, P. Helander *Phys. Plasmas* **6**, 3066 (1999)

.