

Functions of Linear Regression

Xueyuan Gong
Department of Computer and Information Science
Faculty of Science and Technology
University of Macau
Macau, China
{yb47453}@umac.mo

Given a data set $X = \{X_j^i | X_j^i \in \mathbb{R}^{n \times (m-1)}\}$ and a label set $Y = \{Y^i | Y^i \in \mathbb{R}^{n \times 1}\}$, where $1 \leq i \leq n$ and $2 \leq j \leq m$. For all X^i , insert $X_1^i = 1$ so that $X_j^i \in \mathbb{R}^{n \times m}$. Initialize weight matrix $W = \{W_j | W_j \in \mathbb{R}^{1 \times m}\}$.

Hypothesis function is defined in Equation (1).

$$H(X^i) = \sum_{j=1}^m W_j X_j^i \quad (1)$$

Cost function is defined in Equation (2).

$$C(X^i) = H(X^i) - Y^i \quad (2)$$

Objective function is defined in Equation (3).

$$\begin{aligned} J(W) &= \frac{1}{2n} \sum_{i=1}^n C^2(X^i) \\ &= \frac{1}{2n} \sum_{i=1}^n \left(\sum_{j=1}^m W_j X_j^i - Y^i \right)^2 \end{aligned} \quad (3)$$

The partial derivative of $J(W)$ with respect to W_j is given as follows:

$$\begin{aligned} \frac{\partial J(W)}{\partial W_j} &= \frac{1}{2n} \sum_{i=1}^n \frac{\partial C^2(X^i)}{\partial W_j} \\ &= \frac{1}{2n} \sum_{i=1}^n \frac{\partial C^2(X^i)}{\partial C(X^i)} \cdot \frac{\partial C(X^i)}{\partial H(X^i)} \cdot \frac{\partial H(X^i)}{\partial W_j} \\ &= \frac{1}{2n} \sum_{i=1}^n 2C(X^i) \cdot 1 \cdot X_j^i \\ &= \frac{1}{n} \sum_{i=1}^n (H(X^i) - Y^i) X_j^i \end{aligned}$$

Thus, for all W_j , the update function is given as follows:

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.

Copyright 20XX ACM X-XXXXX-XX-X/XX/XX ...\$15.00.