

# Discrete Structures— Propositional Logic Pt. 2

Dr. Ab Mosca (they/them)

# Plan for Today

- Boolean Algebra
- Propositions
- Quantifiers

One way we model truth values of statements is with **truth tables**.

conjunction

$P$	$Q$	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

disjunction

$P$	$Q$	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

implication

$P$	$Q$	$P \rightarrow Q$
T	T	T
T	F	F
F	T	T
F	F	T

biconditional negation

$P$	$Q$	$P \leftrightarrow Q$
T	T	T
T	F	F
F	T	F
F	F	T

$P$	$\neg P$
T	F
F	T

Warm Up

We also use truth tables to show logical equivalence. Two molecular statements  $P$  and  $Q$  are **logically equivalent** provided  $P$  is true precisely when  $Q$  is true.

Work with 1-2 other people to show that  $\neg(P \wedge Q)$  is logically equivalent to  $\neg P \vee \neg Q$ .

## De Morgan's Laws

Last time we showed

$\neg(P \vee Q)$  is logically equivalent to  $\neg P \wedge \neg Q$ .

You just showed

$\neg(P \wedge Q)$  is logically equivalent to  $\neg P \vee \neg Q$ .

## De Morgan's Laws

$\neg(P \vee Q)$  *is logically equivalent to*  $\neg P \wedge \neg Q$ .

$\neg(P \wedge Q)$  *is logically equivalent to*  $\neg P \vee \neg Q$ .

These equivalences are called ***De Morgan's Laws***.

## De Morgan's Laws

$\neg(P \vee Q)$  *is logically equivalent to*  $\neg P \wedge \neg Q$ .

$\neg(P \wedge Q)$  *is logically equivalent to*  $\neg P \vee \neg Q$ .

These equivalences are called ***De Morgan's Laws***.

Notice that these laws show you how to “distribute” a negation into parentheses.

# Implications are Disjunctions

Last time we also showed that the implication  $P \rightarrow Q$  is logically equivalent to the disjunction  $\neg P \vee Q$ .

# Implications are Disjunctions

Last time we also showed that the implication  $P \rightarrow Q$  is logically equivalent to the disjunction  $\neg P \vee Q$ .

That means at any time we see  $P \rightarrow Q$  in a statement we can replace it with  $\neg P \vee Q$  without changing the truth values of the statement. And vice versa.



# Implications are Disjunctions

Last time we also showed that the implication  $P \rightarrow Q$  is logically equivalent to the disjunction  $\neg P \vee Q$ .

That means at any time we see  $P \rightarrow Q$  in a statement we can replace it with  $\neg P \vee Q$  without changing the truth values of the statement. And vice versa.

(The same is true of De Morgan's Laws, or any logically equivalent statements; we can always replace one with another without changing the truth values of a statement.)

## Double Negation

A double negation cancels itself out.

In other words,  $\neg\neg P$  is logically equivalent to  $P$ .

# Boolean Algebra

***Boolean Algebra*** is the set of rules dictating how to transform one statement into another.

# Boolean Algebra

***Boolean Algebra*** is the set of rules dictating how to transform one statement into another.

Another way to show logical equivalence (besides truth tables) is with ***Boolean Algebra***.

# Boolean Algebra

**Boolean Algebra** is the set of rules dictating how to transform one statement into another.

Another way to show logical equivalence (besides truth tables) is with **Boolean Algebra**.

So far we know:

- De Morgan's Laws
  - $\neg(P \vee Q)$  is logically equivalent to  $\neg P \wedge \neg Q$
  - $\neg(P \wedge Q)$  is logically equivalent to  $\neg P \vee \neg Q$
- Implications are disjunctions
  - $P \rightarrow Q$  is logically equivalent to  $\neg P \vee Q$
- Double negation
  - $\neg\neg P$  is logically equivalent to  $P$

# Boolean Algebra

**Boolean Algebra** is the set of rules dictating how to transform one statement into another.

Another way to show logical equivalence (besides truth tables) is with **Boolean Algebra**.

So far we know:

- De Morgan's Laws
  - $\neg(P \vee Q)$  is logically equivalent to  $\neg P \wedge \neg Q$
  - $\neg(P \wedge Q)$  is logically equivalent to  $\neg P \vee \neg Q$
- Implications are disjunctions
  - $P \rightarrow Q$  is logically equivalent to  $\neg P \vee Q$
- Double negation
  - $\neg\neg P$  is logically equivalent to  $P$

**Practice:** Use Boolean Algebra to show  $\neg(P \rightarrow Q)$  and  $P \wedge \neg Q$  are logically equivalent.

# Boolean Algebra

**Boolean Algebra** is the set of rules dictating how to transform one statement into another.

Another way to show logical equivalence (besides truth tables) is with **Boolean Algebra**.

So far we know:

- De Morgan's Laws
  - $\neg(P \vee Q)$  is logically equivalent to  $\neg P \wedge \neg Q$
  - $\neg(P \wedge Q)$  is logically equivalent to  $\neg P \vee \neg Q$
- Implications are disjunctions
  - $P \rightarrow Q$  is logically equivalent to  $\neg P \vee Q$
- Double negation
  - $\neg\neg P$  is logically equivalent to  $P$

**Practice:** Use Boolean Algebra to show  $\neg((\neg P \wedge Q) \vee \neg(R \vee \neg S))$  and  $(Q \rightarrow P) \wedge (S \rightarrow R)$  are logically equivalent.

# Variables and Statements

Often we want to use variables in our mathematical sentences.

For example, suppose we want to represent the following as a mathematical statement:

If  $n$  is prime, then  $n + 7$  is not prime.



## Variables and Statements

Often we want to use variables in our mathematical sentences.

For example, suppose we want to represent the following as a mathematical statement:

If  $n$  is prime, then  $n + 7$  is not prime.

Hmm... this looks like an implication (if ... then ...), but  $n$  could be anything; it's not limited to one value.

## Variables and Statements

Often we want to use variables in our mathematical sentences.

For example, suppose we want to represent the following as a mathematical statement:

If  $n$  is prime, then  $n + 7$  is not prime.

Hmm... this looks like an implication (if ... then ...), but  $n$  could be anything; it's not limited to one value.

We need a way to symbolically write this statement for different values of  $n$ .

## Variables and Statements

Often we want to use variables in our mathematical sentences.

For example, suppose we want to represent the following as a mathematical statement:

If  $n$  is prime, then  $n + 7$  is not prime.

Hmm... this looks like an implication (if ... then ...), but  $n$  could be anything; it's not limited to one value.

We need a way to symbolically write this statement for different values of  $n$ .

Let's let  $P(n) = n$  is prime

## Variables and Statements

Often we want to use variables in our mathematical sentences.

For example, suppose we want to represent the following as a mathematical statement:

If  $n$  is prime, then  $n + 7$  is not prime.

Hmm... this looks like an implication (if ... then ...), but  $n$  could be anything; it's not limited to one value.

We need a way to symbolically write this statement for different values of  $n$ .

Let's let  $P(n)$  =  $n$  is prime

Now we can represent the statement above as:

$$P(n) \rightarrow \neg P(n + 7)$$

## Variables and Statements

Let  $P(n)$  =  $n$  is prime

Now we can represent the statement If  $n$  is prime, then  $n + 7$  is not prime as  $P(n) \rightarrow \neg P(n + 7)$

## Variables and Statements

Let  $P(n)$  =  $n$  is prime

Now we can represent the statement If  $n$  is prime, then  $n + 7$  is not prime as  $P(n) \rightarrow \neg P(n + 7)$

**Practice:** Write the sentence "Primes greater than 2 are odd" symbolically.

## Variables and Statements

Let  $P(n)$  =  $n$  is prime

Now we can represent the statement If  $n$  is prime, then  $n + 7$  is not prime as  $P(n) \rightarrow \neg P(n + 7)$

Notice that  $P(n) \rightarrow \neg P(n + 7)$  is *not a statement*. We have not specified anything about  $n$ ; it can be any value.

## Variables and Statements

Let  $P(n)$  =  $n$  is prime

Now we can represent the statement If  $n$  is prime, then  $n + 7$  is not prime as  $P(n) \rightarrow \neg P(n + 7)$

Notice that  $P(n) \rightarrow \neg P(n + 7)$  is *not a statement*. We have not specified anything about  $n$ ; it can be any value.

If we plug in a specific value for  $n$ , then we get a statement.



## Variables and Statements

Let  $P(n)$  =  $n$  is prime

Now we can represent the statement If  $n$  is prime, then  $n + 7$  is not prime as  $P(n) \rightarrow \neg P(n + 7)$

Notice that  $P(n) \rightarrow \neg P(n + 7)$  is *not a statement*. We have not specified anything about  $n$ ; it can be any value.

If we plug in a specific value for  $n$ , then we get a statement.

A sentence that contains variables is called a ***predicate***.

# Variables and Statements

Let  $P(n)$  =  $n$  is prime

Now we can represent the statement If  $n$  is prime, then  $n + 7$  is not prime as  $P(n) \rightarrow \neg P(n + 7)$

Notice that  $P(n) \rightarrow \neg P(n + 7)$  is *not a statement*. We have not specified anything about  $n$ ; it can be any value.

If we plug in a specific value for  $n$ , then we get a statement.

A sentence that contains variables is called a ***predicate***.

To make this predicate a statement (something that is always true or false), we need to quantify the variable,  $n$ .

In mathematics we use two quantifiers: existential and universal.

# Quantifiers

# Quantifiers

In mathematics we use two quantifiers: existential and universal.

The ***existential quantifier*** is  $\exists$  and is read “there exists”

Ex.

$$\exists x(x < 0)$$

is read “There exists an  $x$  such that  $x$  is less than 0”

This statement says there exists a number less than 0.

# Quantifiers

In mathematics we use two quantifiers: existential and universal.

The ***universal quantifier*** is  $\forall$  and is read “for all” or “every”

Ex.

$$\forall x(x \geq 0)$$

is read “For all x, x is greater than or equal to 0”

This statement says every number is greater than or equal to 0.

# Quantifiers

The ***existential quantifier*** is  $\exists$  and is read “there exists”

The ***universal quantifier*** is  $\forall$  and is read “for all” or “every”

**Practice:** Write the sentence “All primes greater than 2 are odd” symbolically.

# Quantifiers

The ***existential quantifier*** is  $\exists$  and is read “there is”

The ***universal quantifier*** is  $\forall$  and is read “for all” or “every”

Quantifiers can be combined.

Ex.

$$\forall x \exists y (y < x)$$

is read “for every  $x$  there exists some  $y$  such that  $y$  is less than  $x$ ”

# Quantifiers

The ***existential quantifier*** is  $\exists$  and is read “there is”

The ***universal quantifier*** is  $\forall$  and is read “for all” or “every”

Quantifiers can be combined.

Ex.

$$\forall x \exists y (y < x)$$

is read “for every x there exists some y such that y is less than x”

**Note:** we also need to define the ***domain*** of variables. For example, does every x mean all positive integers, all real numbers, something else? [The ***domain*** of a variable is the set of possible values it can take.]



# Quantifiers

The ***existential quantifier*** is  $\exists$  and is read “there is”

The ***universal quantifier*** is  $\forall$  and is read “for all” or “every”

**Practice:** Once we have defined the domain and quantified variables in a predicate, we have a statement!

Consider this statement:  $\forall x \exists y (y < x)$

Let the domain of  $x, y$  be the ***natural numbers*** (0, 1, 2, ...)

What is the truth value of the statement?

**Note:** we also need to define the ***domain*** of variables. For example, does every  $x$  mean all positive integers, all real numbers, something else? [The ***domain*** of a variable is the set of possible values it can take.]

# Quantifiers

The **existential quantifier** is  $\exists$  and is read “there is”

The **universal quantifier** is  $\forall$  and is read “for all” or “every”

**Practice:** Once we have defined the domain and quantified variables in a predicate, we have a statement!

Consider this statement:  $\exists x \forall y (y \geq x)$

Let the domain of  $x, y$  be the **natural numbers** (0, 1, 2, ...)

What is this statement in English?

What is the truth value of the statement?

**Note:** we also need to define the **domain** of variables. For example, does every  $x$  mean all positive integers, all real numbers, something else? [The **domain** of a variable is the set of possible values it can take.]

## Negation of Quantifiers

The ***existential quantifier*** is  $\exists$  and is read “there exists”

The ***universal quantifier*** is  $\forall$  and is read “for all” or “every”

Quantifiers can be negated:

$\neg \forall x P(x)$  is equivalent to  $\exists x \neg P(x)$

$\neg \exists x P(x)$  is equivalent to  $\forall x \neg P(x)$

## Negation of Quantifiers

The ***existential quantifier*** is  $\exists$  and is read “there is”

The ***universal quantifier*** is  $\forall$  and is read “for all” or “every”

Quantifiers can be negated:

$\neg \forall x P(x)$  is equivalent to  $\exists x \neg P(x)$

$\neg \exists x P(x)$  is equivalent to  $\forall x \neg P(x)$

**Practice:** Rewrite the following statement without negation

$$\neg \exists x \forall y (x \leq y)$$

What is each version in English?

## Back to Implications

In the statement  $P \rightarrow Q$ , we say

- P is the ***hypothesis*** (or antecedent)
- and Q is the ***conclusion*** (or consequent)

## Back to Implications

In the statement  $P \rightarrow Q$ , we say

- P is the ***hypothesis*** (or antecedent)
- and Q is the ***conclusion*** (or consequent)

Recall this statement is false only when P is T and Q is F

$P$	$Q$	$P \rightarrow Q$
T	T	T
T	F	F
F	T	T
F	F	T

## Back to Implications

In the statement  $P \rightarrow Q$ , we say

- P is the ***hypothesis*** (or antecedent)
- and Q is the ***conclusion*** (or consequent)

Recall this statement is false only when P is T and Q is F

$P$	$Q$	$P \rightarrow Q$
T	T	T
T	F	F
F	T	T
F	F	T

We will use this later for proofs. We will assume our hypothesis is true, and show that this implies our conclusion is true.

## Back to Implications

The ***converse*** of an implication  $P \rightarrow Q$  is the implication  $Q \rightarrow P$



## Back to Implications

The ***converse*** of an implication  $P \rightarrow Q$  is the implication  $Q \rightarrow P$

- The converse of an implication is not logically equivalent to the original implication

## Back to Implications

The **converse** of an implication  $P \rightarrow Q$  is the implication  $Q \rightarrow P$

- The converse of an implication is not logically equivalent to the original implication

**Practice:** Come up with an example where  $P \rightarrow Q$  is true and  $Q \rightarrow P$  is false.

## Back to Implications

The ***contrapositive*** of an implication  $P \rightarrow Q$  is the statement  $\neg Q \rightarrow \neg P$

## Back to Implications

The ***contrapositive*** of an implication  $P \rightarrow Q$  is the statement  $\neg Q \rightarrow \neg P$

- The contrapositive of an implication is always logically equivalent to the original implication

## Back to Implications

The ***contrapositive*** of an implication  $P \rightarrow Q$  is the statement  $\neg Q \rightarrow \neg P$

- The contrapositive of an implication is always logically equivalent to the original implication

**Practice:** Show  $P \rightarrow Q$  and  $\neg Q \rightarrow \neg P$  are logically equivalent using truth tables or Boolean algebra.