

Discrete Structures— Proofs: Contradiction and Counter Example

Dr. Ab Mosca (they/them)

Plan for Today

- Proof by contradiction
- Proof by counter example

Warm Up

Direct proof is usually used to prove implications.

To prove $P \rightarrow Q$, we will

1. Assume P is true
2. Deduce that Q must then also be true

Proof by contrapositive

To prove $P \rightarrow Q$, we will

1. Prove the contrapositive ($\neg Q \rightarrow \neg P$)
 1. Assume $\neg Q$ is true
 2. Deduce that $\neg P$ must then also be true

Work with a small group to prove if ab is an even number, then a , or b is even.

Proof by Contradiction

Consider the implication $\neg P \rightarrow Q$

If this implication is true, and Q is false, what does that tell us about $\neg P$?

Proof by Contradiction

Consider the implication $\neg P \rightarrow Q$

If this implication is true, and Q is false, what does that tell us about $\neg P$?

It tells us that $\neg P$ must be false!

P	Q	$P \rightarrow Q$
T	T	T
T	F	F
F	T	T
F	F	T

Proof by Contradiction

Consider the implication $\neg P \rightarrow Q$

If this implication is true, and Q is false, what does that tell us about $\neg P$?

It tells us that $\neg P$ must be false!

P	Q	$P \rightarrow Q$
T	T	T
T	F	F
F	T	T
F	F	T

If $\neg P$ is false, then P must be true.

Proof by Contradiction

Consider the implication $\neg P \rightarrow Q$

If this implication is true, and Q is false, what does that tell us about $\neg P$?

It tells us that $\neg P$ must be false!

P	Q	$P \rightarrow Q$
T	T	T
T	F	F
F	T	T
F	F	T

If $\neg P$ is false, then P must be true.

This gives us another option for proof called ***proof by contradiction***.

Proof by Contradiction

To perform a ***proof by contradiction*** (on P):

1. Assume $\neg P$ is true
2. Show that this assumption leads to a contradiction
3. As a result, the only conclusion is that P is true
(i.e. if it impossible for P to be false, we know it must be true)

Proof: We will show that $\sqrt{2}$ is irrational using proof by contradiction.

Proof by
Contradiction

Proof by Contradiction

Proof: We will show that $\sqrt{2}$ is irrational using proof by contradiction.

Assume $\sqrt{2}$ is rational. This means that we can write it as a fraction, $\frac{a}{b}$, where a and b are integers. We will assume $\frac{a}{b}$ is in lowest terms.

Proof by Contradiction

Proof: We will show that $\sqrt{2}$ is irrational using proof by contradiction.

Assume $\sqrt{2}$ is rational. This means that we can write it as a fraction, $\frac{a}{b}$, where a and b are integers. We will assume $\frac{a}{b}$ is in lowest terms.

Now consider $(\sqrt{2})^2 = \left(\frac{a}{b}\right)^2$

Proof by Contradiction

Proof: We will show that $\sqrt{2}$ is irrational using proof by contradiction.

Assume $\sqrt{2}$ is rational. This means that we can write it as a fraction, $\frac{a}{b}$, where a and b are integers. We will assume $\frac{a}{b}$ is in lowest terms.

Now consider $(\sqrt{2})^2 = \left(\frac{a}{b}\right)^2$

This implies $2 = \frac{a^2}{b^2}$

and $a^2 = 2b^2$, which means a^2 is even and a is even.

Proof by Contradiction

Proof: We will show that $\sqrt{2}$ is irrational using proof by contradiction.

Assume $\sqrt{2}$ is rational. This means that we can write it as a fraction, $\frac{a}{b}$, where a and b are integers. We will assume $\frac{a}{b}$ is in lowest terms.

Now consider $(\sqrt{2})^2 = \left(\frac{a}{b}\right)^2$

This implies $2 = \frac{a^2}{b^2}$

and $a^2 = 2b^2$, which means a^2 is even and a is even.

Therefore, we can rewrite a^2 as $4k^2$ for some integer k .

Proof by Contradiction

Proof: We will show that $\sqrt{2}$ is irrational using proof by contradiction.

Assume $\sqrt{2}$ is rational. This means that we can write it as a fraction, $\frac{a}{b}$, where a and b are integers. We will assume $\frac{a}{b}$ is in lowest terms.

Now consider $(\sqrt{2})^2 = \left(\frac{a}{b}\right)^2$

This implies $2 = \frac{a^2}{b^2}$

and $a^2 = 2b^2$, which means a^2 is even and a is even.

Therefore, we can rewrite a^2 as $4k^2$ for some integer k .

This means $4k^2 = 2b^2$

$b^2 = 2k^2$, which means b^2 is even.

Proof by Contradiction

Proof: We will show that $\sqrt{2}$ is irrational using proof by contradiction.

Assume $\sqrt{2}$ is rational. This means that we can write it as a fraction, $\frac{a}{b}$, where a and b are integers. We will assume $\frac{a}{b}$ is in lowest terms.

Now consider $(\sqrt{2})^2 = \left(\frac{a}{b}\right)^2$

This implies $2 = \frac{a^2}{b^2}$

and $a^2 = 2b^2$, which means a^2 is even and a is even.

Therefore, we can rewrite a^2 as $4k^2$ for some integer k .

This means $4k^2 = 2b^2$

$b^2 = 2k^2$, which means b^2 is even.

However, if a is even and b is even, then $\frac{a}{b}$ is not in lowest terms, which contradicts our original assumption.

Proof by Contradiction

Proof: We will show that $\sqrt{2}$ is irrational using proof by contradiction.

Assume $\sqrt{2}$ is rational. This means that we can write it as a fraction, $\frac{a}{b}$, where a and b are integers. We will assume $\frac{a}{b}$ is in lowest terms.

Now consider $(\sqrt{2})^2 = \left(\frac{a}{b}\right)^2$

This implies $2 = \frac{a^2}{b^2}$

and $a^2 = 2b^2$, which means a^2 is even and a is even.

Therefore, we can rewrite a^2 as $4k^2$ for some integer k .

This means $4k^2 = 2b^2$

$b^2 = 2k^2$, which means b^2 is even.

However, if a is even and b is even, then $\frac{a}{b}$ is not in lowest terms, which contradicts our original assumption.

Thus, $\sqrt{2}$ is irrational. //

Proof by Contradiction

To perform a ***proof by contradiction*** (on P):

1. Assume $\neg P$ is true
2. Show that this assumption leads to a contradiction
3. As a result, the only conclusion is that P is true
(i.e. if it impossible for P to be false, we know it must be true)

Practice: Prove the following...

There are no integers x and y such that $x^2 = 4y + 2$.

Proof by Contradiction

To perform a ***proof by contradiction*** (on P):

1. Assume $\neg P$ is true
2. Show that this assumption leads to a contradiction
3. As a result, the only conclusion is that P is true
(i.e. if it impossible for P to be false, we know it must be true)

Practice: Prove the following...

There are no integers x and y such that x is a prime greater than 5 and $x = 6y + 3$.

Proof by Contradiction

To perform a ***proof by contradiction*** (on P):

1. Assume $\neg P$ is true
2. Show that this assumption leads to a contradiction
3. As a result, the only conclusion is that P is true
(i.e. if it impossible for P to be false, we know it must be true)

Practice: Prove the following...

$\sqrt{3}$ is irrational.

Proof by Contradiction

To perform a ***proof by contradiction*** (on P):

1. Assume $\neg P$ is true
2. Show that this assumption leads to a contradiction
3. As a result, the only conclusion is that P is true
(i.e. if it impossible for P to be false, we know it must be true)

Practice: Prove the following...

For all integers a , b , and c , if $a^2 + b^2 = c^2$, then a or b is even.



Start here on Tuesday

Proof by Contradiction

To perform a ***proof by contradiction*** (on P):

1. Assume $\neg P$ is true
2. Show that this assumption leads to a contradiction
3. As a result, the only conclusion is that P is true
(i.e. if it impossible for P to be false, we know it must be true)

Practice: Prove the following...

Pigeonhole principle: If more than n pigeons fly into n pigeonholes, then at least one pigeonhole will contain at least two pigeons.

Pigeonhole Principle

If more than n pigeons fly into n pigeonholes, then at least one pigeonhole will contain at least two pigeons.

This principle comes up often in proofs. Now that you've proven it, you can use it as justification in other proofs.

Pigeonhole Principle

If more than n pigeons fly into n pigeonholes, then at least one pigeonhole will contain at least two pigeons.

This principle comes up often in proofs. Now that you've proven it, you can use it as justification in other proofs.

This is true of all properties, principles, theorems, etc. you've proven before. They are fair game for future proofs. (Unless the property/principle/theorem/etc. is what you're being asked to prove.)

Proof by Counter Example

Proof by counter example is a useful method for showing something is false.

Proof by Counter Example

Proof by counter example is a useful method for showing something is false.

You CANNOT show a statement of the form $\forall x P(x)$ is true with only one example.

Proof by Counter Example

Proof by counter example is a useful method for showing something is false.

You CANNOT show a statement of the form $\forall x P(x)$ is true with only one example.

But you can prove a statement of the form $\forall x P(x)$ is false by providing a counter example.

Proof by Counter Example

Proof by counter example is a useful method for showing something is false.

You CANNOT show a statement of the form $\forall x P(x)$ is true with only one example.

But you can prove a statement of the form $\forall x P(x)$ is false by providing a counter example.

To perform a ***proof by counter example***:

1. Make sure you're trying to show something is false
2. Find an exception to the statement

Proof by Counter Example

Proof: We will show that the statement for all integers a and b , if a is odd or b is odd, then $a + b$ is odd.

Let $a = 1$, and $b = 3$.

Then $a + b = 1 + 3 = 4$.

4 is not odd; we have found an example where the statement is not true.

Thus, the statement is false.

Proof by Counter Example

To perform a ***proof by counter example***:

1. Make sure you're trying to show something is false
2. Find an exception to the statement

Practice: Prove the following...

For all integers a and b , if ab is a multiple of 6, then a is even and b is a multiple of 3.

Notes about proofs

Sometimes you will need to break a proof into ***cases***.

For example, if you were to prove this:

For any integer n , the number $(n^3 - n)$ is even.

You might look at a case where n is even and a case where n is odd.

Notes about proofs

Sometimes you will need to break a proof into **cases**.

For example, if you were to prove this:

For any integer n , the number $(n^3 - n)$ is even.

You might look at a case where n is even and a case where n is odd.

Practice: Prove the following...

For any integer n , the number $(n^3 - n)$ is even.

Notes about proofs

To prove a biconditional, you must prove *both* implications.

For example, if you were to prove this:

For all integers n , it is the case that n is even if and only if $3n$ is even.

Notes about proofs

To prove a biconditional, you must prove *both* implications.

For example, if you were to prove this:

For all integers n , it is the case that n is even if and only if $3n$ is even.

You would need to prove that for all integers n

(a) If n is even then $3n$ is even
and

(b) If $3n$ is even then n is even

Notes about proofs

To prove a biconditional, you must prove *both* implications.

For example, if you were to prove this:

For all integers n , it is the case that n is even if and only if $3n$ is even.

You would need to prove that for all integers n

(a) If n is even then $3n$ is even
and

(b) If $3n$ is even then n is even

Practice: Prove the following...

For all integers n , it is the case that n is even if and only if $3n$ is even.